

ARTIFACT SHEET

Enter artifact number below. Artifact number is application number + artifact type code (see list below) + sequential letter (A, B, C ...). The first artifact folder for an artifact type receives the letter A, the second B, etc.. Examples: 59123456PA, 59123456PB, 59123456ZA, 59123456ZB

Indicate quantity of a single type of artifact received but not scanned. Create individual artifact folder/box and artifact number for each Artifact Type.

<input type="checkbox"/>	CD(s) containing:	<input type="checkbox"/>
	computer program listing	
	Doc Code: Computer	Artifact Type Code: P
	pages of specification	
	and/or sequence listing	<input type="checkbox"/>
	and/or table	
	Doc Code: Artifact	Artifact Type Code: S
	content unspecified or combined	<input type="checkbox"/>
	Doc Code: Artifact	Artifact Type Code: U

<input type="checkbox"/>	Stapled Set(s) Color Documents or B/W Photographs
	Doc Code: Artifact Artifact Type Code: C

<input type="checkbox"/>	Microfilm(s)
	Doc Code: Artifact Artifact Type Code: F

<input type="checkbox"/>	Video tape(s)
	Doc Code: Artifact Artifact Type Code: V

<input type="checkbox"/>	Model(s)
	Doc Code: Artifact Artifact Type Code: M

<input type="checkbox"/>	Bound Document(s)
	Doc Code: Artifact Artifact Type Code: B

<input type="checkbox"/>	Confidential Information Disclosure Statement or Other Documents marked Proprietary, Trade Secrets, Subject to Protective Order, Material Submitted under MPEP 724.02, etc.
	Doc Code: Artifact Artifact Type Code X

<input type="checkbox"/>	Other, description: _____
	Doc Code: Artifact Artifact Type Code: Z

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 65/24, 65/26, 65/28, 59/68, 59/90	A1	(11) International Publication Number: WO 95/13262
		(43) International Publication Date: 18 May 1995 (18.05.95)

(21) International Application Number: PCT/GB94/02499
(22) International Filing Date: 14 November 1994 (14.11.94)

(30) Priority Data:

9323382.3	12 November 1993 (12.11.93)	GB
9403363.6	22 February 1994 (22.02.94)	GB
9410750.5	27 May 1994 (27.05.94)	GB

(71) Applicant (for all designated States except US): RHONE-POULENC RORER LIMITED [GB/GB]; RPR House, 52 St. Leonards Road, Eastbourne, East Sussex BN21 3YG (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ASTLES, Peter, Charles [GB/GB]; Rhône Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). HARPER, Mark, Francis [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). HARRIS, Neil, Victor [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). McLAY, Iain, McFarlane [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). WALSH, Roger, John, Aitchison [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham

Road South, Dagenham, Essex RM10 7XS (GB). LEWIS, Richard, Alan [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). SMITH, Christopher [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). PORTER, Barry [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB). MCCARTHY, Clive [GB/GB]; Rhône-Poulenc Rorer Limited, Rainham Road South, Dagenham, Essex RM10 7XS (GB).

(74) Agent: CAFFIN, Lee; Rhône-Poulenc Rorer Limited, Patent Dept., Rainham Road South, Dagenham, Essex RM10 7XS (GB).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).

Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: SUBSTITUTED PHENYL COMPOUNDS

(57) Abstract

Compounds of formula (I) are described wherein R¹ is hydrogen, -(lower alkyl)_q(CO₂R⁶ or OH), -CN, -C(R⁷)=NOR⁸, NO₂, -O(lower alkyl)R⁹, -C≡C-R¹⁰, -CR¹¹=C(R¹²)(R¹³), -C(-O)CH₂C(-O)CO₂H, -CO(R¹⁴), alkylthio, alkylsulphanyl, alkylsulphonyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, substituted thiocarbamoyl, sulphamoyl or an optionally substituted nitrogen-containing ring, m, n, o and p are independently zero or 1 and R², R³, R⁴ and R⁵ are various groups; and physiologically acceptable salts, N-oxides and prodrugs thereof. The compounds have endothelin antagonist activity and are useful as pharmaceuticals.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

- 1 -

SUBSTITUTED PHENYL COMPOUNDS

Field of the Invention

10

This invention is directed to substituted phenyl compounds, their preparation, pharmaceutical compositions containing these compounds, and their pharmaceutical use in the treatment of disease states associated with endothelin peptides.

15

Endothelins are a family of peptides, mainly synthesized and released by endothelial cells. The term endothelin (ET) refers to a family of homologous 21-amino acid peptides found in three distinct isoforms: ET-1, ET-2 and ET-3, and in the present specification the term "endothelin" is intended to refer to any or all of the isoforms of endothelin. Each endothelin isopeptide is encoded by a distinct gene with a distinct chromosomal locus for each human gene.

20

Receptor subtypes ET_A and ET_B specific for endothelin have been identified (H. Arai, et al., Nature, 348, 730 (1990) and T. Sakurai et al., Nature, 348, 732 (1990)). ET-1 and ET-2 bind more potently than ET-3 to ET_A, and stimulation of this receptor subtype promotes vasoconstriction. ET-1, ET-2 and ET-3 bind with equal affinity to ET_B receptors, and stimulation of this receptor subtype can evoke vasodilation or promote vasoconstriction.

25

Thus, endothelin is an important potent vasoconstrictor producing long-lasting effects in arteries and veins. Consequently, endothelin causes profound actions on the cardiovascular system in particular the coronary, renal, mesenteric and cerebral circulation. Other biological activities by endothelin are also observed. Thus, disease states associated with a physiologically detrimental excess of endothelin are treatable according to the invention.

30

35

- Intravenous infusion of ET-1 to rats causes a transient hypotensive effect, followed by a sustained increase in blood pressure. Even low doses of endothelin, which alone are without pressor actions, potentiate the effects of other vasoconstrictor agents. Significantly elevated plasma immunoreactive ET-1 levels have been reported in patients with disorders such as myocardial infarction including acute myocardial infarction, coronary heart disease, unstable angina including vasospastic angina, preeclampsia, essential and pulmonary hypertension and congestive heart failure.
- Renal blood vessels are particularly sensitive to the vasoconstrictor effect of ET. It produces a marked reduction in renal blood flow accompanied by reductions in glomerular filtration rate, urine volume and urinary sodium and potassium excretion. Endothelin is also mitogenic for mesangial cells. Thus, endothelin has a role in a number of renal disorders such as acute renal insufficiency and chronic renal insufficiency and cyclosporin induced nephrotoxicity. Furthermore, erythropoietin causes endothelin release and that release plays a role in renal complications and hypertension occurring as side effects in dialysis patients.
- Endothelin induces a proliferative response in vascular smooth muscle cells and this, combined with observations of elevated circulating levels of ET-1 in atherosclerosis, indicates that endothelin contributes to the pathogenesis of this and related diseases. Levels of endothelin are also elevated after angioplasty and is implicated in the high level of restenosis after percutaneous transluminal angioplasty.
- The cerebral vasculature is very sensitive to the pressor actions of the endothelins. A single intrathecal injection of ET-1 in dogs leads to a prolonged constriction of the basilar artery. Hypoxia and ischaemia are potent stimuli for increased release of endothelin by endothelial cells, while the secretion of endogenous vasodilators such as PGI₂ and endothelial derived relaxant factor are reduced. Therefore, endothelin plays an important role in cerebral ischaemia such as stroke and subarachnoid hemorrhage.
- Endothelin is a potent contractor of isolated airway tissue including human bronchus. In addition, endothelin has been shown to induce eicosanoid release, possess mitogenic properties for airway smooth muscle

and has pronounced inflammatory actions. All of these actions confirm an important role for endothelin in pulmonary pathophysiology and in asthma and related conditions.

5 Endothelin levels are elevated during septic shock and other endotoxin induced conditions such as disseminated intravascular coagulation, migraine, gastrointestinal disorders such as ulceration and irritable bowel syndrome, Raynauds disease and haemangioendothelioma.


10 Normal bone remodelling involves the coupling of osteoclast and
osteoblast functions, an imbalance of these events leading to
pathophysiological bone loss. Both cell types produce endothelin and possess
endothelin receptors. Antagonists of selected actions of endothelin would
therefore be useful in the treatment of clinical conditions of bone loss, such as
15 osteoporosis.

Endothelin-1 is produced in the human prostate and endothelin receptors have been identified in this tissue. Since endothelin is a paracrine contractile and proliferative factor in the prostate gland, a role is indicated for endothelin in benign prostatic hyperplasia.

The further actions of endothelin on neurotransmitter release are also observed, indicating a role in certain disorders of the central nervous system.

25 SUMMARY OF THE INVENTION

This invention is directed to compounds useful for inhibiting the production or physiological effects of endothelin in the treatment of a patient suffering from a disease state associated with a physiologically detrimental excess of endothelin. Thus, according to a first aspect of the present invention, we provide a compound of formula I as follows:



wherein

5 R^1 is hydrogen, $-(\text{lower alkyl})_q(\text{CO}_2R^6 \text{ or OH})$, $-\text{CN}$, $-\text{C}(R^7)=\text{NOR}^8$, $-\text{NO}_2$, $-\text{O}(\text{lower alkyl})R^9$, $-\text{C}\equiv\text{C}-R^{10}$, $-\text{CR}^{11}=\text{C}(R^{12})(R^{13})$, $-\text{C}(=\text{O})\text{CH}_2\text{C}(=\text{O})\text{CO}_2\text{H}$, $-\text{CO}(R^{14})$, alkylthio, alkylsulphinyl, alkylsulphonyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, substituted thiocarbamoyl, sulphamoyl or an optionally substituted nitrogen-containing ring;

10 R^2 is aryl lower alkoxy, heteroaryl lower alkoxy, aryl lower alkylthio or heteroaryl lower alkylthio;

15 R^3 is hydroxy, alkoxy, aryloxy, cycloalkyl(lower alkyl)oxy, cycloalkenyl(lower alkyl)oxy, aryl lower alkoxy, heteroaryl lower alkoxy, aryl lower alkylthio, heteroaryl lower alkylthio or aralkynyl;

R^4 is $-\text{Y}-\text{CH}(R^{15})(\text{alkyl or alkenyl})R^{16}$;

R^5 is alkyl, alkenyl or halo;

20 R^6 , R^7 , R^{17} , R^{18} and R^{19} are independently hydrogen or alkyl;

R^8 is hydrogen, aralkyl or $-(\text{lower alkyl})\text{CO}_2R^{17}$;

25 R^9 is $-\text{CN}$, $-\text{CO}_2R^{19}$, $-\text{CH}_2\text{OH}$, or carbamoyl;

R^{10} is $-\text{CO}_2\text{H}$ or carboxyphenyl;

R^{11} is hydrogen, alkyl or aralkyl;

30 R^{12} and R^{13} are independently hydrogen, $-\text{CO}_2R^{18}$, $-\text{CN}$, aryl lower alkyl, heteroaryl lower alkyl or $-\text{NHC}(=\text{O})\text{aryl}$, provided that one of R^{12} and R^{13} is $-\text{CO}_2\text{H}$;

35 R^{14} is hydrogen, alkyl, $-(\text{lower alkyl})\text{carboxy}$, aralkenyl, heteroaralkenyl or carboxy;

R^{15} is aryl or heteroaryl;

R¹⁶ is carboxy or acid isostere;

Y is oxygen or carbonyl; and

5

m, n, o, p and q are independently zero or 1,

with the provisos that (i) when o is zero then m and n are both 1 (ii) when o is 1 then at least one of m and n is 1 (iii) when o is zero then R¹ cannot represent
10 hydrogen,
or a pharmaceutically acceptable salt, N-oxide or prodrug thereof.

DETAILED DESCRIPTION OF THE INVENTION

15

As used above, and throughout the description of the invention, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

20 Definitions

"Patient" includes both human and other mammals.

"Pharmaceutically acceptable salt" means a salt form of the parent
25 compound of formula I which is relatively innocuous to a patient when used in therapeutic doses so that the beneficial pharmaceutical properties of the parent compound of formula I are not vitiated by side-effects ascribable to a counter ion of that salt form. Pharmaceutically acceptable salt also includes a zwitterion or internal salt of the compound of formula I.

30

"Alkyl" means an aliphatic hydrocarbon group which may be straight or branched having about 1 to about 6 carbon atoms in the chain. Preferred alkyl groups have 1 to about 4 carbon atoms in the chain, especially 1 to about 2 carbon atoms. Branched means that one or more lower alkyl groups such as
35 methyl, ethyl or propyl are attached to a linear alkyl chain. "Lower alkyl" means about 1 to about 4 carbon atoms in the chain which may be straight or branched. The alkyl group may be independently substituted by one or more

halo, cycloalkyl or cycloalkenyl. Exemplary alkyl groups include methyl, fluoromethyl, trifluoromethyl, cyclohexylmethyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *n*-pentyl and pentafluoroethyl.

5 "Alkenyl" means an aliphatic hydrocarbon group containing a carbon-carbon double bond and which may be straight or branched having about 2 to about 6 carbon atoms in the chain. Preferred alkenyl groups have 2 to about 3 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl or ethyl are attached to a linear alkenyl chain. "Lower
10 alkenyl" means about 2 to about 4 carbon atoms in the chain which may be straight or branched. The alkenyl group may be independently substituted by one or more halo, cycloalkyl or cycloalkenyl. Exemplary alkenyl groups include ethenyl, propenyl, *n*-butenyl, *i*-butenyl, 3-methylbut-2-enyl, *n*-pentenyl, hexenyl or cyclopentylethenyl.

15 "Alkynyl" means an aliphatic hydrocarbon group containing a carbon-carbon triple bond and which may be straight or branched having about 2 to about 6 carbon atoms in the chain. Preferred alkynyl groups have 2 to about 3 carbon atoms in the chain. Branched means that one or more lower alkyl
20 groups such as methyl are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to about 4 carbon atoms. An exemplary alkynyl groups is ethynyl.

"Alkylenedioxy" means an -O-alkyl-O- group in which the alkyl group is
25 as previously described. Exemplary alkylenedioxy groups include methylenedioxy and ethylenedioxy.

"Cycloalkyl" means a saturated mono- or multicyclic ring system of about 3 to about 10 carbon atoms. The cycloalkyl group may be substituted by one or
30 more halo or alkyl. Preferred monocyclic cycloalkyl rings include cyclopentyl, fluorocyclopentyl and cyclohexyl; more preferred is cyclohexyl. Exemplary multicyclic cycloalkyl rings include 1-decalin and norbornyl.

"Cycloalkenyl" means a non-aromatic monocyclic or multicyclic ring
35 system containing a carbon-carbon double bond and having about 3 to about 10 carbon atoms. Exemplary monocyclic cycloalkenyl rings include cyclopentenyl or cyclohexenyl; preferred is cyclohexenyl. An exemplary

multicyclic cycloalkenyl ring is norbornylenyl. The cycloalkenyl group may be independently substituted by one or more halo or alkyl.


"Aryl" means aromatic carbocyclic radical containing about 6 to about 10 carbon atoms. Exemplary aryl include phenyl or naphthyl, or phenyl or naphthyl substituted with one or more aryl group substituents which may be the same or different, where "aryl group substituent" includes alkyl, preferably lower alkyl, halo for example chloro, bromo or fluoro, CF_3 , amino, (lower alkyl) $_q\text{CO}_2\text{R}^{11}$, carbamoyl, thiocarbamoyl, substituted carbamoyl, substituted thiocarbamoyl, nitro, cyano, alkoxy, preferably lower alkoxy, hydroxy and alkylenedioxy, where q, R^{11} , alkoxy, alkyl, alkylenedioxy, carbamoyl, thiocarbamoyl, substituted carbamoyl and substituted thiocarbamoyl, are as defined herein.

"Heteroaryl" means about a 5- to about a 10- membered aromatic monocyclic or multicyclic hydrocarbon ring system in which one or more of the carbon atoms in the ring system is/are element(s) other than carbon, for example nitrogen, oxygen or sulfur. The "heteroaryl" may also be substituted by one or more aryl group substituents, or aryl, heteroaryl, aralkyl or hydroxyalkyl. Exemplary heteroaryl groups include pyrazinyl, pyrazolyl, tetrazolyl, furanyl, (2- or 3-)thienyl, (2-, 3- or 4-)pyridyl, imidazolyl, pyrimidinyl, isoxazolyl, thiazolyl, isothiazolyl, quinolinyl, indolyl, isoquinolinyl, oxazolyl and 1,2,4-oxadiazolyl. Preferred heteroaryl groups include pyridyl, isothiazolyl and thienyl.

"Optionally substituted nitrogen-containing ring" means a 5- or 6-membered ring containing at least one nitrogen atom and optionally containing one or more additional heteroatoms selected from nitrogen, oxygen and sulphur in addition to the carbon atom(s) present. The ring system may be unsaturated or partially saturated and may optionally be substituted, for example by any of the groups Y^1 , CO_2R^{20} or R as defined hereinafter. Exemplary nitrogen-containing ring systems include tetrazolyl, substituted tetrazolyl, 1,2,4-oxadiazolyl, substituted isoxazolyl, isothiazolyl, substituted thiazolyl, pyrazolyl, substituted pyrazolyl, pyridyl, oxazolyl, substituted oxazolyl and dihydrooxazolyl.


"Prodrug" means a compound, for example an ester, which is convertible in vivo by metabolic means (e.g. by hydrolysis) to a compound of formula I.

5 "Substituted pyrazolyl" means a group of the following formula,



wherein Y¹ is hydroxymethyl, carbamoyl, substituted carbamoyl, thiocarbamoyl, substituted thiocarbamoyl or -(lower alkyl)_kCO₂R²⁰, wherein k is 0 or 1 and R²⁰ is hydrogen and alkyl, and alkyl, carbamoyl, substituted carbamoyl, thiocarbamoyl and substituted thiocarbamoyl are as defined herein. The substituted pyrazolyl may also be C- or N-substituted by lower alkyl, aryl, heteroaryl, aralkyl or -(lower alkyl)_kCO₂R²⁰. The substituted pyrazolyl group is preferably bonded at its 3-position to the phenyl depicted in formula I. The substituted pyrazolyl wherein R¹⁷ is hydrogen is preferred. The Y¹ moiety is preferably substituted at the 5-position. Exemplary substituted pyrazolyl groups include 5-(carboxy)pyrazol-3-yl, N-methyl-5-(carboxymethyl)pyrazol-3-yl, 5-(hydroxymethyl)pyrazol-3-yl, N-phenyl-5-(carboxy)pyrazol-3-yl, N-(1- or 2-)-benzyl-5-(carboxy)pyrazol-3-yl and 2-(2-pyridyl)pyrazol-3-yl.


"Substituted isoxazolyl" means a group of the following formula,



25 wherein R²⁰ is hydrogen and alkyl. The substituted isoxazolyl wherein R²⁰ is hydrogen is preferred. The substituted isoxazolyl is preferably bonded at its 3-position to the phenyl depicted in formula I. The carboxy moiety is preferably substituted at the 5-position.


30

"Substituted oxazolyl" means a group of the following formula,



wherein R^{20} is hydrogen and alkyl. The substituted oxazolyl wherein R^{20} is hydrogen is preferred. The oxazolyl group is preferably bonded at its 2-position to the phenyl depicted in formula I. The carboxy moiety is preferably substituted at the 4-position.

"Substituted thiazolyl" means a group of the following formula,




10

wherein R^{20} is hydrogen and alkyl. The substituted thiazolyl wherein R^{20} is hydrogen is preferred. The thiazolyl group is preferably bonded at its 2-position to the phenyl group depicted in formula I. The moiety $-CO_2R^{20}$ is preferably in the 4-position of the thiazolyl ring.

15

"Substituted tetrazolyl" means a group of the following formulae

20



where R is alkyl, hydroxyalkyl or (lower alkyl) CO_2R^{11} , wherein alkyl and R^{11} are as defined herein.

25

"Aralkyl" means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls are aryl lower alkyls. Exemplary aralkyl groups include benzyl and phenethyl.

30

"Aralkenyl" means an aryl-alkenyl- group in which the aryl and alkenyl are as previously described. Preferred aralkenyls are aryl lower alkenyls. An exemplary aralkenyl group is styryl.

5 "Aralkynyl" means an aryl-alkynyl- group in which the aryl and alkynyl are as previously described. Preferred aralkynyls are aryl lower alkynyls. An exemplary aralkenyl group is phenylethynyl.

10 "Heteroaralkyl" means an heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls are heteroaryl lower alkyls. Exemplary heteroaralkyl groups include pyrid(2- or 3-)ylmethyl, pyrid(2- or 3-)ylethyl, thienylethyl, thienylmethyl, indol-3-ylmethyl or furylmethyl.

15 "Heteroaralkenyl" means an heteroaryl-alkenyl- group in which the heteroaryl and alkenyl are as previously described. Preferred heteroaralkenyls are heteroaryl lower alkenyls. An exemplary heteroaralkenyl group is 3-(2-pyridyl)prop-2-enyl.

20 "Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Preferred alkoxy groups are lower alkoxy groups having 1 to about 3 carbon atoms. Exemplary alkoxy groups include methoxy, ethoxy, *n*-propoxy, *i*-propoxy, *n*-butoxy, *t*-butoxy and hexyloxy.

25 "Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Exemplary aryloxy groups include phenoxy and naphthoxy.

30 "Aralkyloxy" means an aralkyl-O- group in which the aralkyl groups is as previously described. Preferred aralkyloxys are aryl lower alkoxys. Exemplary aralkyloxy groups include benzyloxy, phenylethoxy, phenylpropyloxy, (1- or 2-naphthalene)ethoxy and (o-tolyl)ethoxy; preferred are 1-phenylethoxy and 1-(o-tolyl)ethoxy.

35 "Heteroaralkyloxy" means an heteroaryl-alkyl-O- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyloxys are heteroaryl lower alkoxys. Exemplary heteroaralkyloxy groups include

pyrid(2- or 3-)ylethoxy, pyrid(2- or 3-)ylmethoxy, thienylmethoxy and thienylethoxy.

"Alkylthio" means alkyl-S- in which the alkyl is as previously described.

- 5 Preferred alkylthios are lower alkylthios. An exemplary alkylthio group is methylthio.

- 10 "Alkylsulphinyl means alkyl-SO- in which the alkyl is as previously described. Preferred alkylsulphinyls are lower alkylsulphinyls. An exemplary alkylsulphinyl group is methylsulphinyl.

"Alkoxycarbonyl" means an alkyl-O-CO- group. Exemplary alkoxycarbonyl groups include methoxy- and ethoxycarbonyl.

- 15 "Halo" means fluoro, chloro, bromo, or iodo. Preferred are fluoro, chloro or bromo, and more preferred are fluoro or chloro.

"Carbamoyl" means -CONH₂.

- 20 "Substituted Carbamoyl" means -CONY²Y³ in which Y² and Y³ are independently hydrogen, alkyl, cyano(lower alkyl), arylalkyl, heteroaralkyl, carboxy(lower alkyl), carboxy(aryl substituted lower alkyl), carboxy(carboxy substituted lower alkyl), carboxy(hydroxy substituted lower alkyl), carboxy(heteroaryl substituted lower alkyl), carbamoyl(lower alkyl),
25 alkoxycarbonyl(lower alkyl) or alkoxycarbonyl(aryl substituted lower alkyl), provided that only one of Y² and Y³ may be hydrogen and when one of Y² and Y³ is carboxy(lower alkyl), carboxy(aryl substituted lower alkyl), carbamoyl(lower alkyl), alkoxycarbonyl(lower alkyl) or alkoxycarbonyl(aryl substituted lower alkyl) then the other of Y² and Y³ is hydrogen or alkyl.
30 Preferred for Y² and Y³ are independently hydrogen, alkyl, cyano(lower alkyl), arylalkyl, heteroaralkyl, carboxy(lower alkyl), carboxy(aryl substituted lower alkyl) and carbamoyl(lower alkyl).

"Thiocarbamoyl" means -CSNH₂.

35

"Substituted thiocarbamoyl" means -CSNY²Y³ in which Y² and Y³ are as defined above.

"Alkoxycarbonyl(lower alkyl)" means alkoxy-CO-lower alkyl in which the alkoxy and lower alkyl are as previously described.

5 "Carboxy(aryl substituted lower alkyl)" means a lower alkyl group substituted by an aryl moiety and a carboxy moiety, wherein the alkyl and aryl moieties are as defined herein.

10 "Alkoxycarbonyl(aryl substituted lower alkyl)" means a lower alkyl group substituted by an aryl moiety and an alkoxy moiety, wherein the alkyl, aryl and alkoxy moieties are as defined herein.

"Aryl lower alkylthio" means aryl-lower alkyl-S- in which the aryl and lower alkyl are as previously described.

15 "Heteroaryl lower alkylthio" means heteroaryl-lower alkyl-S- in which the heteroaryl and lower alkyl are as previously described.

20 "Acid isostere" means a group which is significantly ionised at physiological pH. Examples of suitable acid isosteres include sulfo, phosphono, alkylsulphonylcarbamoyl, tetrazolyl, arylsulphonylcarbamoyl or heteroarylsulphonylcarbamoyl.

Preferred Embodiments

25 A compound of formula I is preferred for use in treating a disease state associated with a physiologically detrimental excess of endothelin.

30 Disease states associated with pathological conditions that are modulated by inhibiting endothelin are also preferably treated with a compound of formula I.

According to the compound aspect of the invention, preferred compounds are described by formula I wherein

35 R^1 is hydrogen, $-(\text{lower alkyl})_q(\text{CO}_2\text{H or OH})$, $-\text{CN}$, $-\text{C}(\text{R}^7)=\text{NOR}^8$, $-\text{NO}_2$, $-\text{O}(\text{lower alkyl})\text{R}^9$, $-\text{C}\equiv\text{C}-\text{R}^{10}$, $-\text{CR}^{11}=\text{C}(\text{R}^{12})(\text{R}^{13})$, $-\text{C}(=\text{O})\text{CH}_2\text{C}(=\text{O})\text{CO}_2\text{H}$,

-CO(R¹⁴), alkylthio, alkylsulphinyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted tetrazolyl, 1,2,4-oxadiazolyl, substituted isoxazolyl, pyrazolyl, substituted pyrazolyl, pyridyl, isothiazolyl, oxazolyl, substituted oxazolyl or dihydrooxazolyl;

5

R² is aryl lower alkoxy or heteroaryl lower alkoxy;

R⁴ is -O-CH(aryl)(alkyl)R¹⁶;

10

R⁵ is halo;

R⁸ is hydrogen or -(lower alkyl)CO₂H;

R⁹ is -CN, -CO₂H or carbamoyl;

15

R¹⁰ is -CO₂H;

R¹² and R¹³ are independently hydrogen, -CO₂H, -CN or -NHC(=O)aryl, provided that one of R¹² and R¹³ is -CO₂H;

20

R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, heteroarylalkenyl or carboxy; and

m, o and p are 1 and n is zero.

25

According to another compound aspect of the invention, preferred compounds are described by formula I wherein

R¹ is hydrogen, -(lower alkyl)_q(CO₂H or OH), -CN, -C(R⁷)=NOR⁸, -NO₂,
 30 -O(lower alkyl)R⁹, -C≡C-R¹⁰, -CR¹¹=C(R¹²)(R¹³), -C(=O)CH₂C(=O)CO₂H,
 -CO(R¹⁴), alkylthio, alkylsulphinyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted tetrazolyl, 1,2,4-oxadiazolyl, substituted isoxazolyl, pyrazolyl, substituted pyrazolyl, pyridyl, oxazolyl, substituted oxazolyl or dihydrooxazolyl;

35

R² is aryl lower alkoxy or heteroaryl lower alkoxy;

R⁴ is -O-CH(aryl)(alkyl)R¹⁶;

R⁸ is hydrogen, aralkyl or -(lower alkyl)CO₂H

5 R⁹ is -CN, -CO₂H, -CH₂OH, or carbamoyl;

R¹⁰ is -CO₂H;

10 R¹² and R¹³ are independently hydrogen, -CO₂H, -CN, or
-NHC(=O)aryl, provided that one of R¹² and R¹³ is -CO₂H;

R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, heteroarylalkenyl or
carboxy; and

15 m and o are 1 and n and p are zero.

According to yet another compound aspect of the invention, preferred
compounds are described by formula I wherein

20 R¹ is -(lower alkyl)_q(CO₂H or OH), -CN, -C(R⁷)=NOR⁸, -NO₂, -O(lower
alkyl)R⁹, -C≡C-R¹⁰, -CR¹¹=C(R¹²)(R¹³), -C(=O)CH₂C(=O)CO₂H, -CO(R¹⁴),
carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted
tetrazolyl, substituted isoxazolyl, substituted pyrazolyl or substituted oxazolyl;

25 R² is aryl lower alkoxy or heteroaryl lower alkoxy;

R³ is hydroxy, alkoxy, aryl lower alkoxy, heteroaryl lower alkoxy or
aralkynyl;

30 R⁸ is hydrogen, aralkyl or -(lower alkyl)CO₂H;

R⁹ is -CN, -CO₂H, -CH₂OH, or carbamoyl;

R¹⁰ is -CO₂H;

R¹² and R¹³ are independently hydrogen, -CO₂H, -CN, aryl lower alkyl, heteroaryl lower alkyl or -NHC(=O)aryl, provided that one of R¹² and R¹³ is -CO₂H;

5 R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, heteroarylalkenyl or carboxy;

m and n are 1; and

10 o and p are zero.

More preferred compounds according to the present invention where o is 1 include those wherein R¹ is -CO₂H, -C≡CCO₂H, -C(R¹¹)=CHCO₂H, -(CH₂)₂CO₂H, -O(lower alkyl)CO₂H, -C(=O)NH(lower alkyl)CO₂H, (carboxy)pyrazolyl, (carboxy)(lower alkyl)pyrazolyl, (alkoxycarbonyl)pyrazolyl, -CN, -NO₂, (carboxy)oxazolyl, (carboxy)isoxazolyl, pyrazolyl, formyl, oxazolyl, 1,2,4-oxadiazolyl, acetyl, carboxymethoxyimino, dihydrooxazolyl, pyridyl, carboxamide, thiocarboxamide, pyridylpyrazolyl, cyano(lower alkyl)carboxamide, carboxamidopyrazolyl, isothiazolyl and oxime, wherein R¹¹ is hydrogen or lower alkyl. Further preferred compounds where o is 1 are those wherein R¹ is CN, NO₂, substituted pyrazolyl [e.g. (carboxy)pyrazolyl, (carboxy)(lower alkyl)pyrazolyl, (alkoxycarbonyl)pyrazolyl, pyridylpyrazolyl or carboxamidopyrazolyl], substituted oxazolyl [e.g. (carboxy)oxazolyl] and substituted isoxazolyl [e.g. (carboxy)isoxazolyl], especially where R¹ is CN.

25 More preferred compounds according to the present invention where o is zero include those wherein R¹ is -CO₂H, -C≡CCO₂H, -C(R¹¹)=CHCO₂H, -C(=O)NH(lower alkyl)CO₂H, (carboxy)pyrazolyl, (carboxy)(lower alkyl)pyrazolyl and (carboxy)oxazolyl, wherein R¹¹ is hydrogen or lower alkyl.

30 Further preferred compounds where o is zero are those wherein R¹ is -C(=O)NH(lower alkyl)CO₂H, (carboxy)pyrazolyl and (carboxy)(lower alkyl)pyrazolyl.

35 According to a further compound aspect of the invention, preferred compounds where o is zero are those wherein R² and R³ are substituted on the phenyl moiety at the (2- and 4-) positions relative to R¹.

According to a further compound aspect of the invention, preferred compounds where o is 1 are those wherein R² or R³ is substituted on the phenyl moiety at the 4- position relative to R¹ and R⁴ is substituted on the phenyl moiety at the 2-position relative to R¹; more preferred R² is substituted on the phenyl moiety at the 4- position relative to R¹.

According to a further compound aspect of the invention, preferred compounds where p is 1 are those wherein R⁵ is substituted on the phenyl moiety at the ortho position relative to R¹.

According to another compound aspect of the invention, preferred compounds are those wherein R² and R³ are independently aryl lower alkoxy or heteroaryl lower alkoxy, particularly where R² and R³ are independently heteroaryl lower alkoxy. When o is 1 only one of R² and R³ is preferably present.

According to another compound aspect of the invention, preferred compounds where o is 1 are those wherein R¹⁵ is aryl; more preferably phenyl or particularly phenyl substituted at the ortho position relative to the attachment of the phenyl group to the rest of the R⁴ moiety, and is optionally further substituted.

According to another compound aspect of the invention, preferred compounds where R¹⁵ is a substituted phenyl include those wherein the phenyl moiety is substituted by one or more (e.g. 1, 2 or 3) substituents selected from lower alkyl (e.g. methyl), halo (e.g. chloro), CN, lower alkoxy (e.g. methoxy) or CF₃.

According to the compound aspect of the invention, preferred compounds where o is zero are those wherein R² is 1-(aryl)ethoxy, more preferred 1-(o-tolyl)ethoxy, and R³ is benzyloxy, (3-thienyl)methoxy, (3-pyridyl)methoxy and (4-isothiazolyl)methoxy.

According to the compound aspect of the invention, preferred compounds where o is 1 are those wherein R² is heteroaryl lower alkoxy, more particularly heteroarylmethoxy. Exemplary heteroarylmethoxy groups include

pyridylmethoxy (e.g. 3-pyridylmethoxy), thienylmethoxy (e.g. 3-thienylmethoxy) and isothiazolylmethoxy (e.g. 4-isothiazolylmethoxy).

According to another compound aspect of the invention, preferred
5 compounds where o is 1 are those wherein Y is oxygen.


According to another compound aspect of the invention, preferred compounds are those wherein o is 1 and n is zero.

10 According to another compound aspect of the invention, preferred compounds where o is 1 are those wherein R^4 is $-Y-CH(R^{15})(C_{1-3}alkyl)R^{16}$ in which Y, R^{15} and R^{16} are as defined hereinbefore, especially where R^{16} is carboxy.

15 It is to be understood that the present invention is intended to cover all combinations of particular and preferred groupings as defined herein.

A particular group of compounds of the present invention are compounds of formula Ia as follows:

20




wherein R^1 , R^2 and R^4 are as defined previously.

25 Compounds of formula Ia in which R^1 represents CN, NO_2 , substituted pyrazolyl, substituted oxazolyl, substituted isoxazolyl, and especially CN, are preferred.

30 Compounds of formula Ia in which R^2 represents heteroaryl lower alkoxy, more particularly a heteroarylmethoxy group such as pyridylmethoxy (e.g. 3-pyridylmethoxy), thienylmethoxy (e.g. 3-thienylmethoxy) and isothiazolylmethoxy (e.g. 4-isothiazolylmethoxy), are also preferred.

Compounds of formula Ia in which R^4 represents $-OCH(R^{15})(alkyl)R^{16}$, more particularly where R^{15} is aryl and R^{16} is carboxy, are also preferred. Within $-OCH(R^{15})(alkyl)R^{16}$, R^{15} is preferably phenyl substituted in the ortho position relative to the attachment of the phenyl group to the rest of the R^4 moiety by a lower alkyl (e.g. methyl) or chloro substituent and is optionally further substituted by one or more halo, CF_3 , lower alkyl, CN or lower alkoxy groups. Within $-OCH(R^{15})(alkyl)R^{16}$, the alkyl portion is preferably C_1 -3alkylene, more preferably $-CH_2CH_2-$.

A preferred group of compounds of the present invention are compounds of formula Ib:



wherein R^2 and R^{15} are as defined previously. Particularly preferred are compounds of formula Ib wherein R^2 is heteroarylmethoxy and R^{15} is aryl (e.g. phenyl substituted in the ortho position by lower alkyl such as methyl or chloro, and is optionally further substituted by one or more, e.g. 1, 2 or 3, substituents selected from halo, lower alkyl, CN, CF_3 and lower alkoxy).

Preferred compounds of formulae Ia and Ib above are those wherein the chiral centre associated with the carbon atom α to the oxygen atom within the R^4 group has the (R) configuration.

Compounds of formula Ib wherein R^2 is pyridylmethoxy (e.g. 3-pyridylmethoxy), thienylmethoxy (e.g. 3-thienylmethoxy) or isothiazolylmethoxy (e.g. 4-isothiazolylmethoxy) are particularly preferred.

A further particular group of compounds of the invention are compounds of formula I wherein o is 1, R^4 is $-Y-CH(R^{15})(alkyl \text{ or } alkenyl)R^{16}$ in which Y, R^{15} and R^{16} are as defined in formula I and the groups R^1 , $(R^2)_m$, $(R^3)_n$ and $(R^5)_p$ comprise phenyl substituents wherein the central phenyl ring is

substituted by one or more substituents selected from alkoxy, cycloalkyl(lower alkyl)oxy, arylalkoxy or heteroarylalkoxy and is optionally further substituted by one or more substituents selected from halogen atoms and alkyl, hydroxy, alkanoyl, cyano, carboxy, carbamoyl, alkylcarbamoyl, dialkylcarbamoyl, 5 sulphamoyl, alkylsulphamoyl, dialkylsulphamoyl and pyrazolyl groups.

Another particular group of compounds of the invention are compounds of formula I wherein m and n are 1, o is zero, R¹ is -(lower alkyl)_qCO₂H, CH₂OH, CN, -C(R⁷)=NOH, NO₂, -O(lower alkyl)R⁹, -CH≡C-R¹⁰, 10 -CH=C(R¹²)(R¹³), -C(=O)NH(lower alkyl)CO₂H, -C(=O)CH₂C(=O)CO₂H, -CO(R¹⁴), tetrazolyl, substituted tetrazolyl, substituted isoxazolyl or substituted pyrazolyl, R² is aryl lower alkoxy or heteroaryl lower alkoxy, R³ is hydroxy, alkoxy, phenoxy, cycloalkyl(lower alkyl)oxy, cycloalkenyl(lower alkyl)oxy, aryl lower alkoxy or heteroaryl lower alkoxy, R⁵ is alkyl or alkenyl, R⁷ is hydrogen or alkyl, R⁹ is CN, CO₂H, CH₂OH or CONH₂, R¹⁰ is CO₂H or carboxyphenyl, 15 R¹² and R¹³ are independently hydrogen, CO₂H, CN, aryl lower alkyl, heteroaryl lower alkyl or NHCOaryl provided that one of R¹² and R¹³ is CO₂H, R¹⁴ is hydrogen, alkyl or carboxy lower alkyl, and p and q are independently zero or 1.

20

Particular compounds for use according to the invention are selected from the following:

- A 2-Benzyloxy-4-(4-chlorobenzyloxy)benzoic acid;
- 25 B 2-Benzyloxy-4-(3-phenylpropyloxy)benzoic acid;
- C 2,4-Di-(4-chlorobenzyloxy)benzoic acid;
- D 2-Benzyloxy-4-(2-(3-indolyl)ethoxy)benzoic acid;
- E 2-Hydroxy-4-benzyloxybenzoic acid;
- F 2-(3-Phenylpropyloxy)-4-benzyloxybenzoic acid;
- 30 G 2-Benzyloxy-4-(2-naphthylmethoxy)benzoic acid;
- H 2-Benzyloxy-4-(1-naphthylmethoxy)benzoic acid;
- I 2-Benzyloxy-4-(3,4-methylenedioxybenzyloxy)benzoic acid;
- J 2-(2-Pyridylmethoxy)-4-benzyloxybenzoic acid;
- K 2-(2-Phenylethoxy)-4-benzyloxybenzoic acid;
- 35 L 2-Cyclohexylmethoxy-4-benzyloxybenzoic acid;
- M 2-(4-Chlorobenzyloxy)-4-benzyloxybenzoic acid;
- N 2-Benzyloxy-4-(2-phenylethoxy)benzoic acid;

- O 2-Benzyloxy-4-(4-isopropylbenzyloxy)benzoic acid;
P 2-Benzyloxy-4-(4-nitrobenzyloxy)benzoic acid;
Q 2-Benzyloxy-4-(4-fluorobenzyloxy)benzoic acid;
R 4-Benzyloxy-2-(methylpropoxy)benzoic acid;
5 S 2-(4-Pyridylmethoxy)-4-benzyloxybenzoic acid;
T 2-Benzyloxy-4-(3,4-dichlorobenzyloxy)benzoic acid;
U 2-Benzyloxy-4-(4-methoxybenzyloxy)benzoic acid;
V 2-Benzyloxy-4-(3-methoxybenzyloxy)benzoic acid;
W 2,4-Dibenzyloxybenzoic acid;
10 X (E)-3-(2,4-Dibenzyloxyphenyl)prop-2-enoic acid;
Y 3-(2,4-Dibenzoyloxyphenyl)prop-2-ynoic acid;
Z N-(2,4-Dibenzyloxybenzoyl)glycine;
AA N-(4-Benzyloxy-2-(1-o-tolylethoxy)benzoyl)glycine;
AB 3-(2,4-Dibenzyloxyphenyl)-pyrazole-5-carboxylic acid;
15 AC 2-Benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid;
AD 1-Benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid;
AE 5-(2,4-Dibenzyloxyphenyl)-2H-tetrazole;
AF (Z)-2-Benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid;
AG (E)-2-Benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid;
20 AH E-2-Benzyl-3-(2,4-dibenzyloxyphenyl)acrylic acid;
AI (E)-3-(2,4-Dibenzyloxyphenyl)-2-cyanoacrylic acid;
AJ 3-(2,4-Dibenzyloxyphenyl)isoxazole-5-carboxylic acid;
AK 4-(2,4-Dibenzyloxyphenyl)-2,4-dioxobutanoic acid;
AL 2,4-Dibenzyloxybenzaldehyde oxime;
25 AM 2,4-Dibenzyloxybenzaldehyde;
AN 2,4-Dibenzyloxybenzyl alcohol;
AO 2,4-Dibenzyloxybenzonitrile;
AP 2,4-Dibenzyloxynitrobenzene;
AQ 2,4-Dibenzyloxyacetophenone;
30 AR 2,4-Dibenzyloxyphenol;
AS 2,4-Dibenzyloxyacetophenone oxime;
AT 2-Benzyloxy-4-phenylethynylbenzoic acid;
AU 2-(2,4-Dibenzyloxyphenoxy)ethanol hemihydrate;
AV 2-(2,4-Dibenzyloxyphenoxy)acetamide;
35 AW (RS)-5-(3-Benzyloxyphenyl)-5-oxo-4-phenylpentanoic acid;
AX (RS)-5-(3-Benzyloxyphenyl)-4-(2-chlorophenyl)-5-oxopentanoic acid;
AY (RS)-5-(3-Benzyloxyphenyl)-4-(2-methoxyphenyl)-5-oxopentanoic acid;

- AZ (RS)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
BA (RS)-5-(3-Benzoyloxyphenyl)-4-(2-ethylphenyl)-5-oxopentanoic acid;
BB (RS)-5-(3-Benzoyloxyphenyl)-4-(2-chloro-6-fluorophenyl)-5-oxopentanoic acid;
5 BC (RS)-5-(3-Benzoyloxyphenyl)-4-(2,6-dichlorophenyl)-5-oxopentanoic acid;
BD (RS)-5-(3-Benzoyloxyphenyl)-4-(4-methoxyphenyl)-5-oxopentanoic acid;
BE (RS)-5-(3-benzoyloxyphenyl)-4-(3,4-dichlorophenyl)-5-oxopentanoic acid;
10 BF (RS)-5-(3-Benzoyloxyphenyl)-4-(4-chlorophenyl)-5-oxopentanoic acid;
BG (RS)-5-(3-Benzoyloxyphenyl)-4-(4-methylphenyl)-5-oxopentanoic acid;
BH (RS)-5-(3-Benzoyloxyphenyl)-4-(3-chlorophenyl)-5-oxopentanoic acid;
BI (RS)-5-(3-Benzoyloxyphenyl)-4-(3-methoxyphenyl)-5-oxopentanoic acid;
BJ (RS)-5-(4-Benzoyloxyphenyl)-5-oxo-4-phenylpentanoic acid;
15 BK (RS)-4-(3-Benzoyloxyphenyl)-4-oxo-3-phenylbutanoic acid;
BL (RS)-6-(3-Benzoyloxyphenyl)-6-oxo-5-phenylhexanoic acid;
BM (RS)-7-(3-Benzoyloxyphenyl)-7-oxo-6-phenylheptanoic acid;
BN (RS)-6-(3-Benzoyloxyphenyl)-6-oxo-5-phenylhex-2-enoic acid;
BO (RS)-5-(3-Benzoyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoic acid;
20 BP (3RS,4RS)-5-(3-Benzoyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoic acid;
BQ (2RS,4RS)-5-(3-Benzoyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoic acid;
BR (RS)-5-(3-Methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
25 BS (RS)-5-[3-(2-Methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid;
BT (RS)-5-[3-(3,4-Methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid;
BU (RS)-4-(2-Methylphenyl)-5-[3-(2-methylpropoxy)phenyl]-5-oxopentanoic acid;
30 BV (RS)-5-[3-(4-Chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
BW (RS)-5-(3-Cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
BX (RS)-5-[3-(3-Thienylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
35 BY (RS)-5-[3-(2-Fluorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;

- BZ (RS)-5-[3-(2-Furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CA (RS)-5-[3-(3-Furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 5 CB (RS)-5-[3-(3-Chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CC (RS)-5-[3-(4-Methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CD (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(2-pyridylmethoxy)phenyl]pentanoic acid;
- 10 CE (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]pentanoic acid;
- CF (RS)-5-(2-Methyl-3-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 15 CG (RS)-5-(3,5-Dibenzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- CH (RS)-5-[3-(3-Methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CI (RS)-5-[3-(3-Aminobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 20 CJ (RS)-5-[3-(3-Isothiazolylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CK (RS)-5-(3-benzyloxy-5-hydroxyphenyl)-4-(2,3-dimethylphenyl)-5-oxopentanoic acid;
- 25 CL (RS)-5-(3-Benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- CM (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(4-pyridylmethoxy)phenyl]pentanoic acid;
- CN (RS)-4-(2-Methylphenyl)-5-[3-(3-methyl-2-thienylmethoxy)phenyl]-5-oxopentanoic acid;
- 30 CO (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(3-pyridylmethoxy)phenyl]pentanoic acid;
- CP (RS)-5-(3-Benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoic acid;
- 35 CQ (RS)-5-(3-Benzyloxy-6-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- CR (RS)-5-(3-Benzyloxyphenyl)-4-(3-methylpyrid-4-yl)-5-oxopentanoic acid;

- CS (RS)-5-[3-(1,2,4-Oxadiazol-3-ylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- CT Methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate;
- CU Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chlorophenyl)-5-oxopentanoate;
- 5 CV Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methoxyphenyl)-5-oxopentanoate;
- CW Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 10 CX Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-ethylphenyl)-5-oxopentanoate;
- CY Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chloro-6-fluorophenyl)-5-oxopentanoate;
- CZ Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2,6-dichlorophenyl)-5-oxopentanoate;
- 15 DA Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3,4-dichlorophenyl)-5-oxopentanoate;
- DB Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(4-chlorophenyl)-5-oxopentanoate;
- 20 DC Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(4-methylphenyl)-5-oxopentanoate;
- DD Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3-chlorophenyl)-5-oxopentanoate;
- DE Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3-methoxyphenyl)-5-oxopentanoate;
- 25 DF Methyl (RS)-5-(4-benzyloxyphenyl)-5-oxo-4-phenylpentanoate;
- DG Methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoate;
- DH Methyl (3RS,4RS)-5-(3-benzyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoate;
- 30 DI Methyl (2RS,4RS)-5-(3-benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoate;
- DJ Methyl (RS)-5-[3-(2-methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate;
- DK Methyl (RS)-5-[3-(3,4-methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate;
- 35 DL Methyl (RS)-5-(3-benzyloxy-2-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate;

- DM Methyl (RS)-5-(3,5-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- DN Methyl (RS)-5-(3-benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- 5 DO Methyl (RS)-5-(3-benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoate;
- DP Methyl (RS)-4-(3-benzyloxyphenyl)-4-oxo-3-phenylbutanoate;
- DQ Ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhexanoate;
- DR Ethyl (RS)-7-(3-benzyloxyphenyl)-7-oxo-6-phenylheptanoate;
- 10 DS Ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoate;
- DT Methyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- DU Ethyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 15 DV Methyl (RS)-5-(3-cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- DW Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(3-thienylmethoxy)phenyl]pentanoate;
- DX Methyl (RS)-5-[3-(2-fluorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 20 DY Methyl (RS)-5-[3-(2-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- DZ Methyl (RS)-5-[3-(3-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 25 EA Methyl (RS)-5-[3-(3-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- EB Methyl (RS)-5-[3-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- EC Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-pyridylmethoxy)phenyl]pentanoate;
- 30 ED Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]pentanoate;
- EE Methyl (RS)-5-[3-(3-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 35 EF Methyl (RS)-5-[3-(3-isothiazolylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;

- EG Methyl (RS)-5-[3-(4-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- EH Methyl (RS)-5-[3-(3-nitrobenzyloxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- 5 EI Methyl (RS)-5-[3-(3-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- EJ Methyl (RS)-5-(3-methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- EK Methyl (RS)-5-(3-isopropoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- 10 EL Methyl (RS)-5-[3-(1,2,4-oxadiazol-3-yl-methoxy)phenyl]-4-(2-methylphenyl)-5-oxo-pentanoate;
- EM Methyl (RS)-5-[3-(3-methyl-2-thienylmethoxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- EN Methyl (RS)-5-[3-(3-aminobenzyloxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- 15 EO Ethyl (RS)-5-[3-benzyloxy-5-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- EP (RS)-5-(3-Benzyloxy-5-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 20 EQ 5-[4-(3-Benzyloxyphenyl)-4-oxo-3-phenylbutyl]-1H-tetrazole;
- ER 5-[3-(3-Benzyloxyphenyl)-3-oxo-2-(2-methylphenyl)propyl]-1H-tetrazole;
- ES (RS)-4-(3-Benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylsulphonic acid;
- ET (RS)-4-[3-(3-Thienylmethoxy)phenyl]-3-(2-methylphenyl)-4-oxobutylsulphonic acid;
- 25 EU (RS)-6-(3-Benzyloxyphenyl)-5-(2-methyl-phenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide;
- EV (RS)-6-(3-(3-Thienylmethoxy)phenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide;
- 30 EW Ethyl (RS)-4-(3-benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate;
- EX Diethyl (RS)-4-(3-benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate;
- EY (RS)-4-(3-Benzyloxyphenoxy)-4-phenylbutanoic acid;
- 35 EZ (RS)-4-(2-Carboxy-5-benzyloxyphenoxy)-4-phenylbutanoic acid;
- FA (RS)-4-[2-(N-Ethylcarbamoyl)-5-benzyloxy-phenoxy]-4-phenylbutanoic acid;

- FB (RS)-4-(2-Acetyl-5-benzyloxyphenoxy)-4-phenylbutanoic acid;
FC (RS)-4-[3-(3-Thienylmethoxy)phenoxy]-4-phenylbutanoic acid;
FD (RS)-4-(3-Benzyloxy)phenoxy-4-(2-methylphenyl)butanoic acid;
FE (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-
5 butanoic acid;
FF (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-phenylbutanoic acid;
FG (RS)-4-[2-Acetyl-5-(3-pyridylmethoxy)-phenoxy]-4-phenylbutanoic acid;
FH (RS)-4-[2-Carbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoic acid;
FI (RS)-4-[2-Cyano-5-benzyloxyphenoxy]-4-phenyl-butanoic acid;
10 FJ (RS)-4-[2-(N,N-Dimethylcarbamoyl)-5-benzyloxyphenoxy]-4-phenylbutanoic acid;
FK (RS)-4-[2-(3-Pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
FL (RS)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-
15 methylphenyl)butanoic acid;
FM Ethyl (RS)-4-(3-benzyloxyphenoxy)-4-phenylbutanoate;
FN Ethyl (RS)-4-(2-methoxycarbonyl-5-benzyloxyphenoxy)-4-phenylbutanoate;
FO Ethyl (RS)-4-[(3-benzyloxy)phenoxy]-4-(2-methylphenyl)butanoate;
20 FP Ethyl (RS)-4-[3-(3-thienylmethoxy)phenoxy]-4-phenylbutanoate;
FQ Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-phenylbutanoate;
FR Ethyl (RS)-4-[2-(N-ethylcarbamoyl)-5-benzyloxyphenoxy]-4-phenylbutanoate;
25 FS Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
FT Ethyl (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)-phenoxy]-4-phenylbutanoate;
FU Ethyl (RS)-4-[2-carbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoate;
30 FV Ethyl (RS)-4-[2-N,N-dimethylcarbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoate;
FW Ethyl (RS)-4-[2-carbamoyl-5-(3-thienyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoate
FX Ethyl (RS)-4-(2-acetyl-5-benzyloxyphenoxy)-4-phenylbutanoate;
35 FY Ethyl (RS)-4-(2-cyano-5-benzyloxyphenoxy)-4-phenylbutanoate;
FZ Ethyl (RS)-4-[2-cyano-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;

- GA Ethyl (RS)-4-[2-(3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- GB (R)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- GC (S)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 5 GD (R)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid,
- GE (S)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid.
- GF (RS)-4-[2-Acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 10 GG (RS)-4-(2-Methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- GH (RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- 15 GI (RS)-4-(2-Methylphenyl)-4-[2-thiocarbamoyl-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- GJ (E)-(RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)prop-2-enoyl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- GK Ethyl (RS)-4-[2-acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate,
- 20 GL Ethyl (RS)-4-(2-methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoate;
- GM Ethyl (RS)-4-(2-methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]butanoate;
- 25 GN Ethyl (RS)-4-(2-methylphenyl)-4-[5-(3-thienylmethoxy)-2-thiocarbamoylphenoxy]butanoate;
- GO (RS)-4-[5-Benzoyloxy-2-(methylthio)phenoxy]-4-phenylbutanoic acid;
- GP (E)-(RS)-4-[5-Benzoyloxy-2-(2-carboxyethenyl)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 30 GQ (RS)-4-[2-(1-Methyl-2-carboxyethenyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- GR (RS)-4-[5-Benzoyloxy-2-hydroxyiminomethylphenoxy]-4-(2-methylphenyl)butanoic acid;
- GS (RS)-4-[2-[N-(Carboxymethoxy)iminomethyl]-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 35 GT (RS,RS)-4-[2-(1-Hydroxyethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;

- GU (RS)-4-(2-Methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)-phenoxy]butanoic acid;
- GV (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 5 GW (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- GX (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)-butanoic acid;
- GY (RS)-4-(2-Methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]-
- 10 butanoic acid;
- GZ (RS,RS)-4-(2-Methylphenyl)-4-[2-methyl-sulphinyl-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- HA (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoic acid;
- 15 HB (RS)-4-(5-Benzoyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoic acid;
- HC (RS)-4-[2-Formyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoic acid;
- HD (RS)-4-(2-Methylphenyl)-4-[5-(3-thienylmethoxy)-2-(trifluoroacetyl)-
- 20 phenoxy]butanoic acid;
- HE (RS)-4-(2-Methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)-phenoxy]butanoic acid;
- HF (RS)-4-[2-Cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 25 HG (RS)-4-[2-Carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- HH (RS)-4-[2-{N-(3-Imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- HI (RS)-4-[2-{N-(2-Carboxyethyl)carbamoyl}-5-(3-thienylmethoxy)-
- 30 phenoxy]-4-(2-methylphenyl)butanoic acid;
- HJ (RS)-4-[2-{N-(Carboxymethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- HK (RS)-4-[2-{N-(2-Cyanoethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid ;
- 35 HL (RS)-4-[2-(5-Carboxy-1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;

- HM (RS)-4-[2-{N-(Carbamoylmethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- HN (RS)-4-[2-{N-(Methoxycarbonylmethyl)-carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 5 HO Ethyl (RS)-4-[5-benzyloxy-2-(methylthio)phenoxy]-4-phenylbutanoate;
- HP Ethyl (RS)-4-(2-methylphenyl)-4-[2-methylthio-5-(3-thienylmethoxy)-phenoxy]butanoate;
- HQ Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoate;
- 10 HR Ethyl (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)-butanoate;
- HS Ethyl (RS)-4-[2-formyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- HT Ethyl (RS)-4-[5-(3-thienylmethoxy)-2-trifluoroacetylphenoxy]-4-(2-methylphenyl)butanoate;
- 15 HU Ethyl (RS)-4-[2-carbamoyl-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate;
- HV Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)-butanoate;
- 20 HW Ethyl (RS)-4-(2-methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]-butanoate;
- HX Ethyl (RS)-4-[2-{N-methoxycarbonylmethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- HY Ethyl (RS)-4-[2-{N-(3-imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- 25 HZ Ethyl (RS)-4-[2-{N-(2-cyanoethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoate;
- IA Ethyl (RS)-4-[2-{N-cyanomethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoate;
- 30 IB Ethyl (RS)-4-[2-{N-(2-methoxycarbonylethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- IC Ethyl (RS,RS)-4-(2-methylphenyl)-4-[2-methylsulphinyl-5-(3-thienylmethoxy)phenoxy]butanoate;
- ID Ethyl (E)-(RS)-4-(2-methylphenyl)-4-[2-(2-methoxycarbonylethenyl)-5-benzyloxyphenoxy]-butanoate;
- 35 IE Ethyl (RS)-4-[2-(2-methoxycarbonyl-1-methylethenyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;

- IF Ethyl (RS)-4-(5-benzyloxy-2-hydroxyiminomethylphenoxy)-4-(2-methylphenyl)butanoate;
- IG Ethyl (RS)-4-[2-hydroxyiminomethyl-5-(3-thienylmethoxy))phenoxy]-4-(2-methylphenyl)-butanoate;
- 5 IH (RS)-4-[2-{N-(ethoxycarbonylmethoxy)iminomethyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- II Ethyl (RS,RS)-4-[2-(1-hydroxyethyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoate;
- IJ Ethyl (RS)-4-(2-methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)-
- 10 phenoxy]butanoate;
- IK Ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-ethylphenyl)butanoate;
- IL Ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- 15 IM Ethyl (RS)-4-(2-methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)phenoxy]-butanoate;
- IN Ethyl (RS)-4-[2-cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.
- IO Methyl (RS)-4-[2-(5-methoxycarbonyl-1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- 20 IP (RS)-N-[4-{2-Cyano-5-(3-thienylmethoxy)-phenoxy}-4-(2-methylphenyl)butanoyl]benzenesulphonamide;
- IQ (R)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 25 IR (S)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- IS (R)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- IT Dicyclohexylammonium (R)-4-[2-cyano-5-(3-pyridylmethoxy)phenoxy]-4-
- 30 (2-methylphenyl)butanoate;
- IU (S)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- IV (RS)-5-(3-Benzylthiophenyl)-4-(2-methylphenyl)-5-oxopentanoic acid hemihydrate;
- 35 IW (RS)-4-(2-Methylphenyl)-5-oxo-(1-oxopyrid-3-ylmethoxy)pentanoic acid hemihydrate;

- IX (RS)-4-[2-(Benzyloxyiminomethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid hydrate;
- IY (RS)-4-(2-Carboxycarbonyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoic acid;
- 5 IZ (R)-4-[2-Benzoyl-5-(pyridin-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid;
- JA (RS)-3-[3-(3-Carboxy-1-(2-methylphenyl)propoxy)-4-nitrophenoxy-methyl]benzoic acid;
- JB (RS)-4-[5-[3-(2-Carboxyethyl)benzyloxy]-2-cyano-phenoxy]-4-(2-methylphenyl)butanoic acid;
- 10 JC (RS)-4-[2-(4,5-Dihydrooxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JD (RS)-4-[2-((1R)-1-Carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid ;
- 15 JE (RS)-4-[2-((1S)-1-Carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- JF (RS)-4-[2-(Oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JG (RS)-2-[2-(3-Carboxy-1-(2-methylphenyl)propoxy)-4-(3-thienylmethoxy)benzoylamino]acrylic acid;
- 20 JH (RS)-4-[2-(4-Carboxyoxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JI (RS)-4-(2-Chloro-6-fluorophenyl)-4-(2-cyano-5-(3-thienylmethoxy)-phenoxy)butanoic acid;
- 25 JJ (R)-4-(2-Methylphenyl)-4-(2-nitro-5-(3-pyridylmethoxy)phenoxy)butanoic acid;
- JK (RS)-4-(2-Cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoic acid;
- JL (RS)-4-(Benzo[1,3]dioxol-4-yl)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]butanoic acid;
- 30 JM (RS)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2,3-dimethylphenyl)-butanoic acid;
- JN (RS)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(5-hydroxy-2-methylphenyl)-butanoic acid;
- 35 JO (RS)-5-[2-Cyano-5-(3-thienylmethoxy)-phenoxy]-5-(2-methylphenyl)-pentanoic acid;

- JP (RS)-4-(2-Cyano-3-fluoro-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl) butanoic acid;
- JQ (RS)-2,4-Dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)phenyl]butanoic acid;
- 5 JR (RS)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JS (R)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JT (R)-4-[2-(5-Carboxypyrazol-3-yl)-5-(pyrid-3-ylmethoxy)phenoxy]-4-(2-methylphenyl) butanoic acid;
- 10 JU (RS)-3-(2-Cyano-5-(3-pyridylmethoxy)phenoxy)-2,2-difluoro-3--(2-methylphenyl)propanoic acid;
- JV (RS)-4-[2-(5-Carboxyisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 15 JW (R)-4-[2-(5-Methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JX (R)-4-[2-(5-Carbamoylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- JY (RS)-4-[(2-Acetyl-5-(3-thienylmethoxy)phenoxy)]-4-(2-cyanophenyl)-butanoic acid;
- 20 JZ (R)-4-[(2-Cyano-5-(isothiazol-4-ylmethoxy)phenoxy)]-4-(2-methylphenyl)-butanoic acid;
- KA 3-(2,4-Dibenzyloxy)phenyl propanoic acid;
- KB 2,4-Dibenzyloxyphenoxyacetic acid;
- 25 KC (R)-4-[2-Nitro-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid;
- KD (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}methane sulphonamide;
- KE (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}trifluoromethane sulphonamide;
- 30 KF (R,S)-4-[2-(3-Carboxypropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid;
- KG (R,S)-4-[2-(1,2,4-Oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid; and
- 35 KH (R,S)-4-[2-(Thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid.

The letters A to KH are allocated to compounds for easy reference in this specification.

Preferred species according to the invention are those identified
5 hereinbefore as compounds FE, FK, FL, GG, GH, GI, GJ, GS, GV, GW, GY, HA, HC, HF, HG, HI, HJ, HK, HL, HM, HN, IQ, IS, IY, JC, JF, JG, JH, JI, JJ, JK, JL, JM, JO, JP, JQ, JR, JS, JT, JV, JW, JX, JZ, KC, KD, KE, KF, KG and KH, and their pharmaceutically acceptable salts, N-oxides and prodrugs.


10 Especially preferred according to the invention is that species identified hereinbefore as compound IS and its pharmaceutically acceptable salts, N-oxide and prodrugs.

Compounds of formula I may be prepared by the application or
15 adaptation of known methods, which means methods used heretofore or described in the literature.

Thus, according to a feature of the present invention, compounds of
formula I containing a carboxy moiety are prepared by hydrolysis of the
20 corresponding compound of formula I containing an alkoxycarbonyl moiety. The hydrolysis may be by alkaline hydrolysis using a base, such as alkali hydroxide or carbonate, for example, NaOH, LiOH, KOH or K₂CO₃, in the presence of an aqueous/organic solvent mixture, using organic solvents such as dioxan, THF or methanol, at a temperature from about ambient to about
25 reflux. The hydrolysis may also be by acid hydrolysis using an inorganic acid, such as HCl, in the presence of an aqueous/inert organic solvent mixture, using organic solvents such as dioxan, THF, at a temperature from about 50°C to about 80°C or where alkoxy moiety of the alkoxycarbonyl (ester) group is *t*-butoxy then one may use trifluoroacetic acid at about ambient temperature.

30 According to a further feature of the present invention, compounds of formula I, wherein R¹ is -C≡CO₂H, and R², R³, R⁴, R⁵, m, n, o and p are as hereinbefore defined, are prepared by the reaction of compounds of the formula II,


35




wherein R^{1a} is $-\text{CH}=\text{CBr}_2$, and R^2 , R^3 , R^4 , R^5 , m , n , o and p are as
 5 tetrahydrofuran, at about -70°C , followed by reaction with carbon dioxide at
 about -40°C .

According to a further feature of the present invention, compounds of
 formula I substituted by a carbamoyl or substituted carbamoyl group are
 10 prepared by the reaction of compounds of the formula I having the
 corresponding alkoxy carbonyl (ester) group with concentrated aqueous
 ammonium hydroxide solution or the appropriate alkylamine, dialkylamine or
 aminoalkanol, optionally in an inert organic solvent, such as methanol, or
 aqueous mixture thereof, preferably at a temperature from about 0°C to about
 15 reflux temperature, and optionally under high pressure. Alternatively
 compounds of formula I substituted by a carbamoyl group are prepared by the
 reaction of compounds of the formula I having the corresponding carboxy
 group with an inorganic halogenating agent, such as SOCl_2 , PCl_3 , PBr_3 or PCl_5
 to form the corresponding acid halide ($-\text{COX}$, wherein X' is halo, preferably
 20 chloro or bromo), which is then reacted with NH_3 preferably in the presence of
 a base such as potassium carbonate or a tertiary amine, such as triethylamine
 or pyridine, optionally in an inert solvent, for example dichloromethane,
 dimethylformamide, or an ether, such as diethyl ether or tetrahydrofuran,
 preferably at a temperature from about 0°C to about reflux temperature.
 25 Alternatively, an acid halide wherein X' is chloro is prepared by reacting oxalyl
 chloride with a carboxy moiety. The reaction preferably takes place in an inert
 solvent, such as methylene chloride, at a temperature from about 0°C to about
 reflux.

30 According to a further feature of the present invention, compounds of
 formula I, wherein R^1 is $-\text{C}\equiv\text{CR}^{10}$, and R^2 , R^3 , R^4 , R^5 , m , n , o and p are as
 hereinbefore defined, are prepared by hydrolyzing the ester product of the
 reaction of compounds of formula III,



wherein R^{1b} is iodo, and R^2 , R^3 , R^4 , R^5 , m , n , o and p are as hereinbefore
 5 defined, with compounds of the formula IV




wherein R^{10a} is alkoxycarbonyl or alkoxycarbonylphenyl, and Q^1 is ethynyl.
 10 Preferably the reaction is carried out with the aid of a catalyst, such as tetrakis(triphenylphosphine)palladium, and cuprous iodide, preferably with the aid of a base such as a tertiary amine, such as triethylamine, preferably in a solvent such as dimethylformamide, at a temperature from about room temperature to about 100°C.
 15 As another example, compounds of formula I wherein R^1 contains or is a -alkylOH moiety, are prepared by the reduction of the corresponding compounds of formula I wherein R^1 contains or is an alkoxycarbonyl or formyl or acetyl moiety by reaction with an alkali metal hydride, such as lithium
 20 aluminum hydride or sodium borohydride respectively, preferably in an inert solvent, such as tetrahydrofuran or methanol, preferably at about room temperature.

According to a further feature of the present invention, compounds of
 25 formula I wherein R^1 is $-\text{CH}=\text{CHCO}_2\text{H}$ are prepared by the reaction of compounds of formula I wherein R^1 is $-\text{CHO}$ with diethyl phosphonoacetic acid in the presence of a base such as sodium hydride or *n*-butyl lithium. The reaction preferably takes place in an inert solvent, such as tetrahydrofuran, at a temperature from about -70°C to about room temperature. Alternatively
 30 compounds of formula I wherein R^1 is $-\text{CH}=\text{CHCO}_2\text{H}$ are prepared by the reaction of compounds of formula I, wherein R^1 is $-\text{CHO}$ with malonic acid in the presence of a base such as piperidine in a solvent such as pyridine, at a temperature from about 50° to about 200°C.

According to a further feature of the present invention, compounds of formula I wherein R¹ is -alkenyl(carbonylalkoxy or alkyl) are prepared by the reaction of compounds of formula I, wherein R¹ is -CHO or -C(alkyl)=O with a compound of the formula V

5




wherein Q² is aryl, such as phenyl, Q³ is (alkoxycarbonyl- or alkyl) and X is halo, preferably bromo, with a base such as an alkali metal alkoxide, such as potassium *t*-butoxide. The reaction is preferably carried out in an inert solvent, such as tetrahydrofuran, at a temperature from about 0°C to about reflux temperature. The alkoxycarbonyl moiety of the -CH=CH(carbonylalkoxy) group formed according to this method may be hydrolyzed to the corresponding -CH=CHCO₂H group as described herein.

15

Alternatively, compounds of formula I as described herein except for comprising hydroxy or carboxy moieties, are prepared by reacting compounds of the formula VIIa

20



wherein R^{2a} is hydroxy, wherein R¹ and R³ are as described herein except for being or having thereon hydroxy or carboxy moieties, and R⁴ is as described herein except for having thereon hydroxy moieties or acidic hydrogen atoms, such as carboxy, phosphonic or sulphonic groups, and R⁵, n, o, and p are as described herein, with compounds of formula VIIIa

25



30

wherein Q⁴ is aryl lower alkyl or heteroaryl lower alkyl except for having thereon hydroxy or carboxy moieties. The reaction is carried out in the presence of a triarylphosphine, such as triphenylphosphine, and a dialkyl ester, such as the diisopropyl or diethyl ester, of azodicarboxylic acid,

preferably in an inert solvent such as tetrahydrofuran, preferably at a temperature from about 0°C to about room temperature.


Alternatively, compounds of formula I as described herein except for comprising hydroxy or carboxy moieties, are prepared by reaction with compounds of the formula VIIa as described herein, with compounds of formula IXa

Q⁴X

IXa

wherein Q⁴ is as described herein but with the exclusion of moieties containing hydroxy groups, or acidic hydrogen atoms, such as in carboxy groups, and X is a leaving group, such as halo or alkylsulphonyloxy, such as methane-sulphonyloxy, or arylsulphonyloxy, such as p-toluenesulphonyloxy. The reaction is carried out in the presence of a base, such as sodium hydride or alkali metal alkoxide, such as potassium tert-butoxide, in an inert solvent, such as a ketone, such as acetone or methyl ethyl ketone, or a dipolar aprotic solvent, such as dimethylformamide, at a temperature from about room temperature to about 100°C. Additionally, since VIIa may have a mercapto group in place of the free hydroxyl of R^{2a}, as described herein regarding introduction of mercapto moieties on phenyl groups, regarding introduction of mercapto moieties, then the reaction of VIIa bearing such a mercapto group with Q⁴X would result in the introduction of a Q⁴S- moiety, for example benzylthio, in place of R^{2a}.

Compounds of formula I, wherein R¹ is as described herein except for comprising hydroxy or carboxy moieties, are prepared by reacting compounds of the formula VIIb



VIIb


wherein R^{3a} is hydroxy, wherein R¹ and R² are as described herein except for having thereon hydroxy or carboxy moieties, and R⁴ is as described herein except for having thereon hydroxy moieties or acidic hydrogen atoms, such as

in carboxy, phosphonic or sulphonic groups, and R^5 , o, and p are as described herein, with compounds of formula VIIIb




5 wherein Q^5 is alkyl, cycloalkyl(lower alkyl), cycloalkenyl(lower alkyl), aryl lower alkyl or heteroaryl lower alkyl except for having thereon hydroxy or carboxy moieties. The reaction is carried out in the presence of a triarylphosphine, such as triphenylphosphine, and a dialkyl ester, such as the diisopropyl or
10 diethyl ester, of azodicarboxylic acid, preferably in an inert solvent such as tetrahydrofuran, preferably at a temperature from about 0°C to about room temperature.

Alternatively, compounds of formula I as described herein except for
15 comprising hydroxy or carboxy moieties, are prepared by reaction with compounds of the formula VIIa as described herein, with compounds of formula IXb



20 wherein Q^5 is as described herein but with the exclusion of moieties containing hydroxy groups, or acidic hydrogen atoms, such as in carboxy groups, and X is a leaving group, such as halo or alkylsulphonyloxy, such as methane-sulphonyloxy, or arylsulphonyloxy, such as p-toluenesulphonyloxy. The
25 reaction is carried out in the presence of a base, such as sodium hydride or alkali metal alkoxide, such as potassium tert-butoxide, in an inert solvent, such as a ketone, such as acetone or methyl ethyl ketone, or a dipolar aprotic solvent, such as dimethylformamide, at a temperature from about room temperature to about 100°C. Additionally, since VIIb may have a mercapto
30 group in place of the free hydroxyl of R^{3a} , as described herein regarding introduction of mercapto moieties on phenyl groups, then the reaction of VIIA bearing such a mercapto group with Q^5X would result in the introduction of a Q^5S - moiety, for example benzylthio, in place of R^{3a} .

35 Alternatively, compounds of formula I as described herein except for comprising hydroxy or carboxy moieties, are prepared by reacting compounds of the formula VIIc




5 wherein R^{2a} and R^{3a} are hydroxy, wherein R¹ is as described herein except for being or having thereon hydroxy or carboxy moieties, and R⁴ is as described herein except for having thereon hydroxy moieties or acidic hydrogen atoms, such as carboxy, phosphonic or sulphonic groups, and R⁵, o, and p are as described herein, with compounds of formula IXa

10




wherein Q⁴ is as described herein but with the exclusion of moieties containing hydroxy groups, or acidic hydrogen atoms, such as in carboxy groups, and X is
 15 a leaving group, such as halo or alkylsulphonyloxy, such as methane-sulphonyloxy, or arylsulphonyloxy, such as p-toluenesulphonyloxy. The reaction is carried out in the presence of a base, such as sodium hydride or alkali metal alkoxide, such as potassium tert-butoxide, in an inert solvent, such as a ketone, such as acetone or methyl ethyl ketone, or a dipolar aprotic
 20 solvent, such as dimethylformamide, at a temperature from about room temperature to about 100°C. Additionally, since VIIc may have a mercapto group in place of the free hydroxyls of R^{2a} and/or R^{3a}, as described herein regarding introduction of mercapto moieties on phenyl groups, then the reaction of VIIc bearing such a mercapto group with Q⁴X would result in the
 25 introduction of a Q⁵S- moiety, for example benzylthio, in place of R^{3a} and/or R^{2a}.

Thus, according to a feature of the present invention, compounds of formula I, as hereinbefore defined but with the exclusion of groups having
 30 thereon hydroxy moieties or acidic hydrogen atoms, such as carboxy, phosphonic or sulphonic groups, are prepared by the reaction of compounds of formula X




X

wherein R^1 , R^2 , R^3 , R^5 , R^{15} , m , n and p are as hereinbefore defined but with the exclusion of groups being or having thereon hydroxy moieties or acidic hydrogen atoms, such as carboxy groups, with compounds of formulae XI or XII




wherein X^1 is halo, such as chloro or, preferably, bromo, and R^{16a} is esterified carboxy, in the presence of an alkali metal alkoxide, such as potassium tert-butoxide. The reaction preferably takes place in an inert solvent, such as tetrahydrofuran, at temperatures from about -40°C to about reflux.

According to a further feature of the present invention, compounds of formula I, as hereinbefore defined but with the exclusion of groups having thereon hydroxy moieties or acidic hydrogen atoms, such as carboxy, phosphonic or sulphonic groups, are prepared by the reaction of compounds of formula X with compounds of formulae XIII




wherein R^{16b} represents esterified carboxy, sulphonic or phosphonic, in the presence of a base, for example an alkali metal alkoxide, such as potassium tert-butoxide. The reaction preferably takes place in an inert solvent, such as tetrahydrofuran, at temperatures from about room temperature to about reflux.

Compounds of formula I, wherein R^1 is as described herein except for comprising hydroxy or carboxy moieties, are prepared by reacting compounds of the formula VIId



wherein R^{4a} is hydroxy, wherein R^1 , R^2 and R^3 are as described herein except for being or having thereon hydroxy or carboxy moieties, and R^5 , m , n and p are as described herein, with compounds of formula XIV



wherein X^1 is a leaving group, such as halo or alkylsulphonyloxy, such as methanesulphonyloxy, or arylsulphonyloxy, such as *p*-toluenesulphonyloxy, and R^{15} and R^{16a} are as herein defined, in the presence of an alkali metal alkoxide, such as potassium *tert*-butoxide. The reaction preferably takes place in an inert solvent, such as a ketone, such as acetone or methyl ethyl ketone, or a dipolar aprotic solvent, such as dimethylformamide, at a temperature from about room temperature to about reflux.

Compounds of formula I, wherein R^1 is as described herein except for being or having thereon a OH, CO_2H or NH group, may be prepared by reacting compounds of formula VIId wherein R^{4a} is hydroxy, with compounds of formula XIVa



wherein X^{1a} is hydroxy and R^{16a} is esterified carboxy. The reaction is carried out in the presence of a triarylphosphine, such as triphenylphosphine, and a dialkyl ester, such as diisopropyl or diethyl ester, of azodicarboxylic acid, preferably in an inert solvent, such as tetrahydrofuran, preferably at a temperature from about $0^\circ C$ to about room temperature. The reaction may be particularly suitable for the preparation of compounds with defined chirality at the carbon to which R^{15} is attached.

Compounds of formula I, wherein R^1 is as described herein may also be prepared by reacting compounds of formula VIId wherein R^{4a} is fluorine with

alkali metal salts of compounds of formula XIVa wherein X^{1a} is hydroxy and R^{16a} is carboxy or esterified carboxy. The reaction preferably takes place in an inert solvent such as THF at a temperature between about room temperature and the reflux temperature. The reaction may be particularly
5 suitable for the preparation of compounds with defined chirality at the carbon to which R^{15} is attached.

Compounds of formula XIVa, where X^{1a} is hydroxy and R^{16a} is CO_2tBu , and with defined chirality at the carbon to which R^{15} is attached, may
10 be prepared by reduction of the corresponding ketones with bis[(1S,2R,3S,5S)-pinan-3-yl]chloroborane, or bis[(1R,2R,3R,5R)-pinan-3-yl]chloroborane, in an inert solvent, such as diethyl ether, at a temperature between about $-40^\circ C$ and about room temperature.

15 According to a further feature of the present invention, compounds of formula I, wherein R^1 or R^2 is tetrazolyl, are prepared by the reaction of the corresponding compounds of formula I, wherein R^1 or R^{16} is $-CN$, with a source of trimethyltin azide, such as sodium azide and trimethyltin chloride, in an inert solvent, such as toluene, preferably at about the reflux temperature.
20

According to a further feature of the present invention, compounds of formula I containing an amino moiety, are preferably prepared by the reduction of the corresponding compounds of formula I containing a nitro moiety. When compounds of formula I contain alkyl moieties rather than alkenyl or alkynyl
25 moieties, the reduction is preferably carried out by means of catalytic hydrogenation, such as using a palladium on carbon catalyst. Preferably the reaction is carried out in an inert solvent, such as an alkyl alkanoate ester, at or near room temperature. When the starting material of formula I contains an ester moiety, the ester used as reaction medium is preferably chosen so as to avoid transesterification, such as methyl esters of formula I are preferably
30 reduced in a solvent which is a methyl ester, such as methyl acetate. When the compound of formula I contains an alkenyl or alkynyl moieties, the reduction is preferably carried out in a system such as iron and hydrochloric acid, in the presence of a solvent such as an alkanol. When the starting material of
35 formula I is an ester, the alkanol used as reaction medium is preferably chosen so as to avoid transesterification, such as methyl esters of formula I are

preferably reduced in the presence of methanol and ethyl esters of formula I are preferably reduced in the presence of ethanol.

According to a further feature of the present invention, compounds of formula I containing a hydroxyaryl group are prepared from the corresponding compounds of formula I containing an arylmethoxy group, such as a p-methoxybenzyloxy group, by reaction with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone in an inert solvent system, such as dichloromethane and tert-butanol, preferably in the presence of a buffer (pH about 6 to 8), such as a phosphate buffer, preferably at about room temperature, or are prepared from the corresponding compounds of formula I containing an arylmethoxy group such as a benzyloxy group instead of the said hydroxy group, by hydrogenolysis, for example in solution in a mixture of an alkanol and an alkyl alkanoate ester, containing dilute hydrochloric acid, in the presence of a catalyst of palladium on carbon. When the starting material of formula I is an ester, the alkanol and ester used as reaction medium are preferably chosen so as to avoid transesterification, such as methyl esters of formula I are preferably hydrogenolysed in a solvent which is a mixture of methanol and a methyl ester, such as methyl acetate.

20

According to a further feature of the present invention, compounds of general formula I containing a cyano group are prepared from the corresponding compounds of formula I containing a carbamoyl group by heating with an alkanoic acid anhydride, such as acetic anhydride.

25

According to a further feature of the present invention, compounds of general formula I containing a pyrazolyl group are prepared from the corresponding compounds of formula I containing an alkylcarbonyl group, such as acetyl group, by reaction with an alkyl formate, such as ethyl formate, in the presence of sodium hydride, followed by treatment with dilute hydrochloric acid and then with hydrazine hydrate.

30

According to a further feature of the present invention, for preparing compounds of formula I wherein R¹ contains a substituted carbamoyl having the nitrogen thereof substituted by a cyano(lower alkyl), aralkyl, heteroaralkyl, alkoxycarbonyl(lower alkyl) or alkoxycarbonyl(aryl substituted lower alkyl) moiety may be prepared by reacting the corresponding compounds of formula I

35

wherein R¹ contains a carboxy moiety with cyano(lower alkyl)NH₂, such as aminopropionitrile, aryalkylNH₂, such as benzylamine, heteroaralkylNH₂, such as aminomethyl-pyridine, alkoxycarbonyl(lower alkyl)NH₂, such as glycine methyl ester, or alkoxycarbonyl(aryl substituted lower alkyl)NH₂, such as phenylalanine ethyl ester, in the presence of an activating agent such as 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide in an inert solvent such as dimethylformamide, and preferably takes place in the presence of a catalyst, such as 1-hydroxy-benzotriazole, at a temperature from about 0°C to about 60°C.

10

Alternatively, according to a further feature of the present invention, compounds of formula I wherein R¹ contains a substituted carbamoyl having the nitrogen thereof substituted by a cyano(lower alkyl), aryalkyl, heteroaralkyl, alkoxycarbonyl(lower alkyl) or alkoxycarbonyl(aryl substituted lower alkyl) moieties may be prepared by reacting the corresponding compounds of formula I wherein R¹ contains a carboxy moiety that is converted to an acid halide as hereinbefore described, with cyano(lower alkyl)NH₂, aryalkylNH₂, heteroaralkylNH₂, alkoxycarbonyl(lower alkyl)NH₂ or alkoxycarbonyl(aryl substituted lower alkyl)NH₂, in the presence of a base, such as triethylamine in an inert solvent, such as dichloromethane, at a temperature from about 0°C to about reflux.

20

According to a further feature of the present invention, for preparing compounds of formula I wherein R¹⁶ is arylsulphonylcarbamoyl or heteroarylsulphonylcarbamoyl may be prepared by reacting the corresponding compounds of formula I wherein R¹⁶ is carboxy with an activating agent such as N,N-carbonyldiimidazole in an inert solvent such as dichloromethane followed by reaction with an arylsulphonamide, alkylsulphonamide or heteroarylsulphonamide in dimethylformamide at or near room temperature.

30

Compounds of formula I, wherein R¹ is alkoxycarbonylpyrazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, are prepared by reaction of compounds of the formula I, wherein R¹ -C(=O)CH₃, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, with diethyl oxalate, in an inert solvent, such as toluene, in the presence of an alkali metal hydride, such as sodium hydride, or an alkali metal alkoxide, such as potassium ethoxide, at temperatures from room temperature

35

to about reflux temperature, to prepare compounds of formula I wherein R^1 is $-\text{COCH}_2\text{COCO}_2\text{alkyl}$, which may be treated as a crude product with hydrazine hydrate in an inert solvent such as ethanol and in the presence of acetic acid, at a temperature from about ambient to about reflux to yield the

5 alkoxy carbonylpyrazole. The pyrazolyl nitrogen atom may be further substituted by reacting it with a lower alkyl halide or aralkyl halide or a lower alkyl ester of a haloalkanoic acid, e.g. ethyl bromoacetate, in an inert solvent, such as DMF or THF, in the presence of base, such as potassium carbonate or sodium hydride, at temperatures between room temperature and the reflux

10 temperature.

According to a further feature of the present invention compounds of formula I wherein R^1 is oxazolyl or alkoxy carbonyloxazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, may be prepared by

15 the oxidation of the corresponding dihydrooxazole or alkoxy carbonyldihydrooxazole, with nickel peroxide or 2,3-dichloro-5,6-dicyanobenzoquinone, in an inert solvent, such as toluene, at a temperature at or about reflux.

20 Compounds of formula I or VIId wherein R^1 is dihydrooxazole or alkoxy carbonyldihydrooxazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, may be prepared by reaction of compounds of formula I or VIId wherein R^1 is $\text{HOCH}_2\text{CH}_2\text{NHC}(=\text{O})-$ or $\text{HOCH}_2\text{CH}(\text{CO}_2\text{alkyl})\text{NHC}(=\text{O})-$, with thionyl chloride in an inert solvent, such as

25 dichloromethane, at a temperature at or about room temperature. Alternatively this dehydration reaction can be performed by reaction with Burgess reagent [(methoxycarbonylsulphamoyl)triethylammonium hydroxide, inner salt] in an inert solvent such as tetrahydrofuran at a temperature at or about reflux.

30 According to a further feature of the present invention compounds of formula I wherein R^1 is alkoxy carbonylisoxazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, may be prepared by reaction of compounds of formula I wherein $R^1 = -\text{C}(=\text{O})\text{CH}_3$ and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, with diethyl oxalate, in an inert

35 solvent, such as toluene, in the presence of an alkali metal hydride, such as sodium hydride, at a temperature from room temperature to about reflux temperature, to prepare compounds of formula I wherein R^1 is

-C(=O)CH₂COCO₂alkyl, which may be treated as a crude product with hydroxylamine hydrochloride, in the presence of an acid, such as hydrochloric acid, at a temperature from room temperature to about reflux temperature to yield the alkoxy carbonyl isoxazole.

- 5 According to a further feature of the present invention compounds of formula I wherein R¹ is thiazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, may be prepared by reaction of compounds of formula I wherein R¹ is -C(=S)NH₂ and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, with chloroacetaldehyde, in an inert
10 solvent, such as aqueous ethanol, at a temperature from room temperature to about reflux temperature.

- According to a further feature of the present invention compounds of formula I wherein R¹ is oxadiazolyl, and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, may be prepared by reaction of
15 compounds of formula I wherein R¹ is -CN and preferably Y is an oxygen atom and carboxy moieties thereon are esterified, with hydroxylamine hydrochloride, in an inert solvent, such as aqueous ethanol, in the presence of an alkali metal hydroxide, such as sodium hydroxide, at a temperature from room temperature to about reflux temperature, followed by treatment with trimethyl orthoformate,
20 at a temperature from room temperature to about reflux temperature.

- Compounds of formula I wherein R¹ is alkylsulphinyl, are prepared by oxidising compounds of the formula I wherein R¹ is alkylthio. The reaction preferably takes place in an inert solvent, such as acetone, in the presence an oxidising agent, such as potassium peroxy monosulfate, at temperatures
25 between about 0°C and room temperature.

- Compounds of formula I wherein R¹ is alkylthio are prepared by alkylating the corresponding compounds of formula I wherein R¹ is mercapto. The reaction preferably takes place in an inert solvent, such as acetone, in the presence of potassium carbonate with an alkyl halide at temperatures between
30 about room temperature and the reflux temperature of the solvent.

- Compounds of formula I, wherein R¹ -CH=NHOH, and preferably Y is oxygen and carboxy moieties thereon are esterified, are prepared by reacting compounds of formula I, wherein R¹ is formyl, and preferably Y is oxygen and carboxy moieties thereon are esterified, with hydroxylamine hydrochloride, in
35 the presence of a base, such as pyridine, at temperatures from about room temperature to about reflux.

Compounds of formula I, wherein R¹ is -CH=NHOalkylCO₂alkyl, and preferably Y is oxygen and carboxy moieties thereon are esterified, are prepared by reacting compounds of formula I, wherein R¹ -CH=NHOH, and preferably Y is oxygen and carboxy moieties thereon are esterified, with an alkyl bromoalkanoate, such as ethyl bromoacetate, in the presence of an alkali metal hydride, such as sodium hydride. The reaction is preferably carried out in an inert solvent, such as dimethylformamide, at temperatures from about room temperature to about 80°C.

Compounds of formula I, having a CF₂alkyl wherein the alkyl may be halogenated as defined, such as trifluoromethyl, Y is oxygen atom and carboxy moieties thereon are esterified, are prepared by reacting compounds of the formula I, wherein R¹ is -C(=O)alkyl, Y is oxygen atom and carboxy moieties thereon are esterified, with diethylaminosulfur trifluoride. The reaction is preferably carried out in an inert solvent, such as dichloromethane, at a temperature between about -5°C and room temperature.

Compounds of formula I, wherein R¹ is -C(=O)NHalkylCONH₂ or -C(=O)NHalkylCO₂alkyl, are prepared by reacting compounds of the formula I, wherein R¹ is -C(=O)NHalkylCN, with potassium carbonate in alkylOH at room temperature.

Compounds of formula I, wherein R¹ is -C(=O)(alkenylaryl or alkenylheteroaryl), are prepared by reaction of compounds of the formula I, wherein R¹ -C(=O)CH₃, with (aryl or heteroaryl)carboxaldehyde, in the presence of aqueous sodium hydroxide. The reaction is preferably carried out in an alcohol, such as ethanol, at about room temperature.

As another example, compounds of formula I containing a thiocarbamoyl group are prepared from the corresponding compounds of formula I containing a carbamoyl group, by reaction with phosphorus pentasulfide or 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide, preferably in a solvent such as pyridine or toluene, and preferably at a temperature from about 0°C to about reflux.

As another example according to the invention, the compounds of formula I wherein a heteroaryl group thereon contains one or more nitrogen ring atoms, can be converted to the corresponding N-oxides preferably by means of reaction with a mixture of hydrogen peroxide and an organic acid, e.g. acetic acid, preferably at about room temperature to about 90°C. Alternatively, the oxidation is carried out by reaction with hydrogen peroxide in the presence of sodium tungstate at temperatures between room temperature

and about 60°C. This latter method is preferred for compounds of formula I containing an acid-labile group, such as a carbon-carbon double bond.

As another example according to the invention, the compounds of formula I containing a cyano moiety, may be prepared the corresponding
5 compound of formula I containing a formyl group by reacting with diammonium hydrogen phosphate and an alkyl nitrate, such as n-propyl nitrate, in an organic acid solvent, such as glacial acetic acid, at about reflux.

As another example according to the invention, the compounds of formula I containing a hydroxy moiety on a phenyl moiety, may be prepared
10 from the corresponding compound of formula I containing a formyl group on a phenyl moiety by reacting with hydrogen peroxide.

The compounds of the present invention are useful in the form of the free base or acid or in the form of a pharmaceutically acceptable salt thereof. All
15 forms are within the scope of the invention.

Where the compound of the present invention is substituted with a basic moiety, acid addition salts are formed and are simply a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the free base form. The acids which can be used to prepare the acid addition salts
20 include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the patient in pharmaceutical doses of the salts, so that the beneficial inhibitory effects on endothelin inherent in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said
25 basic compounds are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt, per se, is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification, and identification, or when it is used as intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures.
30 Pharmaceutically acceptable salts within the scope of the invention are those derived from the following acids: mineral acids such as hydrochloric acid, sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid,
35 cyclohexylsulfamic acid, quinic acid, and the like. The corresponding acid addition salts comprise the following: hydrohalides, such as hydrochloride and hydrobromide, sulfate, phosphate, nitrate, sulfamate, acetate, citrate, lactate,

5 tartarate, malonate, oxalate, salicylate, propionate, succinate, fumarate, maleate, methylene-bis- β -hydroxynaphthoates, gentisates, mesylates, isethionates and di-p-toluoyltartrates, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, cyclohexylsulfamate and quinate, respectively.

10 According to a further feature of the invention, acid addition salts of the compounds of this invention are prepared by reaction of the free base with the appropriate acid, by the application or adaptation of known methods. For example, the acid addition salts of the compounds of this invention are prepared either by dissolving the free base in aqueous or aqueous-alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

15 The parent compounds of this invention can be regenerated from the acid addition salts by the application or adaptation of known methods. For example, parent compounds of the invention can be regenerated from their acid addition salts by treatment with an alkali, such as aqueous sodium bicarbonate solution or aqueous ammonia solution.

20 Where the compound of the invention is substituted with an acidic moiety, base addition salts may be formed and are simply a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the free acid form. The bases which can be used to prepare the base addition salts include preferably those which produce, when combined with the free acid, pharmaceutically acceptable salts, that is, salts whose cations are non-
25 toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial inhibitory effects on endothelin inherent in the free acid are not vitiated by side effects ascribable to the cations. Pharmaceutically acceptable salts, including for example alkali and alkaline earth metal salts, within the
30 scope of the invention are those derived from the following bases: sodium hydride, sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminum hydroxide, lithium hydroxide, magnesium hydroxide, zinc hydroxide, ammonia, ethylenediamine, dicyclohexylamine, N-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine,
35 diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, tetramethylammonium hydroxide, and the like.

Metal salts of compounds of the present invention may be obtained by contacting a hydride, hydroxide, carbonate or similar reactive compound of the chosen metal in an aqueous or organic solvent with the free acid form of the compound. The aqueous solvent employed may be water or it may be a
5 mixture of water with an organic solvent, preferably an alcohol such as methanol or ethanol, a ketone such as acetone, an aliphatic ether such as tetrahydrofuran, or an ester such as ethyl acetate. Such reactions are normally conducted at ambient temperature but they may, if desired, be conducted with heating.

10 Amine salts of compounds of the present invention may be obtained by contacting an amine in an aqueous or organic solvent with the free acid form of the compound. Suitable aqueous solvents include water and mixtures of water with alcohols such as methanol or ethanol, ethers such as tetrahydrofuran, nitriles such as acetonitrile, or ketones such as acetone. Amino acid salts may
15 be similarly prepared.

The parent compounds of this invention can be regenerated from the base addition salts by the application or adaptation of known methods. For example, parent compounds of the invention can be regenerated from their base addition salts by treatment with an acid, such as hydrochloric acid.

20 As will be self-evident to those skilled in the art, some of the compounds of this invention do not form stable acid addition salts. However, acid addition salts are most likely to be formed by compounds of this invention having a nitrogen-containing heteroaryl group.

As well as being useful in themselves as active compounds, salts of
25 compounds of the invention are useful for the purposes of purification of the compounds, for example by exploitation of the solubility differences between the salts and the parent compounds, side products and/or starting materials by techniques well known to those skilled in the art.


It will be apparent to those skilled in the art that certain compounds of
30 formula I can exhibit isomerism, for example geometrical isomerism and optical isomerism. Geometrical isomers include the cis and trans forms of compounds of the invention having an alkenyl moiety. Optical isomers contain asymmetric centers. These asymmetric centers may independently be in either the R or S configuration. All isomers within formula I, and their mixtures, are within the
35 scope of the invention.

Such isomers can be separated from their mixtures, by the application or adaptation of known methods, for example chromatographic techniques and

recrystallization techniques, or they are separately prepared from the appropriate isomers of their intermediates, for example by the application or adaptation of methods described herein.


5 The starting materials and intermediates may be prepared by the application or adaptation of known methods. Methods for preparing starting materials and intermediates are described in the Reference Examples or by chemical equivalents thereof.

10 According to a further example of the present invention, compounds of formula X are prepared by reacting solutions of Grignard reagents, derived by the reaction of compounds of formula XV



15


wherein X^1 and R^{15} are as hereinbefore defined, with magnesium in an ethereal solvent, such as diethyl ether, with compounds of the formula XVI




20

wherein R^1 , R^2 , R^3 , R^5 , R^{15} , n , m , and p , are as hereinbefore defined. The reaction is followed by heating to the reflux temperature, followed by reaction with dilute hydrochloric acid.

25 According to a further example of the present invention, compounds of formula XVII



30 wherein R^1 , R^2 , R^3 , R^5 , R^{15} , n , m , and p , are as hereinbefore defined, and Y is carbonyl, are prepared by reacting compounds of formula XVIII




wherein R^1 , R^2 , R^3 , R^5 , R^{15} , n , m , and p , are as hereinbefore defined, with
 5 compounds of formula XIX



wherein X^1 is as hereinbefore defined, in the presence of an alkali metal
 10 alkoxide, such as potassium tert-butoxide. The reaction preferably takes place
 in an inert solvent, such as tetrahydrofuran, at temperatures between -40°C
 and the reflux temperature of the reaction mixture.

Alternatively, compounds of formula XVIIa,



wherein R^1 , R^2 , R^3 , R^5 , R^{15} , n , m , and p , are as hereinbefore defined, and Y is
 carbonyl, are prepared by reacting of compounds of formula XVIII, with
 acrylonitrile, in the presence of a base, for example an alkali metal alkoxide,
 20 such as potassium tert-butoxide. The reaction preferably takes place in an
 inert solvent, such as tetrahydrofuran, at temperatures between room
 temperature and the reflux temperature of the reaction mixture.

Compounds of formula VII(a) or VII(b), as hereinbefore defined, are
 prepared from compounds of formula VIIC, as hereinbefore defined, by reaction
 25 with compounds of formula IXb, as hereinbefore defined, in the presence of a
 base, for example an alkali metal alkoxide, such as potassium tert-butoxide.
 Alternatively, salts, such as alkali metal salts, of the compounds of formula VIIC,
 are prepared from compounds of formula VIIC by known methods beforehand
 or in situ, for example by reaction with an alkali metal carbonate, alkoxide or

hydride), are used for the reaction with the compounds of formula IXb. The reaction preferably takes place in an inert solvent, such as a ketone, such as acetone or methyl ethyl ketone, or a dipolar aprotic solvent, such as dimethyl-formamide, at temperatures between room temperature and the reflux
 5 temperature of the reaction mixture.

Alternatively, compounds of formula VII(a) or VII(b), as hereinbefore defined, are prepared from compounds of formula VIIc, as hereinbefore defined, by reaction with compounds of formula VIIIb, as hereinbefore defined, in the presence of a triarylphosphine, such as triphenylphosphine, and a
 10 dialkyl ester, such as the diisopropyl or diethyl ester, of azodicarboxylic acid, preferably in an inert solvent such as tetrahydrofuran, preferably at temperatures between 0°C and room temperature.

According to a further feature of the invention compounds of formula VIIC, wherein R¹ is pyridyl, R^{2a} and R^{3a} are hydroxy, and o and p are zero, may be
 15 prepared by reaction of compounds of formula VIIC, wherein R¹ is pyridyl, R^{2a} and R^{3a} are methoxy, and o and p are zero, with pyridine hydrochloride at a temperature of about 160°C.


Compounds of formula VIIC, wherein R¹ is pyridyl, R^{2a} and R^{3a} are methoxy, and o and p are zero, may be prepared by reacting solutions of
 20 Grignard reagents derived by reaction of a bromodimethoxybenzene, such as 1-bromo-2,4-dimethoxybenzene with magnesium in an inert solvent, such as tetrahydrofuran, with a bromopyridine, such as 2-bromopyridine, in the presence of bis-triphenylphosphinepalladium(II) chloride at a temperature about the reflux temperature.

25 Compounds of formula XIV as hereinbefore defined except that R^{16a} is an esterified carboxy, and X¹ is a chlorine atom are prepared by the reaction of compounds of formula XX



30 wherein R¹⁵ is as hereinbefore defined, with [(1-ethoxy-cyclopropyloxy)-trimethyl]silane, in the presence of titanium (IV) chloride. The reaction preferably takes place in an inert solvent, such as dichloromethane, at temperatures between about 0°C and about room temperature.

35 Compounds corresponding to compounds of formula I wherein R¹ is mercapto are prepared by reaction of compounds of the formula XXI



wherein R^2 , R^3 , R^5 , R^{15} , X^1 , n , m , o and p , are as hereinbefore defined, with sodium hydrogensulphide at temperatures between room temperature and the reflux temperature of the solvent.

Nitrohalophenyl, preferably nitrofluorophenyl, compounds optionally substituted alkoxy, cycloalkyloxy arylalkoxy, e.g. arylmethoxy, or heteroarylalkoxy, such as a heteroarylalkoxy group, may be reacted with an alkali metal hydroxide or hydrogensulphide, such as KOH or KSH to prepare the corresponding hydroxynitrophenyl or mercaptonitrophenyl compounds, such as the precursors VIIa, VIIb or VIIc. The reaction preferably takes place in an alcohol, such as t-butanol, at reflux temperature. Alternatively, compounds of the formula $Q^4(OH \text{ or } SH)$ may be reacted with the nitrohalophenyl under the aforesaid conditions to introduce Q^4O^- or Q^4S^- moieties on the phenyl moiety in place of the halo groups.

Intermediates for preparing compounds of formula I having a free hydroxyl group thereon, may be prepared from the compound wherein the hydroxyl moiety is protected by a $-CH_2OCH_2CH_2Si(CH_3)_3$ moiety that is removed by hydrochloric acid or tetrabutylammonium fluoride. The reaction is preferably carried out in an inert solvent, such as THF, at a temperature from about room temperature to about reflux.

Intermediates, for preparing compounds of formula I, having a protected hydroxy moiety may be prepared from the corresponding compounds having a free hydroxyl moiety by reaction with 2-(trimethylsilyl)ethoxymethyl chloride. The reaction is preferably carried out in an inert solvent, such as DMF, in the presence of sodium hydride at room temperature.

The following Examples illustrate the preparation of the compounds according to the invention and the Reference Examples illustrate the preparation of the intermediates.

- 5 In the nuclear magnetic resonance (NMR) spectra, chemical shifts are expressed in ppm relative to tetramethylsilane. Abbreviations have the following meanings:

 s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, dd = doublet
10 of doublets.

EXAMPLE 1 Compound A-W

 A solution of methyl 2-benzyloxy-4-(4-chlorobenzyloxy)benzoate (1.88
15 g) in dioxan (37 mL) and 1M NaOH (12.5 mL) is refluxed for 4 hours. The solution is evaporated, the residue diluted with water, and brought to pH 1 with 2 M HCl. The precipitate is extracted with ethyl acetate, washed with water, dried and evaporated. The residue is recrystallized from ethyl acetate to yield
20 2-benzyloxy-4-(4-chlorobenzyloxy)benzoic acid (0.8 g, 44%) as colourless crystals, m.p. 150-152°C. ¹H NMR (DMSO) 5.15 (2H,s), 5.2(2H,s), 6.7(1H,d), 6.8(1H,s), 7.3-7.5(9H,m), 7.7(1H,d).

 By proceeding in a similar manner, but using the appropriate amount of an ester precursor, there are prepared:

- 25 2-Benzyloxy-4-(3-phenylpropyloxy)benzoic acid. m.p. 74-75°C;
 2,4-Di-(4-chlorobenzyloxy)benzoic acid. m.p. 147-149°C;
 2-Benzyloxy-4-(2-(3-indolyl)ethoxy)benzoic acid. m.p. 68-69°C;
 2-Hydroxy-4-benzyloxybenzoic acid. m.p. 182°C;
30 2-(3-Phenylpropyloxy)-4-benzyloxybenzoic acid. m.p. 101-102°C;
 2-Benzyloxy-4-(2-naphthylmethoxy)benzoic acid. m.p. 134-136°C;
 2-Benzyloxy-4-(1-naphthylmethoxy)benzoic acid. m.p. 145-147°C;
 2-Benzyloxy-4-(3,4-methylenedioxybenzyloxy)benzoic acid. m.p. 112-114°C;
 2-(2-Pyridylmethoxy)-4-benzyloxybenzoic acid. m.p. 147-148°C;
35 2-(2-Phenylethoxy)-4-benzyloxybenzoic acid. m.p. 100-101°C;
 2-Cyclohexylmethoxy-4-benzyloxybenzoic acid. m.p. 99-100°C;
 2-(4-Chlorobenzyloxy)-4-benzyloxybenzoic acid. m.p. 145-146°C;

- 2-Benzyloxy-4-(2-phenylethoxy)benzoic acid. m.p. 116-118°C;
2-Benzyloxy-4-(2-methylpropoxy)benzoic acid. m.p. 162-164°C;
2-Benzyloxy-4-(4-nitrobenzyloxy)benzoic acid. m.p. 203-205°C;
2-Benzyloxy-4-(4-fluorobenzyloxy)benzoic acid. m.p. 130-132°C;
5 4-Benzyloxy-2-(2-methylpropoxy)benzoic acid. m.p. 96°C;
2-(4-Pyridylmethoxy)-4-benzyloxybenzoic acid. m.p. 180-181°C;
2-Benzyloxy-4-(3,4-dichlorobenzyloxy)benzoic acid. m.p. 149-150°C;
2-Benzyloxy-4-(4-methoxybenzyloxy)benzoic acid. m.p. 127-129°C;
2-Benzyloxy-4-(3-methoxybenzyloxy)benzoic acid. m.p. 104-105°C; and
10 2,4-Dibenzyloxybenzoic acid. m.p. 220-222°C.

EXAMPLE 2 Compound X

- To a solution of 2.5 M n-butyl lithium (3.64 mL) in dry tetrahydrofuran
15 (THF) (20 mL) that is cooled to -70°C under nitrogen is added a solution of
diethyl phosphonoacetic acid (0.89 g) in THF (5 mL). After 30 minutes a
solution of 2,4-dibenzyloxybenzaldehyde (1.46 g) in THF (5 mL) is slowly
added to the cooled solution so as to maintain the temperature. The resulting
mixture is stirred at -70°C for 3 hours and then at 25°C for 18 hours. The
20 resulting suspension is treated with water, the THF is removed in vacuo, the
residue is extracted with ethyl acetate, the extract is dried and evaporated.
The residual solid is recrystallized from isopropanol to yield (E)-3-(2,4-
dibenzyloxyphenyl)prop-2-enoic acid as colourless crystals (0.6 g), m.p. 153-
155°C. ¹H NMR (DMSO) 5.15 (2H,s), 5.21(2H,s), 6.4(1H,d,J=16Hz), 6.7(1H,d),
25 6.85(1H,s), 7.3-7.5(10H,m), 7.6(1H,d), 7.8(1H,d,J=16Hz).

EXAMPLE 3 Compound Y

- A solution of 1-(2,4-dibenzyloxyphenyl)-2,2-dibromoethylene (4.85 g) in
30 dry THF (100 mL) is treated with 2.5 M n-butyl lithium (9 mL) at -70°C under
nitrogen. The mixture is stirred for 45 minutes at -70°C and then at 25°C for 45
minutes. The mixture is then cooled to -40°C, treated with solid CO₂ (10 g) and
then stirred at 25°C for 24 hours. The solution is evaporated, the residue taken
up in water. and the pH is adjusted to pH 14 with 1 M NaOH. The water is
35 washed with pentane, the pH adjusted to pH 1 and extracted with
dichloromethane. The extract is washed with brine, dried and evaporated. The
residue is recrystallized from ethyl acetate/pentane to yield 3-(2,4-

dibenzoyloxyphenyl)prop-2-ynoic acid as yellow crystals (1.5 g, 42%), m.p. 125°C (dec.). ¹H NMR (DMSO) 5.15 (2H,s), 5.25(2H,s), 6.7(1H,d), 6.85(1H,s), 7.3-7.5(11H,m).

5 EXAMPLE 4 Compound Z and AA

A solution of methyl N-(2,4-dibenzyloxybenzoyl)glycinate (900 mg) in methanol (20 mL) is treated with 1 M NaOH (20 mL) and stirred at 25°C for 2 hours. The solution is evaporated, the residue diluted with water, brought to
10 pH 1 and extracted with ethyl acetate. The extract is dried and evaporated. The residue is recrystallized from methanol to yield N-(2,4-dibenzyloxybenzoyl)glycine as a colourless solid (300 mg), m.p. 183°C. ¹H NMR (CDCl₃) 4.15(2H,d), 5.05 (2H,s), 5.2(2H,s), 6.6(1H,s), 6.7(1H,d), 7.3-7.5(10H,m), 8.2(1H,d), 8.4(1H,t).

15

By proceeding in a similar manner, but using the appropriate amount of an ester precursor, there is prepared N-(4-Benzyloxy-2-(1-o-tolyloethoxy)-benzoyl)glycine, m.p. 168-169°C.

20

EXAMPLE 5 Compound AB

A solution of ethyl 3-(2,4-dibenzyloxyphenyl)-pyrazole-5-carboxylate (1.5 g) in methanol (20 mL) and THF(10 mL) is treated with aqueous 10% KOH
25 (7.5 mL) and refluxed for 2 hours. The solution is evaporated, brought to pH 1 with 1M HCl, and the precipitate filtered. The solid is purified by flash chromatography on silica, eluting with dichloromethane/methanol (98:2). Fractions homogeneous in the required product are combined and evaporated. The residue is recrystallized from ethyl acetate to give 3-(2,4-
30 dibenzoyloxyphenyl)-pyrazole-5-carboxylic acid as colourless crystals (0.13 g), m.p. 218-219°C. ¹H NMR (CDCl₃) 5.1 (2H,s), 5.2(2H,s), 6.68(1H,d), 6.72(1H,s), 7.3-7.5(11H,m), 7.7(1H,d).

35

EXAMPLE 6 Compounds AC and AD

A solution of ethyl 2-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate (1.2 g) in methanol (125 mL) is treated with 10% aqueous

potassium hydroxide solution (10 mL, w/v) and the mixture is heated to reflux for 2 hours. The reaction mixture is evaporated in vacuo, stirred with 1N hydrochloric acid (50 mL) and extracted with ethyl acetate (100 mL). The organic phase is dried over magnesium sulphate, evaporated in vacuo, and
5 the residue crystallised from a mixture of ethyl acetate and cyclohexane to give 2-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid (0.55 g), m.p. 207-208°C. [Elemental analysis:- C,75.2; H,5.20; N,7.30%. Calculated:- C,75.0; H,5.20; N,7.30%].

10 By proceeding in a similar manner but replacing the ethyl 2-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate by ethyl 1-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate there is prepared 1-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid hydrate, m.p. 141-146°C. [Elemental analysis:- C,74.2; H,5.13; N,7.10%. Calculated for
15 $C_{24}H_{20}N_2O_3 \cdot 0.2H_2O$:- C,74.3; H,5.26; N,7.20%].

EXAMPLE 7 Compound AE

A suspension of 2,4-dibenzyloxybenzonitrile (1.89 g) in dry toluene (200
20 mL) is treated with trimethyltin chloride (6 g) and sodium azide (2 g). The mixture is stirred at reflux for 3 days, evaporated and the residue partitioned between dichloromethane (250 mL) and water (165 mL). The organic phase is washed with brine (165 mL), dried over magnesium sulphate and evaporated. The resulting semi-solid is triturated with ether (30 mL) and the solid washed
25 with pentane and crystallised from a mixture of ethyl acetate and pentane to give 5-(2,4-dibenzyloxyphenyl)-2H-tetrazole hydrate (1.55 g) as a white solid, m.p. 178-179°C. [Elemental analysis:- C,69.7; H,5.02; N,15.4%. Calculated for $C_{21}H_{18}N_4O_2 \cdot 0.2H_2O$:- C,68.68; H,5.12; N,15.48%].

30 EXAMPLE 8 Compounds AF and AG

A mixture of methyl (Z)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)-acrylate (1.6 g) and aqueous potassium hydroxide in methanol (35 mL, 15%) is stirred and heated at 60°C for 6 hours. The reaction mixture is diluted with
35 water (20 mL), acidified to pH 1 by addition of concentrated hydrochloric acid and the resulting solid filtered. Recrystallisation from ethyl acetate gives

(Z)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid (0.72 g) as a white solid, m.p. 212-213°C. [Elemental analysis:- C,75.2; H,5.28; N,2.83%. Calculated :- C,75.1; H,5.26; N,2.92%].

- 5 By proceeding in a similar manner but replacing the methyl
Z-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylate by the appropriate
quantity of methyl (E)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylate there
is prepared (E)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid, m.p.
149-150°C. [Elemental analysis:- C,74.8; H,5.24; N,2.84%. Calculated :-
10 C,75.1; H,5.26; N,2.92%].

EXAMPLE 9 Compound AH

- A mixture of ethyl (E)-2-benzyl-3-(2,4-dibenzyloxyphenyl)acrylate (1.5 g)
15 and aqueous potassium hydroxide in methanol (50 mL, 15%) is stirred and
heated at reflux for 3 hours. The reaction mixture is diluted with water (30 mL)
plus 3 N potassium hydroxide solution (5 mL), acidified to pH 1 by addition of
concentrated hydrochloric acid and the resulting solid filtered. Recrystallisation
from a mixture of ethyl acetate and pentane gives E-2-benzyl-3-(2,4-
20 dibenzyloxyphenyl)acrylic acid (1.13 g) as a white solid, m.p. 175-176°C.
[Elemental analysis:- C,79.7; H,5.72%. Calculated :- C,80.0; H,5.82%].

EXAMPLE 10 Compound AI

- 25 A solution of methyl (E)-3-(2,4-dibenzyloxyphenyl)-2-cyanoacrylate
(400 mg) in tetrahydrofuran (20 mL) containing 0.5 N aqueous lithium
hydroxide solution (4 mL) is stirred at ambient temperature for 2 hours. The
reaction is concentrated under reduced pressure and the residue taken up in
30 water. This solution is acidified to pH 1 and extracted with diethyl ether. The
organic layer is dried over magnesium sulphate, filtered and concentrated
under reduced pressure to leave (E)-3-(2,4-dibenzyloxyphenyl)-2-cyano-
acrylic acid hydrate as a yellow solid (30 mg). [Elemental analysis:- C, 71.43;
H, 5.25; N, 3.47%; Calculated for $C_{24}H_{19}NO_4 \cdot H_2O$:- C, 71.40; H, 4.81; N,
35 3.72%].

EXAMPLE 11 Compound AJ

A solution of ethyl 3-(2,4-dibenzyloxyphenyl)isoxazole-5-carboxylate (2.2 g) in hot methanol (100 mL) is treated with 10% aqueous potassium hydroxide solution (10 mL w/v) and the mixture is heated to reflux for 2 hours.

5 The reaction mixture is evaporated to low bulk, stirred with 1 N hydrochloric acid (50 mL) for 15 minutes and filtered. The resulting solid is washed with water, dried and crystallised from ethyl acetate to give 3-(2,4-dibenzyloxyphenyl)isoxazole-5-carboxylic acid (1 g), in the form of a white solid, m.p. 192-193°C. [Elemental analysis:- C,71.7; H,4.67; N,3.34%. Calculated:- C,71.8;

10 H,4.77; N,3.49%].

EXAMPLE 12 Compound AK

A solution of ethyl 4-(2,4-dibenzyloxyphenyl)-2,4-dioxobutanoate (2 g) in

15 methanol (100 mL) is treated with 10% aqueous potassium hydroxide (10 mL) and the mixture stirred at room temperature for 18 hours. The reaction mixture is evaporated to low bulk and the residue partitioned between ethyl acetate (100 mL) and 1N hydrochloric acid (50 mL). The organic phase is washed with water, dried over magnesium sulphate and evaporated in vacuo. The residue

20 isocrystallised from a mixture of ethyl acetate and cyclohexane to give 4-(2,4-dibenzyloxyphenyl)-2,4-dioxobutanoic acid (0.6 g), in the form of a white solid, m.p. 160-164°C. [Elemental analysis:- C,71.3; H,4.93%. Calculated:- C,71.3; H,4.99%].

25 EXAMPLE 13 Compound AL

A mixture of 2,4-dibenzyloxybenzaldehyde (2 g), hydroxylamine hydrochloride (0.45 g) and pyridine (0.53 mL) in ethanol (20 mL) is heated at reflux for 1 hour. The solvent is then removed in vacuo, the residue dissolved

30 in ethyl acetate and the solution washed with 1 N hydrochloric acid then with brine. Removal of the solvent in vacuo gives a beige coloured solid which crystallises from a mixture of ether and petroleum ether to give 2,4-dibenzyloxybenzaldehyde oxime in the form of a white solid, m.p., 130°C. [Elemental analysis:- C,76.1; H,5.75; N,4.28%. Calculated:- C,75.7; H,5.74;

35 N,4.20%].

EXAMPLE 14 Compound AM

A mixture of 2,4-dihydroxybenzaldehyde (17.5 g), potassium carbonate (42 g) and benzyl bromide (35 mL) in dimethyl formamide (100 mL) is stirred at room temperature for 18 hours. The reaction mixture is poured into vigorously stirred ice-water (300 mL) and stirring is continued for 30 minutes. The resulting brown solid is filtered and dried to give 2,4-dibenzyloxybenzaldehyde (28.5 g), m.p. 80-81°C. [Elemental analysis:- C,79.5; H,6.10%. Calculated:- C,79.2; H,5.70%].

10 EXAMPLE 15 Compound AN

A solution of 2,4-dibenzyloxybenzaldehyde (2.26 g) in methanol (20 mL) is treated portionwise with sodium borohydride (0.268 g) over 5 minutes. Water (5 mL) is added dropwise to the reaction mixture. The resulting white solid is filtered, dried and crystallised from a mixture of ether and petroleum ether to give 2,4-dibenzyloxybenzyl alcohol as a white solid, m.p. 87°C. [Elemental analysis:- C,79.0; H,6.42%. Calculated:- C,78.7; H,6.29%].

20 EXAMPLE 16 Compound AO

A mixture of 2,4-dibenzyloxybenzaldehyde (1.59 g), diammonium hydrogen phosphate (3.5 g) and *n*-propyl nitrate (15 mL) in glacial acetic acid (5 mL) is refluxed for 15 hours. The reaction mixture is evaporated to dryness in vacuo and the residue is stirred with water (100 mL). The insoluble material is filtered and dried at 50°C to yield 2,4-dibenzyloxybenzonitrile as a tan coloured solid (0.6 g, 40%), m.p. 100°C. [Elemental analysis:- C,79.8; H,5.47; N,4.48%. Calculated:- C,80.0; H,5.43; N,4.44%].

30 EXAMPLE 17 Compound AP

Benzyl alcohol (5.4 g) is added slowly to a stirred suspension of sodium hydride (2 g, 60% dispersion in mineral oil) in dry dimethylformamide (100 mL). The mixture is stirred for 30 minutes then 2,4-difluoronitrobenzene (3.2 g) is added in one portion. The resulting black solution is stirred at room temperature for 18 hours, evaporated and the residue partitioned between ether (200 mL) and water (100 mL). The organic phase is washed thoroughly with water, dried and evaporated. The residue is triturated with hot ethanol

(50 mL) and the mixture cooled in an ice-bath. The insoluble material is purified by filtration through a bed of silica eluting with a mixture of dichloromethane and petroleum ether, b.p.40-60°C, (1:1 v/v). The filtrate is evaporated and the solid crystallised from ethanol to give 2,4-dibenzyloxy-nitrobenzene (2.3 g) as a cream coloured solid, m.p. 101-103°C. [Elemental analysis:- C,71.4; H,5.07; N,4.09%. Calculated:- C,71.6; H,5.11; N,4.18%].

EXAMPLE 18 Compound AQ

10 A mixture of 2,4-dihydroxyacetophenone (15.2 g), potassium carbonate (30.4 g), potassium iodide (0.5 g) and benzyl bromide (37.6 g) in dimethyl formamide (200 mL) is stirred at room temperature for 18 hours. The reaction mixture is filtered and the insoluble material washed with a little dimethyl formamide. The combined filtrate plus washings are evaporated to dryness and the residue partitioned between ether and water. The organic phase is
15 dried over magnesium sulphate, evaporated and the residual pink-red coloured solid crystallised from cyclohexane to give 2,4-dibenzyloxyacetophenone (28.4 g) as a pale pink coloured powder, m.p. 64-65°C. [Elemental analysis:- C,79.5; H,5.94%. Calculated:- C,79.5; H,6.07%].

20

EXAMPLE 19 Compound AR

Concentrated sulphuric acid (2-3 drops) is added to a stirred solution of dibenzyloxybenzaldehyde (1.5 g) in methanol (50 mL), resulting in a thick
25 precipitate. The mixture is stirred for 5 minutes then hydrogen peroxide (1 g; 27.5% in water) is added in one portion and stirring is continued for 2 hours. The resulting brown solution is treated with 10% aqueous sodium sulphite solution (10 mL) then evaporated to low bulk. The residue is treated with water (50 mL), the solid washed with water, dried and crystallised from cyclohexane,
30 with charcoal treatment, to give 2,4-dibenzyloxyphenol (0.3 g) as a white powder, m.p. 82-83°C. [Elemental analysis:- C,78.2; H,5.79%. Calculated:- C,78.4; H,5.92%].

EXAMPLE 20 Compound AS

35

A mixture of 2,4-dibenzyloxyacetophenone (6.6 g), hydroxylamine hydrochloride (2.8 g) and pyridine (10 mL) in ethanol (100 mL) is heated at

reflux for 1 hour. The clear solution is cooled in an ice-bath and the solid filtered. The solid is washed with ice-cold ethanol then with ether to give 2,4-dibenzyloxyacetophenone oxime (5.4 g) as a white solid, m.p. 150°C. [Elemental analysis:- C,76.3; H,6.10; N,4.02%. Calculated:- C,76.1; H,6.09; N,4.03%]

EXAMPLE 21 Compound AT

A stirred solution of benzyl 2-benzyloxy-4-phenylethynylbenzoate (2 g) and 1 N sodium hydroxide (5 mL) in dioxan (80 mL) is heated at reflux for 1 hour. The reaction mixture is evaporated, the residue dissolved in water (6 mL) and the solution washed twice with ethyl acetate (20 mL). The pH of the aqueous phase is adjusted to pH 1 by addition of 2 N hydrochloric acid and resulting solid is filtered and washed well with water to give 2-benzyloxy-4-phenylethynylbenzoic acid (0.6 g), m.p. 125-127°C. [Elemental analysis:- C,80.2; H,4.86%. Calculated:- C,80.5; H,4.91%].

EXAMPLE 22 Compound AU

Ethyl (2,4-dibenzyloxyphenoxy)acetate (3.9 g) is added in small portions to a stirred suspension of lithium aluminium hydride (0.21 g) in dry tetrahydrofuran (100 mL) with ice-cooling. The mixture is allowed to warm to room temperature and stirred for a further 30 minutes. The reaction mixture is cooled in an ice-bath and water is added dropwise. After stirring for 1 hour the mixture is filtered through hyflo and the insolubles washed with tetrahydrofuran. Evaporation of the combined filtrate plus washings and crystallisation of the resulting solid from petroleum ether (bp 40-60°C) containing a little ethyl acetate gives 2-(2,4-dibenzyloxy-phenoxy)ethanol hemihydrate (1.8 g) as a white solid, m.p. 47-52°C. [Elemental analysis:- C,73.3; H,6.25%. Calculated for $C_{22}H_{22}O_4 \cdot 0.5H_2O$:- C,73.5; H,6.45%].

EXAMPLE 23 Compound AV

A mixture of ethyl (2,4-dibenzyloxyphenoxy)acetate (2 g), concentrated ammonia (25 mL) and methanol (50 mL) is stirred at room temperature for 18 hours. The reaction mixture is diluted with water (50 mL) then evaporated to low bulk and filtered. The solid is washed with water and dried to give

2-(2,4-dibenzyloxyphenoxy)acetamide (1.1 g) as a white powder, m.p. 120°C. [Elemental analysis:- C,73.1; H,5.78; N,3.79%. Calculated :- C,72.7; H,5.83; N,3.85%].

5 EXAMPLE 24 Compounds AW-AZ, BA-BZ, and CA-CQ

- (i) A mixture of methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate (11.6 g), sodium hydroxide (60 mL; 1 N) and 1,4-dioxane is stirred at ambient temperature for 18 hours. The reaction mixture is then concentrated to dryness and dissolved in water (50 mL), acidified to pH 1 by treatment with concentrated hydrochloric acid, and extracted with chloroform. The organic layer is dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave an oil. Trituration with a mixture of pentane and diethyl ether (1:1 v/v) and recrystallisation of the resulting solid with a mixture of diethyl ether and pentane gives (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoic acid (5.1 g), in the form of a white solid, m.p. 127-129°C. [Elemental analysis:- C,76.9;H,5.94%; Calculated:- C,76.98;H,5.92%]. (ii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2-chlorophenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2-chlorophenyl)-5-oxopentanoic acid in the form of a white solid after recrystallisation from ethyl acetate and pentane, m.p. 131-132°C. [Elemental analysis:- C,70.8;H,5.17;Cl,8.7%; Calculated; C,70.5;H,5.18;Cl,8.67%]. (iii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2-methoxyphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1 hour, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2-methoxyphenyl)-5-oxopentanoic acid in the form of a white solid after trituration with diisopropyl ether and pentane and recrystallisation from ethyl acetate, m.p. 143-144°C. [Elemental analysis:- C,74.3;H,6.10%. Calculated:- C,74.24;H,6.10%]. (iv) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1.5 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2-methylphenyl)-5-

- oxopentanoic acid in the form of a pale yellow solid after recrystallisation from a mixture of ethyl acetate and pentane, m.p. 121-122°C. [Elemental analysis:- C,77.3;H,6.28%. Calculated:- C,77.3; H,6.23%]. (v) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2-ethylphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 2.5 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2-ethylphenyl)-5-oxopentanoic acid in the form of a white solid after recrystallisation from a mixture of ethyl acetate and pentane, m.p. 85-86°C. [Elemental analysis:- C,77.0;H,6.38%; Calculated:- C,77.58;H,6.51%]. (vi) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2-chloro-6-fluorophenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1 hour, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2-chloro-6-fluorophenyl)-5-oxopentanoic acid in the form of a white solid after trituration with diisopropyl ether/pentane and recrystallisation from ethyl acetate/pentane, m.p. 95-96°C. [Elemental analysis:- C,67.5; H,4.72;Cl,8.20%. Calculated:- C,67.53;H,4.72; Cl,8.31%]. (vii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2,6-dichlorophenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 30 minutes, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2,6-dichlorophenyl)-5-oxopentanoic acid in the form of a white solid after trituration with a mixture of diisopropyl ether and pentane and recrystallisation from a mixture of ethyl acetate and pentane, m.p. 113-114°C. [Elemental analysis:- C,64.6;H,4.43; Cl,15.70%. Calculated:-C,65.02;H,4.55; Cl,15.99%]. (viii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(4-methoxyphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1.5 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(4-methoxyphenyl)-5-oxopentanoic acid in the form of a white solid after trituration with pentane, m.p. 74°C. [Elemental analysis:- C 74.3;H 5.95%. Calculated:- C 74.24;H,5.98%]. (ix) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(3,4-dichloro-

phenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 3 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(3,4-dichlorophenyl)-5-oxopentanoic acid in the form of a yellow solid after trituration from pentane, m.p. 87-88°C. [Elemental analysis:-

- 5 C,64.50;H,4.51;Cl,15.60%. Calculated:- C,65.02;H,4.55;Cl,15.99%]. (x) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(4-chlorophenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1 hour, there is
- 10 prepared (RS)-5-(3-benzyloxyphenyl)-4-(4-chlorophenyl)-5-oxopentanoic acid in the form of a white solid after chromatography on silica gel using a mixture of ethyl acetate and pentane (1:1v/v) and trituration of the resulting solid with diisopropyl ether, m.p. 91°C. [Elemental analysis:- C,70.4;H,5.14; Cl,8.6%. Calculated:- C,75.5;H,5.17;Cl,8.67%]. (xi) By proceeding in a similar manner
- 15 but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(4-methylphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 3.5 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(4-methylphenyl)-5-oxopentanoic acid in the form of a white solid after
- 20 recrystallisation from a mixture of diisopropyl ether and pentane, m.p. 105-106°C. [Elemental analysis:- C,77.1;H,6.17%. Calculated:- C,77.3;H,6.23%]. (xii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(3-chlorophenyl)-5-
- 25 oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1.5 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(3-chlorophenyl)-5-oxopentanoic acid in the form of a white solid after trituration with a mixture of diethyl ether and pentane and recrystallisation from a mixture of ethyl acetate and pentane, m.p. 110-111°C. [Elemental analysis:- C,70.3;H,5.10;Cl,8.50%. Calculated:-C,70.5;H,5.18;Cl,8.67%]. (xiii) By proceeding in a similar manner
- 30 but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(3-methoxyphenyl)-5-oxopentanoate and using 15% potassium hydroxide in methanol at reflux for 1 hour, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-
- 35 (3-methoxyphenyl)-5-oxopentanoic acid in the form of a white solid after trituration with a mixture of diisopropyl ether and pentane and recrystallisation from a mixture of ethyl acetate and pentane, m.p. 94-95°C. [Elemental

- analysis:- C,73.0;H,5.93%; Calculated for $C_{25}H_{24}O_5 \cdot 0.25AcOEt$:-
C,73.22;H,6.14%]. (xiv) By proceeding in a similar manner but replacing the
methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the
appropriate quantity of methyl (RS)-5-(4-benzyloxyphenyl)-5-oxo-4-
5 phenylpentanoate, there is prepared (RS)-5-(4-benzyloxyphenyl)-5-oxo-4-
phenylpentanoic acid in the form of a white solid after trituration with pentane,
m.p. 147-149°C. [Elemental analysis:- C,77.4;H,6.06%. Calculated:- C,76.98;
H,5.92%]. (xv) By proceeding in a similar manner but replacing the methyl
(RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate
10 quantity of methyl (RS)-4-(3-benzyloxyphenyl)-4-oxo-3-phenylbutanoate, there
is prepared (RS)-4-(3-benzyloxyphenyl)-4-oxo-3-phenylbutanoic acid in the
form of a white solid after recrystallisation from a mixture of ethyl acetate and
pentane, m.p. 150-152°C. [Elemental analysis:- C,76.7;H,5.67%. Calculated:-
C,76.65; H,5.59%]. (xvi) By proceeding in a similar manner but replacing the
15 methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the
appropriate quantity of ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenyl-
hexanoate, there is prepared (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenyl-
hexanoic acid in the form of a white solid after trituration with pentane and
recrystallisation from a mixture of ethyl acetate and cyclohexane, m.p. 107-
20 109°C. [Elemental analysis:- C,77.0;H,6.22%. Calculated:- C,77.3;H,6.23%].
(xvii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-
benzyloxyphenyl)-5-oxo-4-phenylpentanoate by ethyl (RS)-7-(3-benzyloxy-
phenyl)-7-oxo-6-phenylheptanoate, there is prepared (RS)-7-(3-benzyloxy-
phenyl)-7-oxo-6-phenylheptanoic acid in the form of an orange oil after
25 chromatography on silica gel eluting with a mixture of pentane and ethyl
acetate (8:1v/v). [Elemental analysis:- C,77.2;H,6.66%; Calculated:- C,77.59;
H,6.51%]. (xviii) By proceeding in a similar manner but replacing the methyl
(RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate
quantity of ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoate,
30 there is prepared (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoic acid
in the form of an oil after chromatography on silica gel eluting with a mixture of
pentane and ethyl acetate (1:4 v/v). [Elemental analysis:- C,75.6;H,5.87%.
Calculated:- C,75.96;H,5.73%]. (xix) By proceeding in a similar manner but
replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate
35 by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-
pyridyl)pentanoate and acidifying to pH 4.5 rather than pH 1 prior to extraction,
there is prepared (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoic

acid in the form of a white solid after chromatography on silica gel eluting with a mixture of pentane and ethyl acetate (1:1 v/v) followed by trituration with pentane, m.p. 99-104°C. [Elemental analysis:- C,73.2;H,5.91; N,3.5%. Calculated:- C,73.58;H,5.64;N,3.73%]. (xx) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (3RS,4RS)-5-(3-benzyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoate, there is prepared (3RS,4RS)-5-(3-benzyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoic acid in the form of a white solid after trituration with pentane, m.p. 68-70°C. [Elemental analysis:- C,77.0; H,6.28%. Calculated:- C,77.29;H,6.22%]. (xxi) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by methyl (2RS,4RS)-5-(3-benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoate, there is prepared (2RS,4RS)-5-(3-benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoic acid in the form of a white solid after recrystallisation from a mixture of ethyl acetate and pentane, m.p. 98-100°C. [Elemental analysis:- C,77.0;H,6.13%. Calculated:- C,77.29; H,6.22%]. (xxii) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate, and carrying out the reaction at 60°C for 3 hours, there is prepared (RS)-5-(3-methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, in the form of a white solid, m.p. 108-110°C. [Elemental analysis:- C,72.7; H,6.42%; Calculated:- C,73.1;H,6.45%]. (xxiii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(2-methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate, there is prepared (RS)-5-[3-(2-methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid in the form of a white solid after trituration with a mixture of pentane, ethyl acetate and acetic acid (72/28/1) and trituration with pentane, m.p. 111-113°C. [NMR(CDCl₃):- 2.1-2.5(4H,m),3.9(3H,s), 4.6(1H,t),5.1(2H,s),6.9-7.6(13H)]. (xxiv) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3,4-methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate, there is prepared (RS)-5-[3-(3,4-methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid, m.p. 92-94°C. [Elemental analysis:- C,71.4;H,5.46%. Calculated:- C,71.8;H,5.29%]. (xxv) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the

- appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-[3-(2-methylpropoxy)phenyl]-5-oxopentanoate, and carrying out the reaction for 18 hours, there is prepared (RS)-4-(2-methylphenyl)-5-[3-(2-methylpropoxy)phenyl]-5-oxopentanoic acid, in the form of a white solid after
- 5 recrystallisation from a mixture of ethyl acetate and hexane, m.p. 96-98°C. [Elemental analysis:- C,74.0;H,7.3%; Calculated:- C,74.5;H,7.39%]. (xxvi) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(4-chlorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-
- 10 oxopentanoate, there is prepared (RS)-5-[3-(4-chlorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 131-133°C. [Elemental analysis; C,71.2;H,5.48%. Calculated:- C,71.0;H,5.48%]. (xxvii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-
- 15 5-(3-cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 104-105°C. [Elemental analysis:- C,75.7;H,7.60%. Calculated:- C,75.8; H,7.42%]. (xxviii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate
- 20 quantity of methyl (RS)-5-[3-(3-thienylmethoxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(3-thienylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid hydrate, m.p. 94-98°C. [Elemental analysis:- C,67.1;H,5.60%. Calculated for $C_{23}H_{22}O_4S \cdot H_2O$:- C,67.1;H,5.85%]. (xxix) By proceeding in a similar manner but replacing the
- 25 methyl (RS)-5-(3-benzyloxy-phenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(2-fluorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(2-fluorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 108-110°C. [Elemental analysis:- C, 73.5;H,5.65%. Calculated:- C,73.9;H,5.70%]. (xxx) By proceeding in a similar manner but
- 30 replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(2-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(2-furylmethoxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 92-94°C. [Elemental
- 35 analysis:- C,72.9;H,5.81%. Calculated:- C,73.0;H,5.86%]. (xxxi) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3-

furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(3-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 83-85°C. [Elemental analysis:- C,72.6;H,5.66%. Calculated:- C,73.0; H,5.86%]. (xxxii) By proceeding in a similar manner but replacing the methyl

5 (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3-chlorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(3-chlorobenzyl-oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 103-105°C. [Elemental analysis:- C,71.1;H,5.45%. Calculated:- C,71;H,5.48%]. (xxxiii) By proceeding in a

10 similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 132-133°C. [Elemental analysis; C,74.4;H,6.25%. Calculated:-

15 C,74.4;H,6.25%]. (xxxiv) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-oxo-5-(3-(2-pyridylmethoxy)phenyl)pentanoate, there is prepared (RS)-4-(2-methylphenyl)-5-oxo-5-(3-(2-pyridylmethoxy)phenyl)pentanoic acid, m.p. 152-154°C. [Elemental

20 analysis:- C,74.0;H,5.95%. Calculated:- C,74.0;H,5.95%]. (xxxv) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]-pentanoate, there is prepared (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-

25 thienylmethoxy)phenyl]-pentanoic acid, m.p. 96-98°C. [Elemental analysis:- C,70.1; H,5.67%. Calculated:- C,70.0;H,5.62%]. (xxxvi) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(2-methyl-3-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate, and carrying out the

30 reaction for 18 hours, there is prepared (RS)-5-(2-methyl-3-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, in the form of a white solid after recrystallisation from ethyl acetate, m.p. 177-178°C. [Elemental analysis:- C,77.7; H,6.54%; Calculated:- C,77.6;H,6.51%]. (xxxvii) By proceeding in a

35 similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3,5-dibenzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate, and using a solution of potassium hydroxide in methanol (15% w/v) at reflux for 1.5 hours, and

- subjecting to flash chromatography on silica gel, eluting with a mixture of ethyl acetate and pentane (2:3 v/v), there is prepared (RS)-5-(3,5-dibenzyloxyphenyl)-4-(2-methylphenyl)-5-oxo-pentanoic acid, in the form of a pale yellow oil. [Elemental analysis:- C,76.6;H,6.08%; Calculated for:
- 5 C₃₂H₃₀O₅·0.25AcOEt:- C,76.7;H,6.24%]. (xxxviii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared
- 10 (RS)-5-[3-(3-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 72-74°C. [Elemental analysis:- C,74.6;H,6.27%. Calculated:- C,74.6; H,6.26%]. (xxxix) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3-aminobenzyloxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(3-aminobenzyl-
- 15 oxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid in the form of an oil. [Elemental analysis:- C,73.9; H,6.34;N,3.43%. Calculated:- C,74.4;H,6.24; N,3.47%]. (xxxx) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-[3-(3-isothiazolylmethoxy)phenyl]-4-(2-methyl-
- 20 phenyl)-5-oxopentanoate, there is prepared (RS)-5-[3-(3-isothiazolyl-methoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, m.p. 102-104°C. [Elemental analysis:- C,66.8;H,5.33;N,3.44%. Calculated:- C,66.8;H,5.35; N,3.54%]. (xxxxi) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate
- 25 quantity of methyl (RS)-5-(3-benzyloxy-5-hydroxyphenyl)-4-(2,3-dimethylphenyl)-5-oxopentanoate, there is prepared (RS)-5-(3-benzyloxy-5-hydroxyphenyl)-4-(2,3-dimethylphenyl)-5-oxopentanoic acid. (xxxxii) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyl-oxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl
- 30 (RS)-5-(3-benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate, and using a solution of potassium hydroxide in methanol (15% w/v) at reflux for 1 hour, there is prepared (RS)-5-(3-benzyloxy-4-methyl-phenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, in the form of a white solid after trituration with pentane, m.p. 94-98°C. [Elemental analysis:- C,77.1; H,6.58%. Calculated:- C,77.6;H,6.51%]. (xxxxiii) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(4-
- 35

- pyridylmethoxy)phenyl]pentanoate, there is prepared (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(4-pyridylmethoxy) phenyl]pentanoic acid, m.p. 124-125°C. [Elemental analysis:-C,74.0;H,6.05;N,3.43%. Calculated:- C,74.0;H,5.95; N,3.60%]. (xxxxiv) By proceeding in a similar manner but replacing the methyl
- 5 (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-[3-(3-methyl-2-thienylmethoxy)-phenyl]-5-oxopentanoate, there is prepared (RS)-4-(2-methylphenyl)-5-[3-(3-methyl-2-thienylmethoxy)phenyl]-5-oxopentanoic acid, m.p. 88-90°C. [Elemental analysis:- C,70.3;H,5.98%; C₂₄H₂₄O₄S requires C,70.6; H,5.92%].
- 10 (xxxxv) By proceeding in a similar manner but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(3-pyridylmethoxy)phenyl]pentanoate, there is prepared (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(3-pyridylmethoxy)phenyl]pentanoic acid hydrate in the form of an oil. [Elemental
- 15 analysis:-C,72.9;H,6.44; N,3.27%; Calculated for C₂₄H₂₃NO₄·0.3H₂O :- C,72.9;H,6.03;N,3.54%]. (xxxxvi) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoate, and using a solution of potassium
- 20 hydroxide in methanol (15% w/v) at reflux for 0.75 hours, there is prepared (RS)-5-(3-benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoic acid, in the form of a white solid after trituration with diisopropyl ether, m.p. 146-148°C. [Elemental analysis:- C,77.2;H,6.5%. Calculated:- C,77.6;H,6.51%]. (xxxxvii) By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyl-
- 25 oxyphenyl)-5-oxo-4-phenylpentanoate by the appropriate quantity of methyl (RS)-5-(3-benzyloxy-6-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate, there is prepared (RS)-5-(3-benzyloxy-6-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, in the form of a yellow oil.

30 EXAMPLE 25 Compounds CR and CS

- A mixture of methyl (RS)-5-(3-benzyloxy-phenyl)-4-(3-methylpyrid-4-yl)-5-oxopentanoate (3.26 g), dioxan (50 mL), water (9 mL) and concentrated hydrochloric acid (6 mL) is stirred at 60°C for 1 hour and then evaporated to
- 35 dryness under reduced pressure. The residue is treated with saturated aqueous sodium bicarbonate solution to give a solution of pH 7 and extracted with ethyl acetate. Evaporation, followed by crystallisation from a mixture of

ethyl acetate and pentane affords (RS)-5-(3-benzyloxyphenyl)-4-(3-methylpyrid-4-yl)-5-oxopentanoic acid (1.56 g) in the form of a cream coloured solid, m.p. 55-57°C. [Elemental analysis:- C,73.7;H,5.92; N,3.41%. Calculated:- C,74.0;H,5.95;N,3.60%].

5

By proceeding in a similar manner, but replacing the methyl (RS)-5-(3-benzyloxyphenyl)-4-(3-methylpyridin-4-yl)-5-oxopentanoate by the appropriate quantity of methyl (RS)-5-[3-(1,2,4-oxadiazol-3-ylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate and carrying out the reaction for 4 hours, there is prepared, after flash chromatography on silica eluting with a mixture of dichloromethane and methanol (96.5:3.5 v/v), (RS)-5-[3-(1,2,4-oxadiazol-3-ylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid, in the form of a colourless oil.

15 EXAMPLE 26 Compounds CT-CZ and DA-DO

A stirred solution of benzyl (3-benzyloxyphenyl) ketone (10 g) in tetrahydrofuran (100 mL) cooled at 0°C is treated with potassium tert-butoxide (370 mg). After 40 minutes at 0°C, the mixture is treated with methyl acrylate (2.9 g) and stirring is continued at ambient temperature. The reaction is then concentrated under reduced pressure, partitioned between water and ethyl acetate and the organic layer is dried over magnesium sulphate. Concentration gives methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenyl-pentanoate (11.6 g), in the form of a colourless oil.

25

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared:- methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chlorophenyl)-5-oxopentanoate, in the form of a yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methoxyphenyl)-5-oxopentanoate, in the form of an orange oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methylphenyl)-5-oxo-pentanoate, in the form of a pale yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-ethylphenyl)-5-oxo-pentanoate, in the form of a pale yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chloro-6-fluorophenyl)-5-oxo-pentanoate, in the form of a yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2,6-dichlorophenyl)-5-oxo-pentanoate, in the form of a pale yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3,4-dichlorophenyl)-5-oxo-pentanoate, in the form of an orange oil; methyl (RS)-5-[3-(benzyloxy)-

35

- phenyl]-4-(4-chlorophenyl)-5-oxopentanoate, in the form of a yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(4-methylphenyl)-5-oxopentanoate, in the form of a yellow oil; methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3-chlorophenyl)-5-oxopentanoate, in the form of a white solid, m.p. 110-111°C.; methyl
- 5 (RS)-5-[3-(benzyloxy)phenyl]-4-(3-methoxyphenyl)-5-oxopentanoate, in the form of a pale yellow oil; methyl (RS)-5-(4-benzyloxyphenyl)-5-oxo-4-phenylpentanoate, in the form of a colourless solid, m.p. 74-76°C.; methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoate, in the form of a brown oil; methyl (3RS,4RS)-5-(3-benzyloxyphenyl)-3-methyl-5-oxo-4-
- 10 phenylpentanoate, in the form of an oil; methyl (2RS,4RS)-5-(3-benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoate, in the form of an oil; methyl (RS)-5-[3-(2-methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate, in the form of a colourless oil; methyl (RS)-5-[3-(3,4-methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate, in the form of an oil; methyl (RS)-5-(3-benzyloxy-2-
- 15 methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-(3,5-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-(3-benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate; and methyl (RS)-5-(3-benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoate.

20 EXAMPLE 27 Compounds DP-DS

- A stirred solution of benzyl 3-benzyloxyphenyl ketone (3.02 g) in dry tetrahydrofuran (70 mL) at -40°C is treated with potassium tert-butoxide (1.12 g), portionwise during 10 minutes. The reaction is stirred at -40°C for
- 25 1 hour and is then treated dropwise with a solution of methyl bromoacetate (1.04 g) in tetrahydrofuran (10 mL). The mixture is allowed to stir at ambient temperature for 18 hours. After this time, the reaction is concentrated to dryness and partitioned between ethyl acetate and water. The organic layer is dried over magnesium sulphate and concentrated under reduced pressure.
- 30 The residue is recrystallised from a mixture of ethyl acetate and pentane, to give methyl (RS)-4-(3-benzyloxyphenyl)-4-oxo-3-phenylbutanoate (1.4 g), m.p. 88-89°C.

- By proceeding in a similar manner but using the appropriate quantities
- 35 of the corresponding starting materials, there are prepared :
- ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhexanoate, in the form of a light green oil; ethyl (RS)-7-(3-benzyloxyphenyl)-7-oxo-6-phenylheptanoate, in

the form of a colourless oil; and ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoate, in the form of an oil.

EXAMPLE 28 Compounds DT-DZ and EA-EL

5

A stirred solution of methyl (RS)-5-[3-(hydroxy)phenyl]-4-(2-methylphenyl)-5-oxo-pentanoate (1.68 g) in dimethylformamide (20 mL) is treated with 60% sodium hydride in oil (237 mg). After 15 minutes the mixture is treated with 4-chlorobenzyl chloride (953 mg) and heated at 60°C for 1 hour.

10 The reaction mixture is evaporated, the residue is dissolved in diethyl ether, washed with water, dried, and evaporated, to give methyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate (2.05 g), in the form of a colourless oil. [NMR(CDCI₃):- 2.0-2.5(4H,m), 2.5(3H,s), 3.65(3H,s), 4.8(1H,m), 5.0(2H,q), 7.0-7.5(12H)].

15

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared :

ethyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxo-pentanoate; methyl (RS)-5-(3-cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-
20 5-oxopentanoate; methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(3-thienylmethoxy)phenyl]pentanoate; methyl (RS)-5-[3-(2-fluorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-[3-(2-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-[3-(3-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-[3-(3-chlorobenzyloxy)-
25 phenyl]-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-[3-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-pyridylmethoxy)phenyl]pentanoate; methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]pentanoate; methyl (RS)-5-[3-(3-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
30 methyl (RS)-5-[3-(3-isothiazolylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)-pentanoate; methyl (RS)-5-[3-(4-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate; methyl (RS)-5-[3-(3-nitrobenzyloxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate; methyl (RS)-5-[3-(3-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate; methyl (RS)-5-(3-methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate; methyl (RS)-5-(3-isopropoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate; and methyl (RS)-5-[3-(1,2,4-oxadiazol-3-ylmethoxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate.

EXAMPLE 29 Compound EM

A solution of triphenylphosphine (1.13 g) in tetrahydrofuran (30 mL) is
5 treated with diisopropyl azodicarboxylate (0.84 mL) at 0-5°C. The suspension
is then treated with methyl (RS)-5-(3-hydroxyphenyl)-5-oxo-4-(2-methyl-
phenyl)pentanoate (671 mg) and 3-methyl-2-thienylmethanol (275 mg), and
stirred at 5°C for 15 minutes. The solution is evaporated and the residue is
10 subjected to flash chromatography on silica gel, eluting with a mixture of
pentane and ethyl acetate (5:1v/v), to give methyl (RS)-5-[3-(3-methyl-2-
thienylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate (350 mg), in the
form of an oil. [NMR(CDCl₃):- 2.0-2.5(4H,m), 2.25(3H,s), 2.5(3H,s), 3.65(3H,s),
4.8(1H,m), 5.1(2H,q), 6.85-7.5(10H)].

15 EXAMPLE 30 Compound EN

A solution of methyl (RS)-5-[3-(3-nitro-benzyloxy)phenyl]-5-oxo-4-(2-
methylphenyl)pentanoate (1.2 g) in methyl acetate (30 mL) is treated with
palladium on carbon catalyst (5%; 200 mg) and shaken under hydrogen at
20 atmospheric pressure and room temperature for 3 hours. The mixture is filtered
and evaporated giving methyl (RS)-5-[3-(3-aminobenzyloxy)- phenyl]-5-oxo-4-
(2-methylphenyl)pentanoate (1.3 g) in the form of a brown oil.

EXAMPLE 31 Compound EO

25 A solution of [3-benzyloxy-5-(4-methoxybenzyloxy)] (2-methylbenzyl)
ketone (18.8 g) in dry tetrahydrofuran (200 mL) is treated with potassium tert-
butoxide and stirred at ambient temperature for 15 minutes. It is then treated
with methyl acrylate (4 g) and stirring is continued for 2 hours. After this time,
30 the mixture is concentrated and partitioned between ethyl acetate (250 mL)
and water (150 mL). The organic layer is dried over magnesium sulphate and
concentrated. The residue is chromatographed on silica gel, eluting with ethyl
acetate/pentane (1/4), to give methyl (RS)-5-[3-benzyloxy-5-(4-methoxy-
benzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate (15.3 g), in the form of
35 a pale yellow oil.

EXAMPLE 32 Compound EP

A solution of methyl (RS)-5-[3-benzyloxy-5-(4-methoxybenzyloxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate (0.54 g) in dichloromethane (10 mL), pH 7.3 phosphate buffer and tert-butanol (0.1 mL) is treated at 0°C with 2,3-dichloro-5,6-dicyanobenzoquinone (0.34 g) and stirred at ambient temperature for 48 hours. It is then diluted with water (100 mL) and extracted with dichloromethane (3 x 30 mL). The combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate and concentrated. The resulting oil (0.4 g) is dissolved in a solution of potassium hydroxide in methanol (20 mL; 15%w/v) and heated at reflux for 5 hours and is then allowed to cool. The mixture is concentrated, diluted with water (30 mL) and acidified to pH1 by treatment with concentrated hydrochloric acid. The mixture is extracted with ethyl acetate (3 x 50 mL) and the combined organic extracts are washed with brine and dried over magnesium sulphate, and concentrated. The resulting residue is chromatographed on silica gel, eluting with a mixture of ethyl acetate and pentane (1:1 v/v). The resulting oil is triturated with a mixture of diisopropyl ether and pentane, to give (RS)-5-(3-benzyloxy-5-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid in the form of a white solid (80 mg), m.p. 116-117°C. [Elemental analysis:- C,73.80; H,5.90%; Calculated:- C,74.24;H,5.98%].

EXAMPLE 33 Compounds EQ and ER

A mixture of 1-(3-benzyloxyphenyl)-4-cyano-2-phenylbutan-1-one (2.2 g), toluene (80 mL), trimethyltin chloride (6.17 g) and sodium azide is stirred at 115°C for 30 hours. The cooled reaction mixture is filtered and the insoluble material is washed with a little ethyl acetate. The combined filtrate plus washings are evaporated and the resulting orange oily solid (3 g) is subjected to flash chromatography on silica gel, eluting with a mixture of dichloromethane and methanol (29:1 v/v), followed by crystallisation from a mixture of ethyl acetate and cyclohexane, to give 5-[4-(3-benzyloxyphenyl)-4-oxo-3-phenylbutyl]-1H-tetrazole (0.86 g), in the form of a white solid, m.p. 136-138°C. [Elemental analysis:- C,72.5;H,5.63; N,14.0%; Calculated:- C,72.3;H,5.56; N,14.1%].

By proceeding in a similar manner, but replacing the 1-(3-benzyloxyphenyl)-4-cyano-2-phenylbutan-1-one by the appropriate quantity of

1-(3-benzyloxyphenyl)-3-cyano-2-(2-methylphenyl)propan-1-one, there is prepared 5-[3-(3-benzyloxyphenyl)-3-oxo-2-(2-methylphenyl)propyl]-1H-tetrazole, in the form of an oil. [Elemental analysis:- C,72.6;H,5.61; N,13.9%; Calculated:- C,72.4;H,5.53;N,14.07%].

5

EXAMPLE 34 Compounds ES and ET

A mixture of (RS)-6-(3-benzyloxyphenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide (1.6 g), aqueous sodium hydroxide solution (10 mL; 1 N) and 1,4-dioxane (20 mL) is heated at reflux for 3 hours. The cooled reaction mixture is partitioned between ethyl acetate (50 mL) and water (50 mL), and the aqueous phase is acidified with dilute hydrochloric acid (2 N) and extracted with ethyl acetate (2 x 50 mL). The combined extracts are washed with water, dried over magnesium sulphate and evaporated, and the residual brown oil is subjected to chromatography on silica gel, eluting with a mixture of dichloromethane and methanol (9:1 v/v) to remove starting material, then eluting with a mixture of dichloromethane and methanol (4:2 v/v), to give (RS)-4-(3-benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylsulphonic acid hydrate (0.95 g), in the form of a brown oil. [Elemental analysis:- C,65.1;H,6.37%; Calculated for $C_{24}H_{24}O_5S \cdot H_2O$:- C,65.1;H,5.92%].

20

By proceeding in a similar manner, but replacing the (RS)-6-(3-benzyloxyphenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide by the appropriate quantity of (RS)-6-[3-(3-thienylmethoxy)phenyl]-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide, there is prepared (RS)-4-[3-(3-thienylmethoxy)phenyl]-3-(2-methylphenyl)-4-oxobutylsulphonic acid, in the form of a pale brown hygroscopic solid, which turns into a brown oil on contact with the atmosphere. [NMR $\{(CD_3)_2SO\}$:- 1.4-2.5(4H,m), 2.41(3H,s), 5.1(2H,q),5.3(1H,t),6.96-7.6(10H,m)].

30

EXAMPLE 35 Compounds EU and EV

A stirred solution of benzyl 3-benzyloxy-phenyl ketone (4.6 g) in tetrahydrofuran (100 mL) is treated with potassium tert-butoxide (1.9 g). After stirring for 20 minutes it is treated with phenyl vinylsulphonate (3.04 g), dropwise, and stirring is continued for a further 3 days. The reaction mixture is then concentrated under reduced pressure and partitioned between water and

35

ethyl acetate. The organic layer is dried over magnesium sulphate and evaporated and the resulting residue is triturated with a mixture of ethyl acetate and petroleum ether (1:4v/v), to give (RS)-6-(3-benzyloxyphenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide, in the form of a white solid
5 (1.6 g), m.p. 257-259°C.

By proceeding in a similar manner, but replacing the benzyl 3-benzyloxyphenyl ketone by the appropriate quantity of 1-[3-(3-thienylmethoxy)phenyl]-2-(2-methylphenyl)ethan-1-one, there is prepared (RS)-6-[3-
10 (3-thienylmethoxy)-phenyl]-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide, in the form of a cream coloured solid, m.p. 56-57°C.

EXAMPLE 36 Compound EW

15 By proceeding in a manner similar to that described hereinbefore in Example 34 but replacing the (RS)-6-(3-benzyloxyphenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide by the appropriate quantity of diethyl (RS)-4-(3-benzyloxy)phenyl-3-(2-methylphenyl)-4-oxobutylphosphonate, and heating at reflux for 8 hours, there is prepared ethyl (RS)-4-(3-benzyloxy-
20 phenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate (0.38 g), in the form of a brown gum.

EXAMPLE 37 Compound EX

25 By proceeding in a manner similar to that described hereinbefore in Example 35, but replacing the benzyl 3-benzyloxyphenyl ketone by the appropriate quantity of 1-[3-(3-thienyl-methoxy)phenyl]-2-(2-methylphenyl)-ethan-1-one and the phenyl vinylsulphonate by the appropriate quantity of diethyl vinylphosphonate, there is prepared diethyl (RS)-4-(3-benzyloxy-
30 phenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate, in the form of a colourless oil.

EXAMPLE 38 Compounds EY-EZ and FA-FL

35 A mixture of ethyl (RS)-4-(3-benzyloxy-phenoxy)-4-phenylbutanoate (1 g), aqueous potassium hydroxide solution (5 mL; 10% w/v) and methanol

(50 mL) is stirred at reflux for 30 minutes. The reaction mixture is then evaporated to dryness and the residue is partitioned between ethyl acetate (100 mL) and dilute hydrochloric acid (50 mL; 1 N). The organic layer is washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure to give a brown oil, which slowly crystallises.

Recrystallisation from cyclohexane gives (RS)-4-(3-benzyloxyphenoxy)-4-phenylbutanoic acid (0.55 g), in the form of a beige solid, m.p. 100-104°C. [Elemental analysis:- C,76.3;H,6.21%; Calculated:- C,76.2;H,6.12%].

- 10 By proceeding in a similar manner, but replacing the ethyl (RS)-4-(3-benzyloxyphenoxy)-4-phenylbutanoate with the appropriate esters, there are prepared:- (RS)-4-(2-carboxy-5-benzyloxyphenoxy)-4-phenylbutanoic acid, in the form of a white solid after recrystallisation from ethyl acetate/cyclohexane, m.p. 124-126°C. [Elemental analysis:- C,71.1;H,5.44%; Calculated:- C,70.9;H,5.46%]; (RS)-4-[2-(N-ethylcarbamoyl)-5-benzyloxyphenoxy]-4-phenylbutanoic acid in the form of a white solid after recrystallisation from ethyl acetate/cyclohexane, m.p. 137-139°C. [Elemental analysis:- C,72.2;H,6.33;N,3.29%; Calculated:- C,72.0;H,6.28; N,3.23%]; (RS)-4-(2-acetyl-5-benzyloxyphenoxy)-4-phenylbutanoic acid in the form of a white solid after recrystallisation from ethyl acetate/cyclohexane, m.p. 112-115°C. [Elemental analysis:- C,74.1; H,6.00%; Calculated:- C,74.2;H,5.98%]; (RS)-4-[3-(3-thienylmethoxy)phenoxy]-4-phenylbutanoic acid in the form of a white solid after recrystallisation from cyclohexane, m.p. 105-107°C. [Elemental analysis:- C,67.2;H,5.44%; Calculated:- C,67.6;H 5.44%]; (RS)-4-(3-benzyloxy)phenoxy-4-(2-methylphenyl)butanoic acid in the form of a yellow oil [Elemental analysis:- C,76.5;H,6.95%; Calculated:- C,76.6; H,6.43%]; (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid after recrystallisation from cyclohexane/ethyl acetate, m.p. 116-121°C. [Elemental analysis:- C,68.4;H,5.69%; Calculated:- C,67.9;H,5.70%]; (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-phenylbutanoic acid in the form of a white solid after recrystallisation from cyclohexane/ethyl acetate, m.p. 134-135°C. [Elemental analysis:- C,67.5; H,5.37%; Calculated:- C,67.3;H 5.40%]; (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)phenoxy]-4-phenylbutanoic acid in the form of a colourless oil [Elemental analysis:- C,67.5;H,5.84;N,3.22%; Calculated:- C,68.0;H,5.91;N,3.31%]; (RS)-4-[2-carbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoic acid in the form of a white solid after recrystallisation from a mixture of ethyl acetate and acetonitrile, m.p. 170-

- 171°C. [Elemental analysis:- C,71.1;H,5.67;N,3.36%; Calculated:- C,71.1;H,5.72;N,3.44%]; (RS)-4-[2-cyano-5-benzyloxyphenoxy]-4-phenylbutanoic acid in the form of a white solid after recrystallisation from ethyl acetate/cyclohexane, m.p. 134-137°C. [Elemental analysis:- C,74.3; H,5.29;N,3.42%; Calculated:- C,74.4;H,5.46; N,3.61%];
- 5 (RS)-4-[2-(N,N-dimethylcarbamoyl)-5-benzyloxy-phenoxy]-4-phenylbutanoic acid in the form of a yellow semi-solid [Elemental analysis:- C,71.3; H,6.38; N,3.04%; Calculated:- C,71.3;H,6.33; N,3.20%]; (RS)-4-[2-(3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of an off-
- 10 white solid after recrystallisation from ethyl acetate/cyclohexane, m.p. 189-190°C. [Elemental analysis:- C,66.9; H,5.39;N,6.21; Calculated:- C,66.9; H,5.39; N,6.25]; and (RS)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid after recrystallisation from a mixture of ethyl acetate and cyclohexane, m.p. 167-169°C. [Elemental
- 15 analysis:- C,67.4;H,5.11; N,3.41; Calculated:- C,67.2;H,5.19;N,3.44].

EXAMPLE 39 Compound FM

- A mixture of 3-benzyloxyphenol (3 g), ethyl (RS)-4-chloro-4-
- 20 phenylbutanoate (4.53 g), potassium carbonate (4.14 g), potassium iodide (100 mg) and methyl ethyl ketone (100 mL) is heated at reflux for 40 hours. The mixture is evaporated to dryness and the residue is partitioned between ethyl acetate (200 mL) and water (200 mL). The organic layer is dried over magnesium sulphate, filtered, and concentrated in vacuo. Flash
- 25 chromatography of the resulting oil, eluting with a mixture of ethyl acetate and cyclohexane (1:19 v/v), gives ethyl (RS)-4-(3-benzyloxyphenoxy)-4-phenylbutanoate (1 g), in the form of a yellow oil.

EXAMPLE 40 Compounds FN-FW

- 30 A stirred suspension of sodium hydride (340mg) in dry dimethylformamide (100 mL) is treated portionwise with methyl 4-benzyloxy-2-hydroxybenzoate (2 g). After stirring at ambient temperature for 30 minutes, it is treated with ethyl (RS)-4-chloro-4-phenylbutanoate (1.9 g) in one portion and
- 35 stirred for 16 hours at ambient temperature, and then for two hours at 80°C. The reaction mixture is then concentrated in vacuo and the residue is partitioned between ethyl acetate (100 mL) and water (100 mL). The organic

layer is dried over magnesium sulphate, filtered and concentrated in vacuo. Flash chromatography of the resulting oil, eluting with 5-10% ethyl acetate/cyclohexane, affords ethyl (RS)-4-(2-methoxycarbonyl-5-benzyloxyphenoxy)-4-phenylbutanoate (2 g), in the form of a yellow oil.

- 5
- By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 3-benzyloxyphenol and ethyl (RS)-4-chloro-4-phenylbutanoate with ethyl (RS)-4-chloro-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[(3-benzyloxy)phenoxy]-4-(2-methylphenyl)-
- 10 butanoate. By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 3-(3-thienylmethoxy) phenol, there is prepared ethyl (RS)-4-[3-(3-thienylmethoxy)phenoxy]-4-phenylbutanoate. By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 2-hydroxy-4-(3-thienylmethoxy)acetophenone, there is prepared
- 15 ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-phenylbutanoate. By proceeding in a similar manner but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 2-hydroxy-4-benzyloxy-N-ethylbenzamide there is obtained ethyl (RS)-4-[2-(N-ethylcarbamoyl)-5-benzyloxyphenoxy]-4-phenylbutanoate. By proceeding in a similar manner but replacing methyl 4-benzyloxy-2-
- 20 hydroxybenzoate with 2-hydroxy-4-(3-thienylmethoxy)acetophenone and ethyl (RS)-4-chloro-4-phenylbutanoate with ethyl (RS)-4-chloro-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate. By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 2-hydroxy-4-(3-
- 25 pyridylmethoxy)acetophenone, there is prepared ethyl (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)phenoxy]-4-phenylbutanoate. By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 4-benzyloxy-2-hydroxybenzamide, there is prepared ethyl (RS)-4-[2-carbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoate. By proceeding in a similar manner,
- 30 but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 4-benzyloxy-2-hydroxy-N,N-dimethylbenzamide, there is prepared ethyl (RS)-4-[2-N,N-dimethylcarbamoyl-5-benzyloxyphenoxy]-4-phenylbutanoate. By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate with 4-(3-thienylmethoxy)-2-hydroxybenzamide and ethyl (RS)-4-chloro-4-phenyl-
- 35 butanoate with ethyl (RS)-4-chloro-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.

EXAMPLE 41 Compound FX

A mixture of 4-benzyloxy-2-hydroxyaceto-phenone (1.2 g), ethyl (RS)-4-chloro-4-phenyl-butanoate (1.21 g), potassium carbonate (760 mg), potassium iodide (35 mg) and dimethylformamide (30 mL) is stirred at 100°C for 8 hours. It is evaporated to dryness and the resulting residue is partitioned between ethyl acetate (200 mL) and water (200 mL). The organic layer is dried over magnesium sulphate, filtered and concentrated in vacuo. Flash chromatography of the resulting oil, eluting with 15-30% ethyl acetate/hexane, affords ethyl (RS)-4-(2-acetyl-5-benzyloxy-phenoxy)-4-phenylbutanoate (0.73 g), in the form of a yellow oil.

EXAMPLE 42 Compounds FY and FZ

15

A mixture of ethyl (RS)-4-(2-carbamoyl-5-benzyloxyphenoxy)-4-phenylbutanoate (1.9 g) and acetic anhydride (50 mL) is stirred at reflux for 2 hours. The reaction mixture is evaporated to low bulk and the residue is dissolved in ethyl acetate (50 mL). This solution is washed with aqueous sodium hydrogen carbonate and water, dried, and evaporated. Flash chromatography, eluting with a mixture of cyclohexane and dichloromethane (15:85 v/v), gives ethyl (RS)-4-(2-cyano-5-benzyloxyphenoxy)-4-phenylbutanoate (0.6 g).

By proceeding in a similar manner, but replacing ethyl (RS)-4-(2-carbamoyl-5-benzyloxy-phenoxy)-4-phenylbutanoate by ethyl (RS)-4-(2-carbamoyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.

30 EXAMPLE 43 Compound GA

A stirred suspension of sodium hydride (0.1 g of a 60% dispersion in oil) in anhydrous toluene (25 mL) is treated with a mixture of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1 g) and ethyl formate (1 mL) in anhydrous toluene (5 mL). The resulting mixture is stirred at reflux for 2 hours, during which time a yellow precipitate forms. After cooling to room temperature the mixture is filtered and the residue is washed with a little

cyclohexane and partitioned between dilute hydrochloric acid (10 mL; 1 M) and ethyl acetate (50 mL). The organic layer is dried and evaporated and the residue is dissolved in ethanol (25 mL) and treated with hydrazine hydrate (0.12 g). After standing at room temperature overnight, the mixture is
5 partitioned between water (50 mL) and ethyl acetate (50 mL). The organic layer is dried and evaporated flash chromatography, eluting with a mixture of ethyl acetate and cyclohexane (1:4v/v), to give ethyl (RS)-4-[2-(3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (0.25 g) in the form of a colourless oil.

10

EXAMPLE 44 Compounds GB-GE

(RS)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid is separated into its enantiomers, (R)-5-(3-benzoyloxyphenyl)-4-(2-methylphenyl)-
15 5-oxopentanoic acid and (S)-5-(3-benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid, by chiral HPLC, under the following conditions:- Chiralcel OD column; mobile phase of isopropanol/methanol/acetic acid/heptane (1:1:1:97 v/v); flow rate 1 mL/minute; temperature ambient; UV detection 270 nm.

20 By proceeding in a similar manner, (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid is separated into its enantiomers, (R)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid and (S)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, by chiral HPLC, under the following conditions:-
25 Chiralcel OD column; mobile phase of isopropanol/acetic acid/heptane (10:1:100 v/v); flow rate 1mL/minute; temperature ambient; UV detection 270 nm.

EXAMPLE 45 Compounds GF-GI

30

A solution of ethyl (RS)-4-[2-acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (1.1 g) in dioxan (10 mL) and sodium hydroxide (5.3 mL; 1 M) is stirred at 25°C for 45 minutes. The solution is evaporated, the residue is diluted with water, and brought to pH 6 by treatment with hydro-
35 chloric acid (2 M). The precipitate is extracted with ethyl acetate, washed with water, dried and evaporated. The residual oil is subjected to flash chromatography on silica gel, eluting with a mixture of dichloromethane and

methanol (95:5 v/v), followed by trituration with diethyl ether to give (RS)-4-[2-acetyl-5-(5-pyrimidinylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid (0.44 g) in the form of a colourless solid, m.p. 142-144°C. [NMR(CDCl₃):- 2.2-2.7(4H,m), 2.45(3H,s), 2.7 (3H,s), 4.9(2H,q), 5.5(1H,q), 6.2-7.8(7H), 8.7(2H,s), 9.2(1H,s)].

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared:- (RS)-4-(2-methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoic acid, m.p. 65-67°C; (RS)-4-(2-methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]-butanoic acid, m.p. 211-213°C; and (RS)-4-(2-methylphenyl)-4-[2-thiocarbamoyl-5-(3-thienylmethoxy)phenoxy]butanoic acid, m.p. 149-150°C.

EXAMPLE 46 Compound GJ

A solution of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (0.5 g) in ethanol (5 mL) is treated with pyridine-2-carboxaldehyde (0.12 g) and aqueous sodium hydroxide solution (0.95 g; 33% w/w) and stirred at 25°C for 90 minutes. The solution is diluted with water and then brought to pH 7 by treatment with hydrochloric acid (2 M), forming a yellow gum. This gum is extracted with ethyl acetate, and the extract is washed with water, dried and evaporated. The resulting residue is subjected to flash chromatography on silica gel, eluting with a mixture of ethyl acetate and pentane (65:35 v/v), followed by trituration with pentane, to give (E)-(RS)-4-(2-methylphenyl)-4-[2-{3-(2-pyridyl)prop-2-enoyl}-5-(3-thienylmethoxy)phenoxy]butanoic acid (0.22 g) in the form of a yellow solid, m.p. 65-70°C. [NMR(CDCl₃):- 2.1-2.9 (4H,m), 2.4(3H,s), 4.9(2H,q), 5.6(1H,q), 6.2-7.6(12H), 7.7(1H,d,J=16Hz), 7.8(1H,t), 8.0(1H,d,J=16Hz), 8.6(1H,d)].

EXAMPLE 47 Compounds GK and GL

A stirred solution of 2-hydroxy-4-(5-pyrimidinylmethoxy)acetophenone (0.65 g) in dry dimethylformamide (15 mL) is treated with sodium hydride (0.16 g; 60% w/v dispersion in mineral oil; 4 mmol). After 30 minutes the suspension is treated with ethyl (RS)-4-chloro-4-(2-methyl-phenyl)butanoate (0.96 g) and heated to 90°C for 6 hours. The reaction mixture is evaporated

and the residue is dissolved in ethyl acetate, washed with water, dried and evaporated. The residual oil is subjected to flash chromatography on silica gel, eluting with ethyl acetate, to give ethyl (RS)-4-[2-acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1.19 g) in the form of a brown oil. [NMR(CDCl₃):- 1.2(3H,m), 2.2-2.7(4H,m), 2.5(3H,s), 2.7(3H,s), 4.1(2H,t), 4.9(2H,q), 5.5(1H,q), 6.2-7.8(7H), 8.7 (2H,s), 9.2(1H,s).

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there is prepared ethyl (RS)-4-(2-methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]-butanoate in the form of a colourless oil

EXAMPLE 48 Compound GM

A stirred solution of ethyl (RS)-4-(2-methylphenyl)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-butanoate (0.90 g) in toluene (25 mL) is treated with ethyl pyridine-2-carboxylate (0.30 g) and sodium hydride (0.12 g; 60% w/v dispersion in mineral oil; 3 mmol) and heated at reflux for 4 hours. The solution is then evaporated, and the residual brown gum is dissolved in ethanol (40 mL) treated with acetic acid (0.48 g) and hydrazine hydrate (0.10 g) and heated at reflux for 2 hours. The solution is evaporated and the residual gum is subjected to flash chromatography on silica gel, eluting with a mixture of pentane and ethyl acetate (1:2 v/v), to give ethyl (RS)-4-(2-methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)-phenoxy]butanoate (0.20 g) in the form of a light brown oil. [NMR(CDCl₃):- 1.2(3H,t), 2.2-2.7 (4H,m), 2.5(3H,s), 4.1(2H,q), 4.9(2H,q), 5.6(1H,m), 6.3-7.7(13H,m), 7.8(1H,t), 8.0(1H,d), 8.7(1H,d)].

EXAMPLE 49 Compound GN

A stirred solution of ethyl (RS)-4-[2-carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2.44 g) in tetrahydrofuran (49mL) is treated with 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide (1.08 g) and maintained at 25°C for 24 hours. The solution is then evaporated and the residual oil is subjected to flash chromatography on silica gel, eluting with a mixture of pentane and ethyl acetate (2:1 v/v), followed by trituration with diethyl ether, to give ethyl (RS)-4-(2-methylphenyl)-4-[5-(3-thienylmethoxy)-2-thiocarbamoylphenoxy]butanoate (1.79 g) in the form of a

yellow solid, m.p. 134-135°C. [NMR(CDCl₃):- 1.2(3H,t), 2.2-2.6(4H,m), 2.4(3H,s), 4.1(2H,q), 4.9(2H,q), 5.6(1H,m), 6.2-7.8(9H,m), 8.7(1H,d)].

EXAMPLE 50 Compounds GO-GZ and HA-HL

5

By proceeding in a manner similar to that described hereinbefore in Example 38, and using the appropriate quantities of the corresponding esters as starting materials, there are prepared;

- 10 (RS)-4-[5-benzyloxy-2-(methylthio)phenoxy]-4-phenylbutanoic acid hydrate, in the form of a white solid, m.p. 104-105°C after recrystallisation from a mixture of petroleum ether (b.p.40-60°C) and ethyl acetate [Elemental analysis:- C,69.5; H,5.82;S,8.0%; Calculated for C₂₄H₂₄O₄S·0.25H₂O:- C,69.8;H,5.98;S,7.76%];
- 15 (E)-(RS)-4-[5-benzyloxy-2-(2-carboxyethenyl)phenoxy]-4-(2-methylphenyl)butanoic acid hydrate, in the form of a white solid, m.p. 151-153°C, after recrystallisation from a mixture of *t*-butyl methyl ether and pentane [Elemental analysis:- C,71.2;H,5.86%; Calculated for C₂₈H₂₈O₆·0.5H₂O:- C,71.2;H,5.97%]; (RS)-4-[2-(1-methyl-2-carboxyethenyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (E:Z ratio 3:1), in the
- 20 form of a white solid, m.p. 167-170°C, after recrystallisation from a mixture of cyclohexane and ethyl acetate [Elemental analysis:- C,66.8; H,5.65%; Calculated:- C,66.9;H,5.62%]; (RS)-4-[5-benzyloxy-2-hydroxyiminomethylphenoxy]-4-(2-methylphenyl)butanoic acid, (E:Z ratio 4:1), in the form of a yellow oil [Elemental analysis:- C,71.5;H,6.59;N,2.56%;
- 25 Calculated:- C,71.6; H,6.01;N,3.34%]; (RS)-4-[2-(N-(carboxymethoxy)iminomethyl)-5-(3-thienyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of an off-white solid, m.p. 114-116°C after recrystallisation from a mixture of isopropanol and pentane [Elemental analysis:- C,62.4; H,5.20;N,2.96%; Calculated:- C,62.1; H,5.21; N,2.90%];
- 30 (RS,RS)-4-[2-(1-hydroxyethyl)-5-(3-thienyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a colourless oil [Elemental analysis:- C,67.7;H,6.63%; Calculated:- C,67.6;H,6.14%];
- (RS)-4-(2-methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)phenoxy]-butanoic acid in the form of a white solid, m.p. 98-100°C, after recrystallisation
- 35 from heptane [Elemental analysis:- C,71.1;H,6.27%; Calculated:- C,71.1; H,6.20%]; (RS)-4-[2-(5-carboxy-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid characterised as the hydrochloride salt in the

- form of a white solid, m.p. 199-200°C, [Elemental analysis:- C,60.8;H,4.79; N,7.79%; Calculated for $C_{27}H_{25}N_3O_6 \cdot HCl \cdot 0.5H_2O$:- C,60.8; H,5.07;N,7.88%];
- (RS)-4-[2-(5-carboxy-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, in the form of a white solid, m.p. 209-210°C [Elemental analysis:- C,62.4;H,4.97;N,5.39%; Calculated for $C_{26}H_{24}O_6N_2S \cdot 0.5H_2O$:- C,62.3;H,5.02; N,5.59%; (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)butanoic acid in the form of a white solid; m.p. 158-160°C, after recrystallisation from a mixture of cyclohexane and diethyl ether [Elemental analysis:- C,56.3;H,4.42; Br,16.4%; Calculated:- C,56.5;H, 4.32;Br,16.3%; (RS)-
- 4-(2-methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]butanoic acid in the form of a pale yellow solid, m.p. 175-177°C, after recrystallisation from a mixture of ethyl acetate and cyclohexane [Elemental analysis:- C,61.8; H,4.87;N,3.24%; Calculated:- C,61.8;H,4.95; N,3.28%]; (RR,RS,SR,SS)-4-(2-methylphenyl)-4-[2-methyl-sulphinyl-5-(3-thienylmethoxy)phenoxy]butanoic
- acid, in the form of a yellow foam, by stirring the reaction mixture at ambient temperature for sixteen hours [Elemental analysis:- C,62.8; H,5.63;S,11.8%; Calculated:- C,62.2;H,5.41; S,14.41%]; (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoic acid in the form of a beige solid, m.p. 129-130°C, after recrystallisation from a mixture of cyclohexane and ethyl acetate [Elemental analysis:- C,59.8; H,4.41;S,6.71; Cl,7.79%; Calculated:- C,59.7;H, 4.36;S,6.93;Cl,7.66%]; (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoic acid in the form of a yellow solid, m.p. 112-114°C, after recrystallisation from a mixture of diisopropyl ether and pentane [Elemental analysis:- C,74.1; H,6.04%; Calculated:- C,74.2;H,5.98%]; (RS)-4-[2-formyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, in the form of a yellow solid, m.p. 124-126°C [Elemental analysis:- C,66.9; H,5.43;S,7.82%; Calculated:- C,67.1;H,5.63;S,7.81%]; (RS)-4-(2-methylphenyl)-4-[5-(3-thienylmethoxy)-2-(trifluoroacetyl)phenoxy]butanoic acid in the form of a white solid, m.p. 103-104°C, after flash chromatography on silica gel, eluting with a mixture of ethyl acetate and cyclohexane (2:3 v/v), followed by recrystallisation from cyclohexane [Elemental analysis:- C,60.0; H,4.32%; Calculated:- C,60.3;H 4.42%]; (RS)-4-(2-methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)phenoxy]butanoic acid in the form of a white solid, m.p. 109-110°C [using aqueous potassium carbonate solution (10% w/v) instead of the potassium hydroxide solution, and recrystallising from heptane] [Elemental analysis:- C,57.7;H,4.26%; Calculated:- C,57.6; H,4.23%]; (RS)-4-[2-cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the

- form of a white solid, m.p. 125-126°C, after recrystallisation from a mixture of ethyl acetate and cyclohexane [Elemental analysis:- C,68.0; H,5.55;N,2.94%; Calculated:- C,68.4;H,5.50; N,3.32%]; (RS)-4-[2-carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid,
- 5 m.p. 203-205°C, after recrystallisation from a mixture of ethyl acetate and acetonitrile [Elemental analysis:- C,64.7;H,5.48;N,3.09%; Calculated:- C,64.9; H,5.45;N,3.29%]; (RS)-4-[2-{N-(3-imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid, m.p. 146-148°C, after recrystallisation from acetonitrile [Elemental
- 10 analysis:- C,65.1; H,5.85;N,7.55%; Calculated:- C,65.2;H,5.86; N,7.87%]; (RS)-4-[2-{N-(2-carboxyethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid, m.p. 147-149°C, after recrystallisation from a mixture of ethyl acetate and cyclohexane [Elemental
- 15 analysis:- C,62.6;H,5.46;N,2.80%; Calculated:- C,62.8;H,5.47;N,2.82%]; (RS)-4-[2-{N-(carboxymethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid, m.p. 138-140°C, after recrystallisation from a mixture of ethyl acetate and cyclohexane [Elemental
- 20 analysis:- C,61.9;H,5.31;N,2.74%; Calculated:- C,62.1;H,5.21;N,2.90%]; (RS)-4-[2-{N-(2-cyanoethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid in the form of a white solid, m.p. 139-141°C, after recrystallisation from a mixture of ethyl acetate and cyclohexane [Elemental
- 25 [Elemental analysis:- C,63.7;H,5.41; N,5.20%; Calculated:- C,64.0;H,5.17; N,5.53%].

EXAMPLE 51 Compounds HM and HN

- 30 A mixture of ethyl (RS)-4-[2-{N-(cyano-methyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1.80 g), aqueous potassium carbonate solution (20 mL; 10% w/v) and methanol (200 mL) is stirred at ambient temperature for 16 hours. The reaction mixture is concentrated in vacuo and the residue is dissolved in water (50 mL) and
- 35 acidified to pH1 by treatment with hydrochloric acid (1 N), then extracted with ethyl acetate (150 mL). The organic extract is washed with water (50 mL), dried over magnesium sulphate, and evaporated to dryness. Flash

chromatography of the residue on silica gel, eluting with a mixture of methanol and dichloromethane (1:19 v/v) gives two products:- (RS)-4-[2-{N-(carbamoylmethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, in the form of a white solid, m.p. 202-204°C, after recrystallisation from a mixture of ethyl acetate and acetonitrile [Elemental analysis:- C,62.3;H,5.44;N,5.66%; Calculated:- C,62.2;H,5.43;N,5.81%]; and (RS)-4-[2-{N-(methoxycarbonylmethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, in the form of a white solid, m.p. 156-158°C, after recrystallisation from a mixture of ethyl acetate and acetonitrile [Elemental analysis:- C,62.8;H,5.48;N,2.90%; Calculated:- C,62.8;H,5.47;N,2.82%].

EXAMPLE 52 Compounds HO-HZ and IA-IB

By proceeding in a manner similar to that described hereinbefore in Example 40, and using the appropriate quantities of the corresponding starting materials, there are prepared:- ethyl (RS)-4-[5-benzyloxy-2-(methylthio)phenoxy]-4-phenylbutanoate, in the form of a yellow oil; ethyl (RS)-4-(2-methylphenyl)-4-[2-methylthio-5-(3-thienylmethoxy)phenoxy]butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-chloro-6-fluorophenyl)butanoate, in the form of an orange oil; ethyl (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-formyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[5-(3-thienylmethoxy)-2-trifluoroacetylphenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-carbamoyl-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a colourless oil; ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)butanoate, in the form of a colourless oil; ethyl (RS)-4-(2-methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-{N-methoxycarbonylmethyl}carbamoyl]-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-{N-(3-imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-{N-(2-cyanoethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-{N-cyanomethyl}carbamoyl]-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl (RS)-4-[2-{N-(2-methoxycarbonylethyl)carbamoyl}-5-(3-thienyl-

methoxy)phenoxy]-4-(2-methylphenyl)butanoate, in the form of a yellow oil; ethyl 4-[2-(1-methoxycarbonyl-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the diastereomers (1R, 4RS), in the form of a yellow oil; ethyl 4-[2-(1-methoxycarbonyl-2-phenyl-ethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the diastereomers (1S, 4RS), in the form of a yellow oil; methyl 2-[2-hydroxy-4-(3-thienylmethoxy)benzoylamino]-3-phenylpropionate, thought to be the (R)-enantiomer, in the form of a yellow oil; and methyl 2-[2-hydroxy-4-(3-thienyl-methoxy)benzoylamino]-3-phenylpropionate, thought to be the (S)-enantiomer, in the form of a yellow oil.

EXAMPLE 53 Compound IC

A solution of ethyl (RS)-4-(2-methylphenyl)-4-[2-methylthio-5-(3-thienylmethoxy)phenoxy]butanoate in acetone (20 mL), at 0°C. is treated dropwise with a solution of potassium peroxymonosulphate (1.01 g of commercial 2KHSO₅:KHSO₄:K₂SO₄). The reaction mixture is stirred at 0°C for 20 minutes, then treated with saturated aqueous sodium bisulphate solution (20 mL), and extracted with dichloromethane (2 x 50 mL). The combined organic extracts are dried over magnesium sulphate, filtered and concentrated in vacuo, to give ethyl (RS,RS)-4-(2-methylphenyl)-4-[2-methyl-sulphinyl-5-(3-thienylmethoxy)phenoxy]-butanoate (0.75 g) in the form of a yellow oil.

EXAMPLE 54 Compounds ID and IE

A stirred suspension of sodium hydride (0.272 g; 60% w/v dispersion in mineral oil; 6.8 mmol) in tetrahydrofuran (100 mL) at 0°C under nitrogen is treated, dropwise, with methyl (diethylphosphono)acetate (1.43 g) and stirred for 15 minutes. It is then treated with a solution of ethyl (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoate (1.96 g) in tetrahydrofuran (5 mL) in one portion, and the reaction mixture is stirred at ambient temperature for 15 minutes. The mixture is treated with water (100 mL) and extracted with ethyl acetate (3 x 100 mL), and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate, and filtered and the solvent is removed in vacuo, to give ethyl (E)-(RS)-4-(2-methylphenyl)-4-[2-(2-methoxycarbonylethenyl)-5-benzyloxyphenoxy]butanoate, in the form of a yellow oil.

By proceeding in a similar manner but replacing the ethyl (RS)-4-(5-benzyloxy-2-formyl-phenoxy)-4-(2-methylphenyl)butanoate used as starting material by the appropriate quantity of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate, there is prepared
5 ethyl (RS)-4-[2-(2-methoxycarbonyl-1-methylethenyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate (E:Z ratio 3:1), in the form of a yellow oil.

EXAMPLE 55 Compounds IF and IG

10

A mixture of ethyl (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoate (0.49 g), hydroxylamine hydrochloride (0.1 g) and pyridine (0.1 g) in ethanol (10 mL) is heated at reflux for 1 hour. The solvent is then removed in vacuo and the residue is partitioned between water (50 mL) and
15 ethyl acetate (50 mL). The aqueous layer is extracted with ethyl acetate (2 x 50 mL) and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate and filtered and the solvent is removed in vacuo, to give a yellow oil. Flash chromatography on silica gel, eluting with a mixture of petroleum ether (b.p.40-60°C) and ethyl acetate (7:1 v/v), gives ethyl (RS)-4-
20 (5-benzyloxy-2-hydroxyiminomethylphenoxy)-4-(2-methylphenyl)butanoate (E:Z ratio 4:1), (0.24 g).

By proceeding in a similar manner, but replacing the ethyl (RS)-4-(5-benzyloxy-2-formyl-phenoxy)-4-(2-methylphenyl)butanoate used as
25 starting material by the appropriate quantity of ethyl (RS)-4-[2-formyl-5-(3-thienylmethoxy))phenoxy]-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-hydroxyiminomethyl-5-(3-thienylmethoxy))phenoxy]-4-(2-methylphenyl)butanoate (E:Z ratio 4:1).

30 EXAMPLE 56 Compound IH

A stirred solution of ethyl (RS)-4-(5-benzyloxy-2-hydroxyiminomethylphenoxy)-4-(2-methylphenyl)butanoate (3.68 g) in tetrahydrofuran (100 mL) at ambient temperature is treated portionwise with sodium hydride (0.33 g; 60%
35 w/v dispersion in mineral oil; 8.25 mmol) during 30 minutes. It is then treated with ethyl bromoacetate (1.58 g) and stirred for one hour at ambient temperature then for 5 hours at 80°C. It is then treated with a further portion of

ethyl bromoacetate (0.40 g) and heating at 80°C is continued for one hour. The solvent is then removed in vacuo and the residue is partitioned between saturated aqueous potassium carbonate solution (50 mL) and ethyl acetate (50 mL). The aqueous layer is extracted with ethyl acetate (2 x 50 mL) and the
5 combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate and filtered and the solvent is removed in vacuo, to give a yellow oil. Flash chromatography on silica gel, eluting with a mixture of petroleum ether (b.p.40-60°C) and ethyl acetate (2:1 v/v) gives ethyl
(RS)-4-[2-{N-(ethoxycarbonylmethoxy)iminomethyl}-5-(3-thienylmethoxy)-
10 phenoxy]-4-(2-methylphenyl)butanoate (E:Z ratio 4:1) (2.66 g), in the form of a yellow oil.

EXAMPLE 57 Compound II

15 A stirred solution of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1 g) in methanol (50 mL) is treated, portionwise, with sodium borohydride (0.1 g), and stirred at ambient temperature for one hour. It is then treated with dilute acetic acid (50 mL; 1 N) and extracted with ethyl acetate (2 x 50 mL). The combined organic extracts are washed with
20 water (2 x 50 mL) dried over magnesium sulphate, filtered and concentrated in vacuo. Flash chromatography on silica gel, eluting with a mixture of ethyl acetate and cyclohexane (1:5 v/v) gives ethyl (RS,RS)-4-[2-(1-hydroxyethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.

25 EXAMPLE 58 Compound IJ

A stirred suspension of methyltriphenylphosphonium bromide (2.4 g) in tetrahydrofuran (50 mL) at ambient temperature is treated with a solution of butyllithium in hexanes (2.6 mL; 2.5 M). After two hours, the reaction mixture is
30 treated with a solution of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (1 g) in tetrahydrofuran (2 mL) and stirred for a further two hours. It is then partitioned between hydrochloric acid (50 mL; 1 N) and ethyl acetate (100 mL). The organic layer is washed with water (50 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The
35 residue is subjected to flash chromatography on silica gel, eluting with a mixture of ethyl acetate and cyclohexane (1:9 v/v) to give ethyl

(RS)-4-(2-methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)phenoxy]-butanoate (0.7 g), in the form of a colourless oil.

EXAMPLE 59 Compounds IK and IL

- 5 A stirred suspension of sodium hydride (0.64 g; 60% w/v dispersion in mineral oil; 16 mmol) in toluene (100 mL) is treated, dropwise, with a solution of ethyl (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (6.5 g) and diethyl oxalate (6.57 g) in toluene (20 mL), and the
- 10 mixture is heated at reflux for 5 hours. It is then evaporated to dryness, and the resulting residue is dissolved in ethanol (100 mL) and treated with hydrazine hydrate (0.75 g) and glacial acetic acid (3.6 g) and heated at reflux for 3 hours. The mixture is then partitioned between ethyl acetate (100 mL) and aqueous potassium carbonate solution (100 mL; 10% w/v). The organic layer is washed
- 15 with water (100 mL), dried over magnesium sulphate and evaporated to dryness. The resulting residue is subjected to flash chromatography on silica gel, eluting with ethyl acetate, to give ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-ethylphenyl)butanoate (2.6 g).
- 20 By proceeding in a similar manner but replacing the ethyl (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate used as starting material by the appropriate quantity of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-
- 25 4-(2-methylphenyl)butanoate.

EXAMPLE 60 Compound IM

- 30 A solution of ethyl (RS)-4-(2-methylphenyl)-4-[2-trifluoroacetyl-5-(3-thienylmethoxy)-phenoxy]butanoate (1.3 g) and diethylaminosulphur trifluoride (0.64 mL) in dichloromethane is stirred at -5°C for 30 minutes, and then at ambient temperature for 16 hours. The reaction mixture is treated with a further portion of diethylaminosulphur trifluoride (0.32 mL) and stirred for 1 hour. It is then treated with saturated aqueous sodium bicarbonate solution (10 mL) and
- 35 stirred for 15 minutes, then it is diluted with dichloromethane (50 mL) and the organic phase is washed with water (50 mL), dried over magnesium sulphate, filtered and evaporated to dryness. The resulting residue is subjected to flash

chromatography on silica gel, eluting with a mixture of ethyl acetate and cyclohexane (1:19v/v), to give ethyl (RS)-4-(2-methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)phenoxy]butanoate (0.6 g), in the form of a yellow oil.

5 EXAMPLE 61 Compound IN

By proceeding in a manner similar to that described hereinbefore in Example 42, but replacing the ethyl (RS)-4-(5-benzyloxy-2-carbamoylphenoxy)-4-(2-methylphenyl)butanoate with the appropriate quantity of ethyl
10 (RS)-4-[2-carbamoyl-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, there is prepared ethyl (RS)-4-[2-cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.

EXAMPLE 62 Compound IO

15

A mixture of (RS)-4-[2-(5-carboxy-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (0.5 g), iodomethane (0.25 mL), potassium carbonate (0.55 g) and dimethylformamide (20 mL) is stirred at ambient temperature for 36 hours. The solvent is then removed in vacuo and
20 the resulting residue is dissolved in ethyl acetate (50 mL), and this solution is washed with water (3 x 50 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The resulting residue is subjected to flash
chromatography on silica gel, eluting with a mixture of petroleum ether (b.p. 40-60°C) and ethyl acetate (2:1 v/v), to give methyl (RS)-4-[2-(5-methoxycarbonyl-
25 1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (0.37 g), in the form of a colourless oil.

EXAMPLE 63 Compound IP

30 (a) A solution of (RS)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid (500 mg) in dichloromethane (50 mL) is treated with 1,1'-carbonyldiimidazole (215 mg) and stirred at 25°C for 2 hours, by the end of which period the evolution of carbon dioxide gas has ceased. The resulting solution is designated "solution A".

35

(b) A solution of benzenesulphonamide (1.16 g) in dimethylformamide (50 mL) is treated with sodium hydride (201 mg; 60% w/v dispersion

in mineral oil; 5 mmol) and stirred for 2 hours. The solution is then treated, dropwise, with "solution A" prepared as hereinbefore in part (a), and stirred for 18 hours at 25°C. The solution is then evaporated and the resulting residue is partitioned between ethyl acetate and water. The organic layer is dried and
5 evaporated, and the residual oil is subjected to flash chromatography on silica gel, eluting with a mixture of pentane and ethyl acetate (2:1 v/v), to give (RS)-N-[4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoyl]benzenesulphonamide (250 mg), in the form of a colourless solid, m.p. 68-70°C [NMR(CDCI₃):- 2.0-2.8(4H,m), 2.3(3H,s), 4.9(2H,q), 5.3(1H,q), 6.1-
10 8.0(10H)].

EXAMPLE 64 Compounds IQ and IR

(RS)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-
15 butanoic acid is separated into its enantiomers, (R)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid and (S)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, by chiral HPLC, under the following conditions:- Chiralcel OD column; mobile phase of isopropanol/acetic acid/heptane (10:1:100 v/v); flow rate 1 mL/minute;
20 temperature ambient; UV detection 270 nm.

The (R)-enantiomer is thought to be the more mobile of the two.

EXAMPLE 65 Compound IQ

25 A mixture of sodium 4-hydroxy-4-(2-methyl-phenyl)butanoate, thought to be the (R)-enantiomer, (3 g) and sodium hydride (1.5 g; 60% w/v dispersion in mineral oil; 37.5 mmol) in tetrahydrofuran (50 mL) is stirred at ambient temperature for 1 hour under an atmosphere of argon, and is then warmed to
30 55°C. It is then treated with 2-fluoro-4-(3-thienylmethoxy)benzonitrile (3 g), in one portion, and stirring is continued at 55°C for 15 hours. The mixture is then cooled, diluted with hydrochloric acid (200 mL; 1 N) and extracted with ethyl acetate (4 x 100 mL). The combined extracts are dried over magnesium sulphate and concentrated in vacuo, to give an oily residue. This residue is
35 boiled in isopropyl ether, and the resulting mixture is filtered while hot, and then allowed to cool. The yellow solid which separates is collected and recrystallized from a mixture of ethyl acetate and hexane, to give 4-[2-cyano-5-

(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the (R)-enantiomer, (1.5 g) in the form of a white fluffy solid, m.p. 145°C. Chiral HPLC assay showed a 98% enantiomeric excess. [Elemental analysis:- C,67.82;H,5.18; N,3.24%; Calculated:- C,67.80;H,5.19;N,3.44%].

5

EXAMPLE 66 Compound IS

A mixture of sodium 4-hydroxy-4-(2-methyl-phenyl)butanoate, thought to be the (R)-enantiomer, (3 g) and sodium hydride (1.5 g; 60% w/v dispersion in mineral oil; 37.5mmol) in tetrahydrofuran (50 mL) is stirred at ambient temperature for 1 hour under an atmosphere of argon and is then warmed to 55°C. It is then treated with 2-fluoro-4-(3-pyridyl-methoxy)benzonitrile (3 g), in one portion, and stirring is continued at 55°C for 15 hours. The reaction mixture is then cooled, concentrated in vacuo, and diluted with water (100 mL). The pH of the mixture is adjusted to 5 by treatment with concentrated hydrochloric acid and is then extracted with ethyl acetate (4 x 100 mL). The combined extracts are dried over magnesium sulphate and concentrated in vacuo, to give an oily residue. This residue is recrystallized from isopropyl ether to give 4-[2-cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid, thought to be the (R)-enantiomer, (1.5 g) in the form of a white solid, m.p. 130-134°C. Chiral HPLC assay showed a 98% enantiomeric excess. [Elemental analysis:- C,71.12; H,5.49;N,6.62%; Calculated:- C,71.64;H,5.47;N, 6.97%].

25 EXAMPLE 67 Compound IT

A mixture of 4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the (R)-enantiomer, (50 mg), dicyclohexylamine (0.5 mL) and diethyl ether (10 mL) is stirred at ambient temperature for 18 hours. The resulting precipitate is collected and washed with diethyl ether, to give dicyclohexylammonium 4-[2-cyano-5-(3-pyridyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the (R)-enantiomer, (30 mg) in the form of a white solid, m.p. 92-93°C.

35 EXAMPLE 68 Compound IU

A solution of triphenylphosphine (0.57 g) in tetrahydrofuran (10 mL) cooled at 0°C is treated with diisopropyl azodicarboxylate (0.42 mL) and stirred for 20 minutes at 0°C. It is then treated, dropwise, with a solution of *t*-butyl 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (R)-enantiomer, (0.35 g) and 2-hydroxy-4-(3-thienylmethoxy)benzonitrile (0.25 g) in tetrahydrofuran (10 mL). Stirring is continued at 0°C for 30 minutes and then at ambient temperature for 18 hours. The solvent is evaporated in vacuo and the oil is subjected to flash chromatography on silica gel, eluting with a mixture of cyclohexane and dichloromethane (1:3 v/v), to give a yellow oil (0.46 g). This oil is dissolved in methanol (5 mL) and tetrahydrofuran (5 mL). The resulting solution is treated with aqueous sodium hydroxide solution (3 mL; 5 N) and stirred at ambient temperature for 24 hours, and then it is concentrated to dryness in vacuo. The resulting residue is diluted with hydrochloric acid (50 mL; 1 N) and extracted with ethyl acetate (2 x 50 mL) and the combined extracts are dried over magnesium sulphate and concentrated to dryness in vacuo, to give a solid, which is recrystallized twice from a mixture of ethyl acetate and cyclohexane, to give 4-[2-cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the (S)-enantiomer, (0.22 g) in the form of a white fluffy solid, m.p. 143°C. Chiral HPLC assay showed a 99% enantiomeric excess. [Elemental analysis:- C,67.67;H,5.14;N,3.28%; Calculated:- C,67.80;H,5.19;N,3.44%].

EXAMPLE 69 Compound IV

A stirred solution of ethyl (RS)-5-(3-benzylthiophenyl)-4-(2-methylphenyl)-5-oxopentanoate (0.56 g) and 1 N sodium hydroxide (3.4 mL) in dioxan (11 mL) is heated at 70°C for 1.5 hours. The reaction mixture is evaporated and the residue dissolved in water (6 mL). The pH of the solution is adjusted to pH 1 by addition of 2 N hydrochloric acid and the mixture extracted with ether (20 mL). The organic phase is washed twice with water (6 mL), dried over magnesium sulphate and evaporated. The residual orange oil (0.59 g) is triturated with pentane and crystallised from cyclohexane to give (RS)-5-(3-benzylthiophenyl)-4-(2-methylphenyl)-5-oxopentanoic acid hemihydrate as a pale yellow solid (0.37 g), m.p. 105-107°C. [Elemental analysis:- C,72.4; H,5.94%. Calculated for C₂₅H₂₄O₃S·0.5H₂O:- C,72.4; H,5.91%].

EXAMPLE 70 Compound IW

A mixture of (RS)-4-(2-methylphenyl)-5-oxo-(1-pyrid-3-ylmethoxy)-
pentanoic acid (0.3 g) and hydrogen peroxide (0.1 mL, 27.5% w/w) in acetic
5 acid (3 mL) is heated at 60°C for 12 hours. The reaction mixture is evaporated
and the residue partitioned between ethyl acetate and water. The organic
phase is washed with water, dried over magnesium sulphate and evaporated
in vacuo. The residue is purified by flash chromatography on silica eluting with
a mixture of dichloromethane and methanol, (95:5 v/v). Fractions homo-
10 geneous in the required product are combined and evaporated. The residue is
crystallised from ether to give (RS)-4-(2-methylphenyl)-5-oxo-(1-oxopyrid-3-
ylmethoxy)pentanoic acid hemihydrate as an off-white solid. [Elemental
analysis:- C,69.5; H,5.69; N,3.09%. Calculated for $C_{24}H_{23}NO_5 \cdot 0.5H_2O$:-
C,69.5; H,6.08; N,3.33%].

15

EXAMPLE 71 Compound IX

A mixture of (RS)-4-[2-formyl-5-(3-thienylmethoxy)phenoxy]-4-(2-
20 methylphenyl)butanoic acid (0.7 g), O-benzylhydroxylamine hydrochloride
(0.48 g) and pyridine (1 mL) in ethanol (50 mL) is heated at 65°C for 5 hours.
The reaction mixture is evaporated. The residue is dissolved dichloromethane
(80 mL), washed with 1 N hydrochloric acid, then with water, dried over
magnesium sulphate and evaporated. The resulting oil is purified by flash
25 chromatography on silica eluting with a mixture of dichloromethane and
methanol (95:5 v/v). Fractions homogeneous in the required product are
combined, evaporated and the solid residue triturated with pentane to give
(RS)-4-[2-(benzyloxyiminomethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-
methylphenyl)butanoic acid hydrate (0.2 g) as a cream coloured solid, m.p.
30 35-37°C. [Elemental analysis:- C,69.2; H,5.60; N,2.58; S,6.26%. Calculated
for $C_{25}H_{23}NO_8 \cdot 0.3H_2O$:- C,69.2; H,5.72; N,2.69; S,6.16%]

EXAMPLE 72 Compound IY

35 A stirred solution of ethyl (RS)-4-(2-carboxycarbonyl-5-(3-thienyl-
methoxy)phenoxy)-4-(2-methylphenyl)butanoate (0.37 g) and 1 N sodium
hydroxide (2.7 mL) in dioxan (6.6 mL) is heated at 60°C for 1.5 hours. The

reaction mixture is evaporated and the residue dissolved in water (5 mL). The pH of the solution is adjusted to 1 by addition of 1 N hydrochloric acid and the mixture extracted with ethyl acetate (20 mL). The organic phase is dried over magnesium sulphate and evaporated. The residual yellow oil is triturated with ether to give (RS)-4-(2-Carboxycarbonyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoic acid as a yellow solid (0.29 g), m.p. 173-175°C. [Elemental analysis:- C,63.2; H,5.07%. Calculated:- C,63.4; H,4.88%].

EXAMPLE 73 Compound IZ

A solution of tert-butyl 4-[2-benzoyl-5-(pyridin-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the (R)-enantiomer, (1 g) in a mixture of methanol (10 mL) and tetrahydrofuran (10 mL) is treated with 5 M aqueous sodium hydroxide solution (7.5 mL) and the reaction stirred at ambient temperature for 24 hours. The reaction is concentrated to low volume, acidified to pH 5 with dilute hydrochloric acid, and extracted into dichloromethane (2 x 30 mL). The combined organic layers are washed with brine, dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a yellow gum. Trituration with diethyl ether and recrystallization of the resultant solid from acetonitrile containing a little decolourizing charcoal gives 4-[2-benzoyl-5-(pyridin-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the (R)-enantiomer, (0.2 g) as a pale yellow solid, m.p. 185-186°C. [Elemental analysis:- C, 74.83; H, 5.65; N, 2.91%; Calculated:- C, 75.01; H, 5.72; N,2.87%].

EXAMPLE 74 Compound JA

A stirred solution of methyl 3-[3-(3-carboxy-1-(2-methylphenyl)propoxy)-4-nitrophenoxy]benzoate (0.86 g) in a mixture of methanol (50 mL) and 50% aqueous potassium hydroxide solution (10 mL) is heated at reflux for 2 hours. The reaction mixture is evaporated in vacuo and the residue partitioned between ethyl acetate (50 mL) and 2 N hydrochloric acid (50 mL). The organic phase is washed three times with water (50 mL), dried over magnesium sulphate and evaporated to give a yellow solid (0.8 g) which is purified by flash chromatography on silica eluting initially with a mixture of cyclohexane and ethyl acetate (7:3 v/v) to remove the starting material then eluting with a mixture of cyclohexane and ethyl acetate (1:1 v/v). Fractions

homogeneous in the required product are combined and evaporated to give a clear yellow oil (0.19 g) which is triturated with a mixture of ether and pentane to give (RS)-3-[3-(3-carboxy-1-(2-methylphenyl)propoxy)-4-nitrophenoxy-methyl]benzoic acid as a pale yellow solid (0.11 g), m.p. 85-87°C. [Elemental analysis:- C,64.5; H,5.10; N,2.60%. Calculated for $C_{25}H_{23}NO_8 \cdot 0.3H_2O$:- C,64.5; H,4.98; N,3.01%].

EXAMPLE 75 Compound JB

10 A solution of dry sodium (RS)-4-hydroxy-4-(2-methylphenyl)butanoate (0.18 g) in dry tetrahydrofuran (50 mL) is treated with sodium hydride and stirred at 60°C for 15 minutes. tert-Butyl 3-[(4-cyano-3-fluoro-phenoxy-methyl)-phenyl]propionate (0.2 g) is added and the mixture heated at reflux for 4 hours. After standing at room temperature for 18 hours the reaction mixture is filtered.

15 Ethyl acetate (50 mL) and 1 N hydrochloric acid (50 mL) are added to the filtrate and the organic phase is separated. The organic phase is washed with water (10mL), dried over magnesium sulphate, evaporated and the residue purified by flash chromatography on silica eluting with a mixture of dichloromethane and methanol (95:5 v/v). Fractions homogeneous in the

20 required product are combined and evaporated. The residue is dissolved in ether and the solution washed with 1% sodium bicarbonate solution, 0.5 N hydrochloric acid, water, dried over magnesium sulphate and evaporated. The resulting colourless gum is crystallised from a mixture of cyclohexane and ethyl acetate to give (RS)-4-[5-[3-(2-carboxyethyl)benzyloxy]-2-cyano-phenoxy]-4-

25 (2-methylphenyl)butanoic acid (0.12 g) m.p. 136-137°C. [Elemental analysis:- C,71.4; H,6.23; N,2.91%. Calculated:- C,71.0; H,5.75; N,2.96%].

EXAMPLE 76 Compound JC

30 A solution of ethyl (RS)-4-[2-(4,5-dihydrooxazol-2-yl)-5-(3-thienyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoate (500 mg) in methanol (50 mL) containing 10% w/v potassium carbonate (5 mL) is stirred at ambient temperature for 16 hours. The reaction mixture is concentrated under reduced pressure and partitioned between water (100 mL) and ethyl acetate (100 mL)

35 then acidified to pH 5 with 1 N HCl. The organic phase is washed three times with water (50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure. Recrystallisation from cyclohexane/ethyl acetate

gives (RS)-4-[2-(4,5-dihydrooxazol-2-yl))-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid hydrate as a white solid (160 mg), m.p. 107-109°C. [Elemental analysis:- C,65.9, H, 5.69, N, 2.94%. Calculated for $C_{25}H_{25}NO_5S \cdot 0.2H_2O$:- C,66.0, H, 5.59, N, 3.01%].

5

EXAMPLE 77 Compounds JD and JE

A solution of ethyl 4-[2-(1-methoxycarbonyl-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the diastereomers (1R, 4RS), (0.48 g) in methanol (100 mL) is treated with 10% aqueous potassium carbonate solution (10 mL). The reaction mixture is stirred at room temperature for 18 hours and evaporated. The residue is dissolved in water (50 mL), the solution acidified to pH 1 by addition of 1 N hydrochloric acid and extracted with ethyl acetate (120 mL). The organic phase is washed with water (100 mL), brine (100 mL), dried over magnesium sulphate and evaporated. The semi-solid residue (0.44 g) is boiled with acetonitrile (20 mL), the mixture cooled to room temperature and filtered to remove 2-[2-hydroxy-4-(3-thienylmethoxy)benzoylamino]-3-phenylpropionic acid (0.047 g) as a white solid, m.p. 178-180°C. [Elemental analysis:- C,63.52; H,4.76; N,3.15%. Calculated :- C,63.46; H,4.82; N,3.52%]. The filtrate is evaporated to give 4-[2-(1-carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the diastereomers (1R, 4RS), (0.3 g), as an oily semi-solid. [Elemental analysis:- C,66.54; H,5.45; N,2.35%. Calculated :- C,67.00; H,5.45; N,2.44%].

25

By proceeding in a similar manner but replacing the ethyl 4-[2-(1-methoxycarbonyl-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the diastereomers (1R, 4RS), by the appropriate quantity of ethyl 4-[2-(1-methoxycarbonyl-2-phenyl-ethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoate, thought to be the diastereomers (1S, 4RS), there is prepared ethyl 4-[2-(1-carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, thought to be the diastereomers (1S, 4RS), m.p. 149-150°C. [Elemental analysis:- C,74.8; H,5.24; N,2.84%. Calculated :- C,75.1; H,5.26; N,2.92%].

35

EXAMPLE 78 Compound JF

- A solution of ethyl (RS)-4-[2-(oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (140 mg) in methanol (25 mL) containing 10% w/v potassium carbonate (2 mL) is stirred at reflux for 2 hours.
- 5 The reaction mixture is partitioned between 1 N HCl (50 mL) and ethyl acetate (50 mL) then the organic phase is washed with water (2 x 50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure. Recrystallisation from cyclohexane/ethyl acetate gives (RS)-4-[2-(oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid as a white
- 10 solid (70 mg), m.p. 112-114°C. [Elemental analysis:- C,66.8, H, 5.16, N, 3.12%. Calculated:- C,66.5, H, 5.04, N, 3.01%]

EXAMPLE 79 COMPOUND JG

- 15 A solution of methyl (RS)-2-[2-(3-ethoxycarbonyl-1-(2-methylphenyl)propoxy)-4-(3-thienylmethoxy)benzoylamino]acrylate (200 mg) in methanol (20 mL) containing 10% w/v potassium carbonate (2 mL) is stirred at ambient temperature for 16 hours and at reflux for 3 hours. The reaction mixture is partitioned between 1N HCl (50 mL) and ethyl acetate (50 mL) then the organic
- 20 phase is washed with water (2 x 50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure. Recrystallisation from cyclohexane/ethyl acetate gives (RS)-2-[2-(3-carboxy-1-(2-methylphenyl)propoxy)-4-(3-thienylmethoxy)benzoylamino]acrylic acid as a white, fluffy solid (20 mg), m.p. 140-143°C. [Elemental analysis:- C,61.9, H, 5.32, N, 2.53%. Calculated:- C,61.9, H, 5.16, N, 2.78%]
- 25

EXAMPLE 80 Compound JH

- 30 A solution of ethyl (RS)-4-[2-(4-methoxycarbonyloxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl) butanoate (300 mg) in methanol (25 mL) containing 10% w/v potassium carbonate (2 mL) is stirred at reflux for 3 hours. The reaction mixture is partitioned between 1N HCl (50 mL) and ethyl acetate (50 mL) then the organic phase is washed with water (50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure.
- 35 Recrystallisation from cyclohexane/ethyl acetate gives (RS)-4-[2-(4-carboxyoxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid

(100 mg), in the form of a white solid, m.p. 178-179°C. [Elemental analysis:- C, 62.4, H, 4.57, N, 2.79%. Calculated for $C_{26}H_{23}NO_7S \cdot 0.2H_2O$:- C, 62.2, H, 4.78, N, 2.79%].

5 EXAMPLE 81 Compound JI

A solution of ethyl (RS)-4-(2-cyano-5-(3-thienylmethoxy)phenoxy)-4-(2-chloro-6-fluorophenyl)butanoate (360 mg) in methanol (10 mL) containing 1 N sodium hydroxide (3 mL) is stirred at ambient temperature for 16 hours. The
10 reaction mixture is poured into 0.5 N HCl (20 mL) and extracted twice with ethyl acetate (20 mL). The combined organic phases are washed with brine (20 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a dark yellow oil. Crystallisation from a mixture of cyclohexane and ethyl acetate gives (RS)-4-(2-chloro-6-fluorophenyl)-4-(2-
15 cyano-5-(3-thienylmethoxy)phenoxy)-butanoic acid (210 mg) in the form of a beige solid, m.p. 128-130°C.

EXAMPLE 82 Compound JJ

A solution of tert-butyl (R)-4-(2-methylphenyl)-4-(2-nitro-5-(3-pyridylmethoxy)phenoxy)butanoate (800 mg) in methanol (20 mL) containing 5 N sodium hydroxide (10 mL) is stirred at ambient temperature for 48 hours. The reaction mixture is poured into water (50 mL), acidified to pH 5 with 1 N HCl and extracted three times with ethyl acetate (50 mL). The combined organic
25 phases are washed with brine (50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure. The residual brown gum is purified by flash chromatography on silica eluting with ethyl acetate. Fractions homogeneous in the required product are combined and evaporated to give a yellow oil. Trituration with pentane afforded (R)-4-(2-methylphenyl)-4-(2-nitro-
30 5-(3-pyridylmethoxy)phenoxy)butanoic acid as a white solid (150 mg), m.p. 117-119°C. [Elemental analysis:- C, 65.4, H, 5.21, N, 6.64%. Calculated:- C, 65.3, H, 5.17, N, 6.59%]

EXAMPLE 83 Compound JK

35 A solution of ethyl (RS)-4-(2-cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoate (500 mg) in methanol (20 mL) containing 1 N

sodium hydroxide (5 mL) is stirred at ambient temperature for 16 hours. The reaction mixture is poured into 0.5 N HCl (20 mL) and extracted three times with ethyl acetate (20 mL). The combined organic phases are washed with brine (20 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a brown oil. Crystallisation with cyclohexane gives (RS)-4-(2-cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoic acid, (170 mg) in the form of an off-white solid, m.p. 135-136°C. [Elemental analysis:- C,68.4, H, 5.46, N, 3.33%. Calculated:- C,68.7, H, 5.57, N, 3.00%]

10

EXAMPLE 84 Compound JL

A solution of sodium (RS)-4-(benzo[1,3]dioxol-4-yl)-4-hydroxybutanoate (1.23 g) is added to a stirred suspension of sodium hydride (0.6 g, 60% dispersion in mineral oil) in tetrahydrofuran (50 mL). The mixture is stirred at room temperature for 1.5 hours when 2-fluoro-4-(3-pyridylmethoxy)benzonitrile (1.07 g) is added in one portion. The reaction mixture is heated at 55-60°C for 18 hours and evaporated. The residue is dissolved in water (100 mL), the solution acidified to pH 2 with 1 N hydrochloric acid and the mixture extracted with ethyl acetate (100 mL). The organic phase is washed with water (100 mL), with brine (100 mL), dried over magnesium sulphate and evaporated. The yellow semi-solid residue is recrystallised from acetonitrile to give (RS)-4-(benzo[1,3]dioxol-4-yl)-4-[2-cyano-5-(3-pyridylmethoxy)phenoxy]butanoic acid (0.80 g) as a cream solid, m.p. 181-183°C. [Elemental analysis:- C,66.32; H,4.57; N,6.52%. Calculated :- C,66.66; H,4.66; N,6.48%]

25

EXAMPLE 85 Compound JM

A solution of sodium (RS)-4-(2,3-dimethylphenyl)-4-hydroxybutanoate (1 g) is added to a stirred suspension of sodium hydride (0.5 g, 60% dispersion in mineral oil) in tetrahydrofuran (50 mL) under nitrogen. The mixture is stirred at room temperature for 3 hours when 2-fluoro-4-(3-pyridylmethoxy)benzonitrile (1 g) is added in one portion. The reaction mixture is heated at 55-60°C for 18 hours and evaporated. The residue is dissolved in water (100 mL), the solution acidified to pH 1 with 1 N hydrochloric acid. The precipitated solid is filtered, washed with water, ethyl acetate and ether, dried, and recrystallised from ethyl acetate to give (RS)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]-4-

35

(2,3-dimethylphenyl)butanoic acid (0.85 g) as a white solid, m.p. 187-189°C. [Elemental analysis:- C,71.76; H,6.59; N,5.67%. Calculated :- C,72.10; H,6.73; N,5.81%]

5 EXAMPLE 86 Compound JN

A solution of (RS)-4-[2-cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy)phenyl)butanoic acid (1.8 g) in tetrahydrofuran (150 mL) is treated with 1 N hydrochloric acid (10 mL) and the
10 mixture heated at reflux for 1 hour. The reaction mixture is evaporated and the residue partitioned between ethyl acetate (200 mL) and water (100 mL). The organic phase is washed with water (100 mL), brine (100 mL), dried over magnesium sulphate and evaporated. The residual brown gum (1.25 g) is recrystallised from acetonitrile to give (RS)-4-[2-cyano-5-(3-pyridylmethoxy)-
15 phenoxy]-4-(5-hydroxy-2-methylphenyl)butanoic acid (0.5 g) as a cream solid, m.p. 198-200°C. [Elemental analysis:- C,68.32; H,5.45; N,6.93%. Calculated :- C,68.89; H,5.30; N,6.69%]

EXAMPLE 87 Compound JO

20

To a stirred suspension of sodium hydride (0.69 g, 60% dispersion in mineral oil) in dry tetrahydrofuran (50 mL) is added a solution of sodium (RS)-5-hydroxy-5-(2-methylphenyl)pentanoate (1.23 g) in dry tetrahydrofuran (100 mL). The mixture is stirred at room temperature for 1.5 hours when
25 2-fluoro-4-(3-thienylmethoxy)benzonitrile (1.57 g) is added in one portion. The reaction mixture is heated at 55-60°C for 18 hours and evaporated. The residue is dissolved in water (100 mL), the solution acidified to pH 1 with 1 N hydrochloric acid and the mixture extracted three times with ethyl acetate (100 mL). The combined extracts are washed with brine (100 mL), dried over
30 magnesium sulphate and evaporated to give (RS)-5-[2-cyano-5-(3-thienylmethoxy)phenoxy]-5-(2-methylphenyl)pentanoic acid as a white waxy solid (0.4 g), mp.p.106-108°C. [Elemental analysis:- C,67.15; H,5.84; N,2.75; S,6.99%. Calculated for C₂₄H₂₃NO₄S·0.5EtOAc:- C,66.66; H,4.66; N,6.48%].

35 EXAMPLE 88 Compound JP

A solution of ethyl (RS)-4-(2-cyano-3-fluoro-5-(3-thienylmethoxy)-phenoxy)-4-(2-methylphenyl)butanoate (200 mg) in methanol (20 mL) containing 1 N sodium hydroxide (5 mL) is stirred at ambient temperature for 16 hours. The reaction mixture is poured into 0.5 N HCl (30 mL) and extracted
5 three times with ethyl acetate (30 mL). The combined organic phases are washed with brine (20 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a brown oil. Crystallisation with cyclohexane gives (RS)-4-(2-cyano-3-fluoro-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl) butanoic acid, (70 mg) in the form of a white solid, m.p. 150-
10 151°C. [Elemental analysis:- C, 64.9, H, 4.74, N, 3.29%. Calculated:- C, 64.8, H, 4.81, N, 3.13%]

EXAMPLE 89 Compound JQ

15 A solution of ethyl (RS)-2,4-dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)phenyl]butanoate (0.38 g) in 15% methanolic potassium hydroxide (25 mL) is stirred at reflux for 1 hour before being cooled and concentrated under reduced pressure. The residue is dissolved in water, acidified to pH 1 with concentrated hydrochloric acid and extracted with ethyl
20 acetate. The organic layer is dried over magnesium sulphate, filtered and concentrated under reduced pressure. The residue is recrystallised from cyclohexane to give (RS)-2,4-dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)-phenyl]butanoic acid (35 mg) as a yellow solid, m.p. 124-6 °C. [Elemental analysis:- C, 65.82; H, 5.03%; Calculated for C₂₄H₂₂O₆S:- C,
25 65.74; H, 5.06%].

EXAMPLE 90 Compound JR

To a stirred suspension of ethyl (RS)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (4 g) in 15%
30 potassium hydroxide in methanol (100 mL, w/v) at room temperature is added water (10 mL). The solid slowly dissolves and after stirring for 15 minutes the reaction mixture is partitioned between ethyl acetate (200 mL) and 2 N hydrochloric acid (200 mL). The organic phase is washed with water (200 mL),
35 dried over magnesium sulphate and evaporated. The residue is recrystallised from a mixture of ethyl acetate and cyclohexane to give (RS)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-

butanoic acid hydrate (1.9 g), m.p.167-168°C. [Elemental analysis:- C,63.9; H,5.39; N,5.22%. Calculated for $C_{28}H_{28}N_2O_6 \cdot 0.5H_2O$:- C,63.5; H,5.48; N,5.29%].

5 EXAMPLE 91 Compound JS

tert-Butyl (R)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate (1.7 g) is stirred in trifluoroacetic acid (10 mL) at ambient temperature for 15 minutes. The reaction mixture is
10 basified with saturated sodium bicarbonate solution and a solid is precipitated which is taken up in ethyl acetate (50 mL) and washed with water (50 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. Recrystallisation from ethyl acetate/cyclohexane affords
(R)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-
15 methylphenyl)butanoic acid (0.35 g) in the form of a white solid, m.p. 108-110°C. [Elemental analysis:- C,67.9, H, 5.92, N, 7.64%. Calculated:- C,67.6, H, 5.63, N, 8.16%]

EXAMPLE 92 Compound JT

20

A solution of (R)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (150 mg) in methanol (50 mL) containing 10% w/v potassium hydroxide (5 mL) is stirred at reflux for 1 hour. The reaction mixture is partitioned between 1 N acetic acid (50 mL)
25 and ethyl acetate (50 mL). Some precipitation occurred in the organic phase which is washed with water (50 mL) and concentrated under reduced pressure. Trituration with hot ethyl acetate followed by cooling affords a solid which is filtered then washed with ethyl acetate to give (R)-4-[2-(5-carboxypyrazol-3yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl) butanoic acid hydrate (40
30 mg) as a white solid, m.p. 144-146°C. [Elemental analysis:- C,64.1, H, 5.02, N, 8.27%. Calculated for $C_{27}H_{25}N_3O_6 \cdot H_2O$:- C,64.2, H, 5.35, N, 8.32%].

EXAMPLE 93 Compound JU

35

(RS)-2,2-difluoro-3-hydroxy-3-(2-methylphenyl)propanoic acid (1.5 g) is added portionwise to a stirred suspension of sodium hydride (1.1 g, 60% dispersion in mineral oil) in THF (80 mL) at ambient temperature under

nitrogen. After 20 minutes 2-fluoro-4-(3-pyridylmethoxy)benzonitrile (1.6 g) is added in one portion and the reaction stirred at 60°C for 16 hours. The reaction is concentrated in vacuo and the residue acidified with 1 N HCl to pH 1 and extracted with ethyl acetate (100 mL). A white solid precipitates from the ethyl acetate which is filtered off to give (RS)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]-2,2-difluoro-4-(2-methylphenyl)propanoic acid (0.64 g), m.p. 265-267°C. [Elemental analysis:- C,64.7, H, 4.19, N, 6.43%. Calculated:- C,65.1, H, 4.72, N, 6.60%]

10 EXAMPLE 94 Compound JV

A solution of ethyl (RS)-4-[2-(5-ethoxycarbonylisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2.56 g) in methanol (30 mL) is treated with 50% aqueous potassium hydroxide solution (10 mL, w/v) and the mixture is heated to reflux for 3 hours. The reaction mixture is acidified to pH 1 by addition of concentrated hydrochloric acid (50 mL). Water (75 mL) and ethyl acetate (75 mL) are added and the organic phase separated. The aqueous phase is further extracted with ethyl acetate (3 x 50 mL). The combined organic phases are dried over magnesium sulphate and evaporated. During the evaporation a white solid is formed. This is filtered and the remaining solution evaporated to dryness to give an orange oil (2.08 g). This material is partitioned between ethyl acetate (50 mL) and water (50 mL), the pH of the aqueous layer is adjusted to pH 9 by addition of potassium carbonate, and the aqueous layer is separated. Glacial acetic acid is added to the aqueous solution to adjust the pH to pH 4 and the orange oil collected by decantation. Trituration with a mixture of ethyl acetate and cyclohexane gives (RS)-4-[2-(5-carboxyisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (0.42 g) as a cream solid, m.p.162-164°C. [Elemental analysis:- C,62.98; H,4.70; N,2.65%. Calculated:- C,63.30; H,4.70; N,2.84%].

30

EXAMPLE 95 Compound JW

tert-Butyl (R)-4-[2-(5-methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2 g) is dissolved in trifluoroacetic acid (10 mL) and allowed to stand at room temperature for 2 minutes. The solution is quenched with saturated aqueous sodium hydrogen carbonate solution (100 mL) and the product extracted into ethyl acetate

35

(100 mL). This solution is washed with water, dried, and evaporated. The residue is purified by flash chromatography on silica eluting initially with a mixture of ethyl acetate and cyclohexane (3:2 v/v) to remove trace high running impurities, and then eluting with a mixture of methanol and dichloromethane (1:4 v/v). Fractions heterogeneous in the required product are combined and evaporated. The residue is recrystallised from a mixture ethyl acetate and cyclohexane to give (R)-4-[2-(5-methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (0.4 g) as a white solid, m.p. 155-156°C. [Elemental analysis: - C, 66.90; H, 5.49; N, 8.11%. Calculated: - C, 67.00; H, 5.43; N, 8.43%].

EXAMPLE 96 Compound JX

(R)-4-[2-(5-methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (0.2 g) is dissolved in methanol (20 mL) and concentrated aqueous ammonia (20 mL). After standing at room temperature overnight the solution is evaporated to dryness and the residue triturated successively with ether and ethyl acetate to give (R)-4-[2-(5-carbamoylpyrazol-3-yl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid (0.035 g) as a white powder, mp 180-192°C

EXAMPLE 97 Compound JY

A solution of ethyl (RS)-4-[(2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-cyanophenyl)butanoate (250 mg) in methanol (10 ml) containing 1N sodium hydroxide (3 mL) is stirred at ambient temperature for 16 hours. The reaction mixture is poured into 0.5 N hydrochloric acid (20 mL) and extracted twice with ethyl acetate (20 mL). The combined organic phases are washed with brine (20 mL), dried over magnesium sulphate and concentrated under reduced pressure to leave a dark yellow oil. Crystallization from a mixture of cyclohexane and ethyl acetate affords (RS)-4-[(2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-cyanophenyl)butanoic acid (20 mg) in the form of a beige solid, m.p. 132-134°C. [Elemental analysis: - C, 66.0, H, 4.87, N, 3.02%. Calculated: - C, 66.2, H, 4.86, N, 3.22%].

35

2-Hydroxy-4-(3-thienylmethoxy)acetophenone (2.5 g) is added portionwise stirred suspension of sodium hydride (440 mg, 60% dispersion in

mineral oil) in dry dimethyl formamide (30 mL). After stirring at ambient temperature for 30 minutes, ethyl (RS)-4-chloro-4-(2-cyanophenyl)butanoate (2.87 g) is added in one portion and the reaction mixture stirred for 16 hours at 100 °C. The reaction is concentrated in vacuo and the residue partitioned
5 between ethyl acetate (100 mL) and water (100 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. The residual oil is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2, v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-[2-acetyl-5-(3-
10 thienyl-methoxy)phenoxy]-4-(2-cyanophenyl)butanoate (250 mg) as an orange oil.

A solution of 2-cyanobenzaldehyde (5 g) in dichloromethane (10 mL) is added dropwise to a stirred 1 M solution of titanium (IV) chloride in
15 dichloromethane (38 mL) at 0°C under nitrogen. The mixture is stirred for 5 minutes and [1-ethoxycyclopropyloxy]trimethylsilane (8.42 mL) is added dropwise maintaining the temperature at 0-5°C. The mixture is stirred for 16 hours at ambient temperature then quenched with water (200 mL). The mixture is extracted three times with dichloromethane (100 mL) and the combined
20 organic extracts are washed with brine (100 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-chloro-4-(2-cyanophenyl)butanoate (5.5 g) as a yellow oil.

EXAMPLE 98 Compound JZ

25 Sodium (R)-4-Hydroxy-4-(2-methylphenyl)butanoate (0.39 g) is added in one portion to a stirred suspension of sodium hydride (0.11 g, 60% dispersion in mineral oil) in tetrahydrofuran (30 mL) and the mixture is stirred at 50°C for 1 hour. A solution of 2-fluoro-4-(isothiazol-4-ylmethoxy)benzonitrile (0.21 g) in
30 tetrahydrofuran (20 mL) is added and the reaction mixture is stirred at reflux for 16 hours. The reaction mixture is concentrated in vacuo and water is added to the residue. The solution is acidified with 1 N hydrochloric acid and extracted with ethyl acetate (50 mL). The organic phase is washed with water (50 mL), with brine (50 mL), dried over magnesium sulphate and evaporated. The
35 residue is purified by flash chromatography on silica eluting with a mixture of dichloromethane and methanol (19:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give (R)-4-[(2-cyano-5-

(isothiazol-4-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid, (0.02 g) in the form of a yellow oil. [NMR (CDCl₃) (300MHz): -2.2-2.4 (m, 2H), 2.45 (s, 3H), 2.6-2.65 (m), 2.75-2.8 (m, 1H), 5.0 (q, 2H), 5.5 (dd, 1H), 6.15 (d, 1H), 6.5 (dd, 1H), 7.2 (m, 3H), 7.4 (m, 1H), 7.45 (d, 1H), 8.45 (d, 2H)].

5

EXAMPLE 99 Compound KA

A mixture of ethyl 3-(2,4-dibenzyloxy)phenyl propanoate (1.4 g) and 15% w/v potassium hydroxide in methanol (25 mL) is refluxed for 2 hours. The reaction is concentrated in vacuo to give a yellow solid which is partitioned between ethyl acetate (50 mL) and water (50 mL). Acidification with 1 N HCl leads to the formation of a precipitate which is filtered off to give 3-(2,4-dibenzyloxy)phenyl propanoic acid (0.52 g) as a white solid, m.p. 142-144°C [Elemental Analysis:- C, 75.9, H, 6.06% Calculated for C, 76.2, H, 6.12%].

15

EXAMPLE 100 Compound KB

A mixture of ethyl 2,4-dibenzyloxyphenoxyacetate (2 g), 10% w/v potassium hydroxide (10 mL) and methanol (50 mL) is stirred at ambient temperature for 1 hour. The reaction is concentrated in vacuo and acidified with 1 N HCL then extracted with ethyl acetate (2 x 100 mL). The organic layer is washed with water (2 x 100 mL), dried over magnesium sulphate, filtered and concentrated in vacuo to give a white powder. Recrystallisation from ethyl acetate/cyclohexane affords 2,4-dibenzyloxyphenoxyacetic acid (0.4 g) as a white solid, m.p. 72°C [Elemental Analysis:- C, 72.4, H, 5.50% Calculated for C, 72.5 H, 5.53%].

25

EXAMPLE 101 Compound KC

30

A stirred suspension of sodium (R)-4-hydroxy-4-(2-methylphenyl)butanoate (190 mg) in THF (15 mL) is treated with 60% sodium hydride (35 mg) and maintained at 25° for 0.5 hours. The mixture is treated with 2-fluoro-4-(thiophen-3-ylmethoxy)nitrobenzene (220 mg) and maintained at 50°C for 3 hours. The solution is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried and evaporated. The residue is purified by

35

flash chromatography on silica, eluting with dichloromethane followed by ethyl acetate, followed by recrystallisation from ethyl acetate/pentane to give (R)-4-[2-nitro-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid (70 mg), a tan coloured solid, mp. 130-131°C. ¹H NMR (CHCl₃) 2.2-2.9 (4H,m), 2.4 (3H,s), 4.9 (2H, q), 5.5 (1H, q), 6.2 (1H,s), 6.5 (1H,d), 7.0-7.4 (7H,m), 7.9 (1H,d).

EXAMPLE 102 Compounds KD and KE

10 A stirred solution of (R,S)-4-[2-cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid (500 mg) in dichloromethane (80 mL) is treated with 1,1-carbonyldiimidazole (215 mg) and stirred at 25°C for 2 hours. This solution is added to a solution of the sodium salt of methanesulphonamide (which is prepared by adding 60% sodium hydride (201 mg) to a solution of
15 methanesulphonamide (684 mg) in DMF (50 mL). Stirred at 25°C for 18 hours. The solution is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried, and evaporated. The residual oil is purified by flash chromatography on silica, eluting with pentane/ ethyl acetate: 2/1, followed by trituration with pentane to give (R,S)-N-{4-[2-cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}methane sulphonamide
20 (500 mg), colourless solid, mp.78-80°C. [Elemental analysis: C,59.7; H,5.14; N,5.41%. Calculated for C₂₄H₂₄N₂O₅S₂: C,59.5; H,4.99; N,5.78%].

25 The following is prepared similarly according to the instant method:

(R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}trifluoromethane sulphonamide, colourless solid, mp.193-195°C. [Elemental analysis: C,53.3; H,4.29; N,5.78%. Calculated for C₂₄H₂₄N₂O₅S₂: C,53.5; H,3.93; N,5.20%].

30

EXAMPLE 103 Compound KF

A stirred solution of ethyl (R,S)-4-[2-(3-methoxycarbonylpropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1.8 g) in
35 dioxan (20 mL) and 2 M NaOH (5.15 mL) is heated at 40°C for 1.5 hours. The solution is evaporated. The residue is dissolved in water, brought to pH 1 with 1 M HCl, and extracted with ethyl acetate. The organic layer is washed with

- water, dried, and evaporated. The residue is recrystallised from ethyl acetate/cyclohexane to give (R,S)-4-[2-(3-carboxypropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid (1 g), a colourless solid, mp. 150-152°C. [Elemental analysis: C,65.0; H,5.52%. Calculated for C₂₆H₂₆O₇S: C,64.7; H,5.43%].

EXAMPLE 104 Compound KG

- A stirred solution of ethyl (R,S)-4-[2-(1,2,4-oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (61 mg) in dioxan (2 mL) and 1 M HCl (0.38 mL) is heated at 80°C for 5 hours. The solution is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried and evaporated. The residue is purified by flash chromatography on silica, eluting with ether/pentane: 3/1, to give (R,S)-4-[2-(1,2,4-oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid (18 mg), a colourless solid, mp. 145-149°C. ¹H NMR (CHCl₃) 2.2-3.0 (4H,m), 2.4 (3H,s), 4.9 (2H, q), 5.5 (1H, q), 6.2 (1H,s), 6.6 (1H,d), 7.0-7.4 (7H,m), 7.9 (1H,d), 8.8 (1H,s).

20 EXAMPLE 105 Compound KH

- A stirred solution of ethyl (R,S)-4-[2-(thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (500 mg) in dioxan (8.6 mL) and 1 M NaOH (2.5 mL) is heated at 60°C for 1.5 hours. The solution is evaporated. The residue is dissolved in water, brought to pH 1 with 1 M HCl, and extracted with ethyl acetate. The organic layer is washed with water, dried, and evaporated. The residual solid is triturated with ether giving (R,S)-4-[2-(thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid, a colourless solid (320 mg), mp. 174-176°C. [Elemental analysis: C,64.0; H,5.32; N, 2.62%. Calculated for C₂₅H₂₃NO₄S₂: C,64.5; H,4.98; N,3.01%].

REFERENCE EXAMPLE 1

- 35 A solution of methyl 2-hydroxy-4-(4-chlorobenzyloxy)benzoate (5.2 g) in dry dimethylformamide (DMF) (50 mL) is treated with NaH (60% dispersion, 0.71 g) and stirred at 25°C for 20 minutes. Benzyl bromide (2.12 mL) is added

and the solution heated at 60°C for 2 hours. The solution is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried, and evaporated. The residue is recrystallized from ethyl acetate to yield methyl 2-benzyloxy-4-(4-chlorobenzyloxy)benzoate as colourless crystals (5.1 g, 75%),
5 m.p. 99-101°C. ¹H NMR (DMSO) 3.8(3H,s), 5.18 (2H,s), 5.21(2H,s), 6.7(1H,d), 6.85(1H,s), 7.3-7.5(9H,m), 7.75(1H,d).

By proceeding in a similar manner, but using the appropriate amount of a hydroxy substituted benzoate precursor, there are prepared:

10 methyl 2-benzyloxy-4-(3-phenylpropyloxy)benzoate; methyl 2,4-di-(4-chlorobenzyloxy)benzoate; methyl 2-benzyloxy-4-(2-(3-indolyl)ethoxy)-benzoate (solid); methyl 2-(3-phenylpropyloxy)-4-benzyloxybenzoate, m.p. 50-51°C; methyl 2-benzyloxy-4-(2-naphthylmethoxy)benzoate (solid); methyl 2-benzyloxy-4-(1-naphthylmethoxy)benzoate; methyl 2-benzyloxy-4-(3,4-methylenedioxybenzyloxy)benzoate; methyl 2-(2-pyridylmethoxy)-4-benzyl-
15 oxybenzoate, m.p. 120-121°C; methyl 2-(2-phenylethoxy)-4-benzyloxybenzoate (solid); methyl 2-cyclohexylmethoxy-4-benzyloxybenzoate, (colourless solid); methyl 2-(4-chlorobenzyloxy)-4-benzyloxybenzoate, m.p. 95-97°C; methyl 2-benzyloxy-4-(2-phenylethoxy)benzoate; methyl 2-benzyl-oxy-4-(4-isopropylbenzyloxy)benzoate (solid); methyl 2-benzyloxy-4-(4-nitro-benzyloxy)benzoate, m.p. 132-134°C; methyl 2-benzyloxy-4-(4-fluorobenzyl-oxy)benzoate, m.p. 130-132°C; methyl 2-isopropoxy-4-benzyloxybenzoate (solid); methyl 2-(4-pyridylmethoxy)-4-benzyloxybenzoate, m.p. 75°C; methyl 2-
20 benzyloxy-4-(3,4-dichlorobenzyloxy)benzoate, m.p. 94-96°C; methyl 2-benzyloxy-4-(4-methoxybenzyloxy)benzoate; and methyl 2-benzyloxy-4-(3-methoxybenzyloxy)benzoate. m.p. 75-76°C.

REFERENCE EXAMPLE 2

30 A solution of methyl 2,4-dihydroxybenzoate (6.73 g) in acetone (250 mL) is treated with potassium iodide (2.66 g), tetrabutylammonium chloride (0.05 g), potassium carbonate (5.53 g), and 4-chlorobenzyl chloride (7.08 g), and stirred at reflux for 24 hours. The mixture is filtered, and the filtrate
35 evaporated. The residue is taken up in ethyl acetate, washed with water, dried, and evaporated. The residue is recrystallized from ethyl acetate to yield methyl 2-hydroxy-4-(4-chlorobenzyloxy)benzoate as colourless crystals

(5.3 g, 45%), m.p. 119-121°C. ¹H NMR (DMSO) 3.9(3H,s), 5.2(2H,s), 6.6(2H,m), 7.5(4H,m), 7.7(1H,d).

By proceeding in a similar manner, but using the appropriate amount of
5 a hydroxy substituted benzoate precursor, there are prepared:

methyl 2-hydroxy-4-(3-phenylpropyloxy)benzoate; methyl 2-hydroxy-4-(2-(3-indolyl)ethoxy)benzoate; methyl 2-hydroxy-4-benzyloxybenzoate, m.p. 101-102°C; methyl 2-hydroxy-4-(2-naphthylmethoxy)benzoate; methyl 2-hydroxy-4-
10 (1-naphthylmethoxy)benzoate; methyl 2-hydroxy-4-(3,4-methylenedioxybenzyloxy)benzoate (solid); methyl 2-hydroxy-4-(2-phenylethoxy)benzoate; methyl 2-hydroxy-4-(4-isopropylbenzyloxy)benzoate, m.p. 68-70°C; methyl 2-hydroxy-4-(4-nitrobenzyloxy)benzoate, m.p. 178-180°C; methyl 2-hydroxy-4-(4-fluorobenzyloxy)benzoate, m.p. 82-84°C; methyl 2-hydroxy-4-(3,4-
15 dichlorobenzyloxy)benzoate, m.p. 134-136°C; methyl 2-hydroxy-4-(4-methoxybenzyloxy)benzoate; and methyl 2-hydroxy-4-(3-methoxybenzyloxy)benzoate.

REFERENCE EXAMPLE 3

20 A mixture of triphenylphosphine (15.7 g) and carbon tetrabromide (9.93 g) in dry dichloromethane (100 mL) is cooled to 0°C and stirred for 30 minutes. A solution of 2,4-dibenzyloxybenzaldehyde (4.76 g) in dichloromethane (30 mL) is added and the solution stirred at 0°C for 2 hours. The solution is treated with pentane to precipitate triphenylphosphine oxide. The
25 filtrate is evaporated to give 1-(2,4-dibenzyloxyphenyl)-2,2-dibromoethylene as a yellow oil (4.85 g, 67%).

REFERENCE EXAMPLE 4

30 A solution of 2,4-dibenzyloxybenzoic acid (3.34 g) and methyl glycinate hydrochloride (1.25 g) in DMF (40 mL) is treated with triethylamine (2.23 g) and 1-hydroxybenzotriazole (1.53 g) at 0°C. The mixture is stirred for 10 minutes, treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, hydrochloride (2.05 g) and allowed to warm to 25°C. The solution is
35 evaporated. The residue is partitioned between ethyl acetate and water. The organic layer is washed with 1 M HCl, 1 M NaOH, water, dried and evaporated. The residue is recrystallized from methanol to yield methyl

N-(2,4-dibenzyloxybenzoyl)glycinate as a solid (900 mg). ¹H NMR (CDCl₃) 3.7(3H,s), 4.2(2H,d), 5.1 (2H,s), 5.2(2H,s), 6.6(1H,s), 6.7(1H,d), 7.3-7.5(10H,m), 8.2(1H,d), 8.4(1H,t).

5 REFERENCE EXAMPLE 5

A suspension of 4-benzyloxy-2-(1-o-tolyloxy)benzoyl chloride (2.73 g) and glycine methyl ester hydrochloride (1.21 g) in dichloromethane (50 mL) is treated with triethylamine (2.78 g) dropwise at 25°C. The mixture is stirred for 24 hours. The solution is washed with water, 1 M HCl, brine, dried and evaporated. The residual oil is purified by flash chromatography, eluting initially with dichloromethane and then with ether. Fractions homogeneous in the required product are combined and evaporated to yield a red gum. Trituration with cyclohexane/ether yields a solid. Recrystallization from ethyl acetate yields N-(4-benzyloxy-2-(1-o-tolyloxy)benzoyl)glycinate as a colourless solid (240 mg), m.p. 137-139°C. ¹H NMR (CDCl₃) 1.8(3H,d), 2.4(3H,s), 3.8(3H,s), 4.3(2H,d), 4.9(2H,s), 5.6(1H,q), 6.2(1H,s), 6.6(1H,d), 7.1-7.4(9H,m), 8.15(1H,d), 8.7(1H,t).

20 REFERENCE EXAMPLE 6

A solution of 4-benzyloxy-2-(1-o-tolyloxy)benzoic acid (2.5 g) in dichloromethane (50 mL) is treated with a 2 M solution of oxalyl chloride in dichloromethane (6.5 mL) and stirred at 25°C for 3.5 hours. The mixture is evaporated to give 4-benzyloxy-2-(1-o-tolyloxy)benzoyl chloride as a brown liquid (2.73 g).

REFERENCE EXAMPLE 7

A solution of methyl 4-benzyloxy-2-(1-o-tolyloxy)benzoate (2.8 g) in 15% KOH in MeOH (50 mL) is refluxed for 3 hours. The solution is evaporated, the residue treated with water, brought to pH 1 with concentrated HCl, extracted with dichloromethane. The extract is dried and evaporated to give 4-benzyloxy-2-(1-o-tolyloxy)benzoic acid as a colourless solid, m.p. 110-113°C.

REFERENCE EXAMPLE 8

A solution of triphenylphosphine (5.48 g) in dry THF (35 mL) is treated with diisopropyl azodicarboxylate (4.23 g) at 0-5°C and stirred for 15 minutes. The suspension is treated dropwise with a solution of methyl 4-benzyloxy-2-hydroxybenzoate (2.7 g) and 1-o-tolyethanol (1.42 g) in dry THF (35 mL) during 25 minutes at 0-5°C and then stirred at 0-5°C for 1 hour and at 25°C for 24 hours. The mixture is evaporated and the residual gum purified by flash chromatography on silica, eluting with dichloromethane. Fractions homogeneous in the required product are combined and evaporated to yield methyl 4-benzyloxy-2-(1-o-tolyethoxy)benzoate as a colourless oil (2.83 g, 72%).
¹H NMR (CDCl₃) 1.6(3H,d), 2.4(3H,s), 3.9(3H,s), 4.9(2H,s), 5.5(1H,q), 6.3(1H,s), 6.5(1H,d), 7.1-7.6(9H,m), 7.8(1H,d).

REFERENCE EXAMPLE 9

A solution of ethyl 4-(2,4-dibenzyloxyphenyl)-2,4-dioxobutanoate (3.23 g) in ethanol (10 mL) is treated with hydrazine hydrate (0.4 mL), refluxed for 2 hours and then cooled to room temperature. The mixture is filtered to yield ethyl 3-(2,4-dibenzyloxyphenyl)-pyrazole-5-carboxylate as a solid (1.5 g).

REFERENCE EXAMPLE 10

A suspension of potassium ethoxide (0.63 g) in ether (50 mL) is treated with diethyl oxalate (1.2 g), cooled to 0°C, treated with 2,4-dibenzyloxyacetophenone (2.5 g) and stirred at 25°C for 24 hours. The mixture is partitioned between ether and 0.1 M HCl. The ether layer is washed with water, dried, and evaporated to yield 4-(2,4-dibenzyloxyphenyl)-2,4-dioxobutanoate as an oil (3.23 g).

REFERENCE EXAMPLE 11

Ethyl 5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate (6 g) is added portionwise to a stirred suspension of sodium hydride (0.75 g) in tetrahydrofuran (250 mL) at room temperature. Stirring is continued at this temperature for a further 30 minutes when benzyl bromide (3.57 g) is added in one portion. The reaction mixture is heated to reflux for 5 hours, then

evaporated in vacuo. The residue is treated with water (250 mL), the resulting yellow solid washed well with water, then dissolved in tetrahydrofuran and filtered to remove a small amount of insoluble white solid. The filtrate is evaporated and the residual yellow solid purified by flash chromatography on silica eluting initially with dichloromethane then with a mixture of ethyl acetate and dichloromethane (1:9 v/v) to give ethyl 2-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate (1.2 g) and ethyl 1-benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate (2 g).

10 A suspension of ethyl 4-benzyloxyphenyl-2,4-dioxobutanoate (35.86 g) in ethanol (300 mL) is treated with hydrazine hydrate (6 mL) and the mixture is heated at reflux for 2 hours. The reaction mixture is cooled to room temperature, filtered, and the solid washed with ethanol to give ethyl 5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylate (25 g), m.p. 160-170°C.

15 Diethyl oxalate (17.7 g) is added in one portion to a stirred suspension of potassium ethoxide (10.2 g) in anhydrous ether (1000 mL). The stirred mixture is cooled to 0°C, treated with 4-benzyloxyacetophenone (25 g) portionwise, and stirring continued for 3 hours. After standing at room temperature for 18 hours the reaction mixture is poured onto 1 N hydrochloric acid (500 mL) and the organic phase separated. The organic phase is washed with water, dried over magnesium sulphate and evaporated to give ethyl 4-benzyloxyphenyl-2,4-dioxobutanoate (35 g).

25 A mixture of 4-hydroxyacetophenone (30 g), potassium carbonate (33.4 g), potassium iodide (1 g) and benzyl bromide (41.4 g) in methyl ethyl ketone (500 mL) is heated at reflux for 18 hours. The cooled reaction mixture is filtered and the filtrate evaporated. The residue is crystallised from cyclohexane to give 4-benzyloxyacetophenone (45 g, 90%).

30
REFERENCE EXAMPLE 12

A stirred solution of methyl benzoylamino(dimethoxyphosphoryl)acetate (4.25 g) in dry tetrahydrofuran (30 mL) is treated with sodium hydride (0.66 g, 60% dispersion in mineral oil) under nitrogen. After stirring for 15 minutes a solution of 2,4-dibenzyloxybenzaldehyde (4.74 g) in dry tetrahydrofuran

(15 mL) is added dropwise. The reaction mixture is stirred at room temperature for 18 hours, evaporated and the residue partitioned between ethyl acetate (100 mL) and water (75 mL). The organic phase is washed with brine (50 mL), dried over magnesium sulphate and evaporated. The residual yellow oil is
5 purified by flash chromatography on silica eluting with initially with dichloromethane then with a mixture of dichloromethane and ethyl acetate (9:1 v/v). Fractions homogeneous in the required products are evaporated to give methyl (E)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylate (1 g) as a yellow foam, and methyl (Z)-2-benzoylamino-3-(2,4-dibenzyloxyphenyl)-acrylate (1.6 g) as a
10 yellow oil.

A stirred solution of methyl amino(dimethoxyphosphoryl)acetate (6 g) in dichloromethane (50 mL) is treated with benzoyl chloride (5 g) and the mixture allowed to stand at room temperature for 4 hours. The reaction mixture is
15 washed twice with saturated aqueous sodium bicarbonate solution (75 mL), brine (50 mL), dried over magnesium sulphate and evaporated. The residual pale yellow oil is dissolved in ether (5 mL) and pentane added to give methyl benzoylamino(dimethoxyphosphoryl)acetate (4.25 g) as a white solid.

20 A solution of (\pm)-N-(benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester (20 g) in methanol (300 mL) is treated with palladium on carbon catalyst (5%, 2 g) and shaken under hydrogen at atmospheric pressure and room temperature for 18 hours. The mixture is filtered and evaporated giving methyl
amino(dimethoxyphosphoryl)acetate as an oil (12 g).

25

REFERENCE EXAMPLE 13

A stirred solution of triethyl 2-phosphono-3-phenylpropionate (5.17 g) in dry tetrahydrofuran (50 mL) is treated with a solution of *n*-butyl lithium in
30 hexanes (8 mL, 2.5 M) at -78°C. After stirring for 15 minutes a solution of 2,4-dibenzyloxybenzaldehyde (5 g) in dry tetrahydrofuran (50 mL) is added dropwise over 5 minutes at -78°C. The reaction mixture is stirred at -78°C for 1 hour and allowed to warm to room temperature. After standing at room
temperature for 18 hours the reaction mixture is evaporated and the residue
35 partitioned between ethyl acetate (120 mL) and water (75 mL). The organic phase is washed with brine (50 mL), dried over magnesium sulphate and evaporated. The residue is purified by flash chromatography on silica eluting

with dichloromethane. Fractions homogeneous in the required product are evaporated to give ethyl (E)-2-benzyl-3-(2,4-dibenzyloxyphenyl)acrylate (1.5 g) as a yellow oil.

5 REFERENCE EXAMPLE 14

A mixture of 2,4-dibenzyloxybenzaldehyde (1 g), methyl cyanoacetate (0.27 mL), piperidine (0.1 mL) and ethanol (30 mL) is stirred at ambient temperature for 20 minutes and the solvent is removed under reduced
10 pressure. The residue is recrystallized from ethanol to give methyl (E)-3-(2,4-dibenzyloxyphenyl)-2-cyanoacrylate as a yellow solid (400 mg).

REFERENCE EXAMPLE 15

15 A mixture of 2,4-dibenzyloxyacetophenone (33.2 g) and diethyl oxalate (21.9 g) in ethanol (250 mL) is treated with sodium (2.53 g) added in small portions over 2 hours. The resulting mixture is stirred at room temperature for 18 hours and filtered. The solid is washed with a little ethanol and then washed thoroughly with ether. A portion of this solid (5 g) and hydroxylamine
20 hydrochloride (0.84 g) are suspended in ethanol (100 mL) and 1 N hydrochloric acid is added until the pH of the mixture reaches pH 1. The stirred mixture is heated at reflux for 3 hours, during which time all the solids dissolve, then left at room temperature for 18 hours. The resulting solid is filtered and washed with ether to give ethyl 3-(2,4-dibenzyloxyphenyl)-isoxazole-5-
25 carboxylate (2.2 g).

REFERENCE EXAMPLE 16

30 Benzyl 2-benzyloxy-4-iodobenzoate (7.6 g) is added to triethylamine (120 mL) followed by phenylacetylene (2.94 g) then tetrakis(triphenylphosphine)palladium(0) (0.664 g) and copper (I) bromide (0.248 g). The resulting mixture is stirred at room temperature for 3 hours. The triethylamine is distilled off and the residue partitioned between ether (600 mL) and saturated aqueous ammonium chloride solution. The organic phase is dried over magnesium
35 sulphate and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of pentane and ethyl acetate (5:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give

benzyl 2-benzyloxy-4-phenylethynyl-benzoate (2 g) as a cream solid, m.p. 98-100°C.

- Sodium hydride (4.7 g, 60% dispersion in mineral oil) is added
5 portionwise over 15 minutes to a solution of 4-iodosalicylic acid (15.4 g) in dry dimethylformamide (150 mL). After stirring for 40 minutes benzyl bromide (19.9 g) is added and the mixture heated at 60°C for 1.5 hours. The reaction mixture is evaporated and the residue partitioned between chloroform and water. The organic phase is dried over magnesium sulphate, evaporated and
10 the residue triturated with ether to give benzyl 2-benzyloxy-4-iodobenzoate (14.2 g) as a cream coloured solid.

REFERENCE EXAMPLE 17

- 15 2,4-Dibenzyloxyphenol (3.1 g) is added to a stirred suspension of sodium hydride (0.44 g, 60% dispersion in mineral oil) and the mixture is heated at reflux for 30 minutes. Refluxing is allowed to abate when ethyl bromoacetate (1.8 g) is added and the mixture heated at reflux for 1 hour. The reaction mixture is evaporated and the residue partitioned between ethyl
20 acetate and water. The organic phase is washed with water, dried over magnesium sulphate and evaporated to give a light brown oil which is purified by column chromatography on silica eluting with dichloromethane. Fractions homogeneous in the required product are combined and evaporated. The residue is triturated with pentane to give ethyl (2,4-dibenzyloxyphenoxy)-
25 acetate (3.2 g) as a white powder, m.p. 62-64°C. [Elemental analysis:- C,73.3; H,6.13%. Calculated :- C,73.5; H,6.13%].

REFERENCE EXAMPLE 18

- 30 A solution of methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-methylphenyl)-pentanoate (46 g) in methanol (250 mL), methyl acetate (250 mL) and concentrated hydrochloric acid (25 mL) is treated with palladium on carbon catalyst (5%; 1.5 g) and shaken under hydrogen at atmospheric pressure and room temperature for 2 hours. The suspension is filtered, and
35 evaporated. The residue is dissolved in ethyl acetate, washed with water, dried and evaporated, and the residue is recrystallised from a mixture of ethyl

acetate and cyclohexane, giving methyl (RS)-5-(3-hydroxyphenyl)-5-oxo-4-(2-methylphenyl)pentanoate as colourless crystals (21.7 g), m.p. 101-103°C.

REFERENCE EXAMPLE 19

5

A suspension of magnesium turnings (9 g) in dry diethyl ether (100 mL) containing a few crystals of iodine is treated with a solution of benzyl chloride (22.8 g) in diethyl ether (150 mL) at such a rate in the form of to maintain gentle reflux. After the addition is complete, the mixture is treated with 3-
10 benzyloxybenzonitrile (12.7 g) and heated to reflux for 5 hours. The mixture is then treated with hydrochloric acid (1 N) until it is at pH 1, and stirring is continued for 1 hour at ambient temperature. The reaction mixture is separated and the aqueous layer extracted with diethyl ether (3 x 100 mL). The combined organic extracts are washed with water (3 x 100 mL), dried over magnesium
15 sulphate and concentrated under reduced pressure. The resulting residue is recrystallised from methanol, to give benzyl 3-benzyloxyphenyl ketone (14.8 g), in the form of a white solid, m.p. 58-60°C. [Elemental analysis:- C,76.2; H,4.99%; Calculated:- C,76.28;H,4.99%].

20 By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared:- 1-[3-(2-methoxy-benzyloxy)phenyl]-2-phenylethanone; 1-[3-(3,4-methylenedioxybenzyloxy)-phenyl]-2-phenylethanone; and 1-(3-phenoxyphenyl)-2-phenylethanone.

25 REFERENCE EXAMPLE 20

A stirred solution of 3-cyanophenol (20 g) in dry N,N-dimethylformamide (200 mL) is treated with sodium hydride (5.6 g; 60% dispersion in mineral oil), portionwise during 40 minutes, followed by dropwise addition of benzyl
30 bromide (20 mL). The mixture is heated at 90°C for 4 hours, cooled and concentrated under reduced pressure. The residue is partitioned between ethyl acetate and water and the organic layer washed with 1 N sodium hydroxide (100 mL) followed by drying over magnesium sulphate and concentration. The residue is recrystallised from methanol to give
35 3-benzyloxybenzonitrile (24 g), in the form of a cream solid, m.p. 38-40°C.

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared :

- 3-(2-methoxybenzyloxy)benzonitrile; and 3-(3,4-methylenedioxybenzyloxy)-
5 benzonitrile.

REFERENCE EXAMPLE 21

- A stirred solution of benzyl 3-benzyloxyphenyl ketone (3.02 g) in
10 tetrahydrofuran (40 mL) at 0°C is treated with potassium tert-butoxide (56 mg),
followed by acrylonitrile (0.66 mL). The mixture is stirred at room temperature
overnight and then concentrated under reduced pressure and partitioned
between water and ethyl acetate. The organic layer is dried over magnesium
sulphate and evaporated, to give a brown oil (4.3 g), which is subjected to flash
15 chromatography on silica gel, eluting with a mixture of ethyl acetate and
pentane (1:4v/v), to give 1-(3-benzyloxyphenyl)-4-cyano-2-phenylbutanone
(2.2 g), in the form of a light green oil. [Elemental analysis:- C,81.3;H,6.06;
N,3.65%; Calculated:- C,81.1;H,5.96;N,3.94%].

20 REFERENCE EXAMPLE 22

- A stirred solution of 1-(3-benzyloxyphenyl)-2-(2-methylphenyl)ethan-1-
one (4 g) in tetrahydrofuran (100 mL) is treated with potassium tert-butoxide
(200 mg). After stirring for 30 minutes, it is treated with bromoacetonitrile
25 (0.98 mL), dropwise, and stirring is continued for a further 28 hours. The
mixture is then treated with further quantities of potassium tert-butoxide
(200 mg) and bromoacetonitrile (0.98 mL) and allowed to stand at room
temperature overnight. It is then concentrated under reduced pressure and
partitioned between water and ethyl acetate. The organic layer is dried over
30 magnesium sulphate and evaporated, to give a brown oil, which is subjected to
flash chromatography on silica gel, eluting with a mixture of ethyl acetate and
pentane (1:9 v/v), to give 1-(3-benzyloxyphenyl)-3-cyano-2-(2-methyl-
phenyl)propan-1-one, in the form of a colourless oil (0.67 g).

35 REFERENCE EXAMPLE 23

A mixture of 2,4-dihydroxyacetophenone (29 g), 3-(chloromethyl)thiophene (30 g), potassium carbonate (32 g) and potassium iodide (0.1 g) in 2-butanone (250 mL) is stirred at reflux overnight. The reaction mixture is partitioned between ethyl acetate (300 mL) and dilute hydrochloric acid (300 mL; 1 M), and the organic layer is washed with water, dried, and evaporated. Crystallisation of the residue from cyclohexane gives 2-hydroxy-4-(3-thienylmethoxy)acetophenone (21 g) in the form of a white solid.

By proceeding in a similar manner, but replacing 3-(chloromethyl)thiophene with 3-(chloromethyl)pyridine, there is prepared 2-hydroxy-4-(3-pyridylmethoxy)acetophenone, in the form of a white solid.

REFERENCE EXAMPLE 24

A mixture of methyl 2,4-dihydroxybenzoate (50 g), benzyl chloride (37 mL), tetrabutylammonium bromide (0.25 g), potassium carbonate (44.4 g) and potassium iodide (18.4 g) in acetone is stirred at reflux overnight. The reaction mixture is evaporated to dryness and the residue is partitioned between water (300 mL) and ethyl acetate (300 mL). The organic layer is washed with aqueous sodium hydrogen carbonate solution and water, dried and evaporated. Crystallisation of the residue from ethyl acetate gives methyl 4-benzyloxy-2-hydroxybenzoate (45.1 g), in the form of a white solid.

By proceeding in a similar manner, but replacing benzyl chloride by 3-(chloromethyl)thiophene and performing the reaction at room temperature, there is prepared methyl 2-hydroxy-4-(3-thienylmethoxy)benzoate.

REFERENCE EXAMPLE 25

A mixture of methyl 4-benzyloxy-2-hydroxybenzoate (20 g) and concentrated aqueous ammonia (30 mL) in ethanol (100 mL) is heated on the steam bath in a sealed pressure vessel overnight. The contents of the pressure vessel are dissolved in methanol and the solution is evaporated to dryness. The residue is subjected to flash chromatography, eluting first with dichloromethane to remove unreacted starting material, and then with a mixture of ethyl acetate and dichloromethane (3:7 v/v), to obtain 2-hydroxy-4-benzyloxybenzamide (4 g), in the form of a white solid.

By proceeding in a similar manner, but replacing concentrated aqueous ammonia with concentrated aqueous ethylamine, there is prepared 4-benzyloxy-N-ethyl-2-hydroxybenzamide.

5

By proceeding in a similar manner, but replacing concentrated aqueous ammonia with ethanolic dimethylamine, there is prepared 4-benzyloxy-N,N-dimethyl-2-hydroxybenzamide.

10 By proceeding in a similar manner, but replacing methyl 4-benzyloxy-2-hydroxybenzoate by methyl 4-(3-thienylmethoxy)-2-hydroxybenzoate and concentrated aqueous ammonia by ethanolic ammonia solution, there is prepared 2-hydroxy-4-(3-thienylmethoxy)benzamide.

15 REFERENCE EXAMPLE 26

A stirred solution of 2,4-dihydroxyacetophenone (1.03 g) in dry acetone (40 mL) is treated with potassium carbonate (1.88 g), potassium iodide (1.13 g), tetrabutylammonium bromide (0.22 g), and 5-chloromethylpyrimidine (0.95 g) and heated at reflux for 18 hours. The suspension is filtered, the filtrate is evaporated and the resulting residual solid is recrystallised from a mixture of cyclohexane and ethyl acetate to give 2-hydroxy-4-(5-pyrimidinylmethoxy)-acetophenone (0.65 g) in the form of a colourless solid, m.p. 133-134°C. [NMR(CDCl₃):- 2.6(3H,s),5.1(2H,s),6.5-7.7(3H),8.7(2H,s),9.2(1H,s),
25 12.7(1H,s)].

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there is prepared 2-(2-pyridyl)-5-(3-thienylmethoxy)phenol, m.p. 115-117°C.

30

REFERENCE EXAMPLE 27

A mixture of 2-(2,4-dimethoxyphenyl)pyridine (14 g) in pyridine hydrochloride (121 g) is heated to 160°C and the resulting melt is heated at
35 160°C for 7 hours. The mixture is then cooled, diluted with dichloromethane, washed with hydrochloric acid (2 M) and water, dried and evaporated. The residual oil is subjected to flash chromatography on silica gel, eluting with a

mixture of pentane and ethyl acetate (1:1 v/v), to give 2-(2,4-dihydroxyphenyl)pyridine (8.0 g) in the form of a solid, m.p. 170-172°C. [NMR(CDCI₃):- 6.3(1H,s),6.4(1H,d),7.3(1H,m), 7.8(1H,d),7.9(1H,t),8.0(1H,d),8.5(1H,d).

5

REFERENCE EXAMPLE 28

A stirred suspension of magnesium (8.11 g) in dry tetrahydrofuran (100 mL) is heated, and then treated with a solution of 1-bromo-2,4-dimethoxybenzene (48 mL) in tetrahydrofuran (100 mL) at such a rate as to maintain reflux. The resulting solution is then cooled and added, dropwise, to a solution of 2-bromopyridine (50.3 g) and bis-triphenylphosphinepalladium(II) chloride (3.5 g) in tetrahydrofuran (100 mL) at reflux, and the mixture is maintained at reflux for 6 hours. The solution is then evaporated and the residue is dissolved in ethyl acetate, washed with water, dried and evaporated to give a dark oil. This is subjected to flash chromatography on silica gel, eluting with a mixture of pentane and ethyl acetate (2:1v/v) to give 2-(2,4-dimethoxyphenyl)pyridine (28 g) in the form of an oil. [NMR(CDCI₃):- 3.84(3H,s),3.86(3H,s), 6.55(1H,s),6.6(1H,d),7.2(1H,m),7.7(1H,t),7.77 (1H,d),7.8(1H,d),8.7(1H,d)].

20

REFERENCE EXAMPLE 29

A stirred solution of titanium(IV) chloride in dichloromethane (95 mL; 1 M) at 0°C under nitrogen is treated, dropwise, with a solution of 2-chloro-6-fluorobenzaldehyde (15 g) in dichloromethane (20 mL). The mixture is stirred for 5 minutes, and then treated, dropwise, with (1-ethoxycyclopropyloxy)-trimethylsilane (21.1 mL), maintaining the temperature at between 0°C and 5°C. The mixture is stirred for 16 hours at room temperature, and then it is treated with water (200 mL) and extracted with dichloromethane (3 x 100 mL), and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate, and evaporated, to give (RS)-4-chloro-4-(2-chloro-6-fluorophenyl)butanoate, in the form of a yellow oil.

By proceeding in a similar manner, but replacing the 2-chloro-6-fluorobenzaldehyde with the appropriate quantity of 2-bromobenzaldehyde, there is prepared ethyl (RS)-4-(2-bromophenyl)-4-chlorobutanoate, in the form of a yellow oil.

35

REFERENCE EXAMPLE 30

A stirred mixture of 5-benzyloxy-2-mercaptophenol (10 g), potassium
5 carbonate (6 g) and acetone (200 mL) at 0°C is treated, dropwise, with
iodomethane (6.1 g) and stirred for 2 hours at 0°C. The reaction mixture is
evaporated to dryness and the residue is partitioned between ethyl acetate
(200 mL) and water (200 mL). The organic layer is dried over magnesium
sulphate, filtered and concentrated in vacuo to give 5-benzyloxy-2-methylthio-
10 phenol, in the form of a brown oil.

By proceeding in a similar manner, but replacing the 5-benzyloxy-2-
mercaptophenol with the appropriate quantity of 2-mercapto-5-(3-thienyl-
methoxy)phenol, there is prepared 2-methylthio-5-(3-thienylmethoxy)phenol, in
15 the form of a yellow oil, after flash chromatography on silica gel, using a
mixture of petroleum ether and ethyl acetate (9:1 v/v) as eluent.

REFERENCE EXAMPLE 31

20 A mixture of 6-benzyloxy-1,3-benzoxathiol-2-one (35 g), aqueous
sodium hydroxide solution (200 mL; 2 N) and methanol (50 mL) is stirred at
ambient temperature for 16 hours and then it is heated at reflux for 30 minutes.
The reaction mixture is then evaporated to dryness and the residue is acidified
to pH 1 by treatment with concentrated hydrochloric acid (20 mL). The residue
25 is diluted with water (200 mL) and then extracted with diethyl ether
(3 x 200 mL). The combined organic extracts are washed with brine (100 mL),
dried over magnesium sulphate, filtered and concentrated in vacuo, to give
5-benzyloxy-2-mercaptophenol, in the form of a yellow oil which solidified on
standing overnight.

30

By proceeding in a similar manner, but replacing the 6-benzyloxy-1,3-
benzoxathiol-2-one with the appropriate quantity of 6-(3-thienylmethoxy)-1,3-
benzoxathiol-2-one, there is prepared 5-(3-thienylmethoxy)-2-mercaptophenol.

35 A mixture of 6-hydroxy-1,3-benzoxathiol-2-one (15 g) and potassium
carbonate (13.8 g) in dimethylformamide (200 mL) is stirred for 20 minutes, and
is then treated, dropwise, with 3-chloromethylthiophene (11.8 g). The mixture

is stirred at 60°C for 16 hours. The reaction mixture is then concentrated in vacuo, and partitioned between water (200 mL) and ethyl acetate (200 mL). The aqueous layer is extracted with ethyl acetate (100 mL) and with diethyl ether (100 mL), and the combined aqueous layers are washed with brine, dried
5 over magnesium sulphate, and evaporated to give a yellow solid, which is recrystallised from ethanol (200 mL) to give 6-(3-thienylmethoxy)-1,3-benzoxathiol-2-one (8.9 g), in the form of pale yellow crystals.

By proceeding in a similar manner, but replacing the 3-
10 chloromethylthiophene by the appropriate quantity of benzyl chloride, there is prepared 6-benzyloxy-1,3-benzoxathiol-2-one.

REFERENCE EXAMPLE 32

15 A solution of 2,4-dihydroxybenzaldehyde (42.6 g) in dimethylformamide (250 mL) at ambient temperature is treated with sodium hydride (13.5 g; 60% w/v dispersion in mineral oil; 338 mmol), portionwise during 30 minutes. It is then treated with 3-chloromethylthiophene and the reaction mixture is heated at 60°C for 3 hours. The mixture is concentrated in vacuo and the residue is
20 partitioned between hydrochloric acid (200 mL; 0.5 N) and ethyl acetate (200 mL). The aqueous layer is extracted with ethyl acetate (3 x 100 mL) and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate, filtered and the solvent is removed in vacuo. Flash chromatography on silica gel, using a mixture of petroleum ether (b.p. 40-60°C)
25 and ethyl acetate (4:1 v/v) as eluent, gives 2-hydroxy-4-(3-thienylmethoxy)-benzaldehyde (11.8 g), in the form of a white solid.

By proceeding in a similar manner, but using the appropriate quantities of the corresponding starting materials, there are prepared 4-benzyloxy-2-
30 hydroxybenzaldehyde and 5-(3-thienylmethoxy)-2-trifluoroacetylphenol.

REFERENCE EXAMPLE 33

By proceeding in a manner similar to that described hereinbefore in
35 Reference Example 24 but replacing the methyl 4-benzyloxy-2-hydroxybenzoate by the appropriate quantity of ethyl 2-hydroxy-6-methyl-4-(3-thienylmethoxy)benzoate, and the concentrated aqueous ammonia solution by

ethanolic ammonia solution, there is prepared 2-hydroxy-6-methyl-4-(3-thienylmethoxy)benzamide.

REFERENCE EXAMPLE 34

5

By proceeding in a manner similar to that described hereinbefore in Reference Example 23 but replacing the methyl 2,4-dihydroxybenzoate by the appropriate quantity of ethyl 2,4-dihydroxy-6-methylbenzoate, there is prepared ethyl 2-hydroxy-6-methyl-4-(3-thienylmethoxy)benzoate.

10

REFERENCE EXAMPLE 35

A mixture of 2-fluoro-4-(3-thienylmethoxy)nitrobenzene (4.21 g), aqueous potassium hydroxide solution (50 mL; 50% w/v) and *t*-butanol (20 mL) is heated at reflux for 4 hours. It is then evaporated to dryness and the residue is partitioned between ethyl acetate (100 mL) and hydrochloric acid (100 mL; 1 N). The organic phase is washed with water (3 x 100 mL), dried over magnesium sulphate, filtered and evaporated, to give 2-nitro-5-(3-thienylmethoxy)phenol, in the form of a yellow solid.

20

REFERENCE EXAMPLE 36

A mixture of 1-[N-(methoxycarbonylmethyl)carbamoyl]-2-[2-(trimethylsilyl)ethoxy]methoxy-4-(3-thienylmethoxy)benzene (3.3 g), tetrabutylammonium fluoride (20 mL of a 1 M solution in tetrahydrofuran) and tetrahydrofuran (30 mL) is stirred at ambient temperature for one hour and at reflux for 45 minutes, then it is cooled to ambient temperature and concentrated in vacuo. The resulting residue is partitioned between ethyl acetate (100 mL) and saturated brine (100 mL), and the organic layer is washed with brine (2 x 50 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. Flash chromatography of the residual oil on silica gel, eluting with a mixture of methanol and dichloromethane (2:98 v/v) gives 2-[N-(methoxycarbonylmethyl)carbamoyl]-5-(3-thienylmethoxy)phenol (2.3 g), in the form of a white solid.

30

By proceeding in a similar manner, but replacing the 1-[N-(methoxycarbonylmethyl)-carbamoyl]-2-[2-(trimethylsilyl)ethoxy]methoxy-4-(3-thienyl-

methoxy)benzene used as starting material by the appropriate quantities of the corresponding starting materials, there are prepared;

2-[N-(3-imidazol-1-ylpropyl)carbamoyl]-5-(3-thienylmethoxy)phenol, in the form of a yellow oil; 2-[N-(2-methoxycarbonylethyl)carbamoyl]-5-(3-thienylmethoxy)-phenol, in the form of a yellow oil; 2-[N-(2-cyanoethyl)carbamoyl]-5-(3-thienylmethoxy)phenol, in the form of a yellow oil; and 2-[N-(cyanomethyl)carbamoyl]-5-(3-thienylmethoxy)phenol, in the form of a yellow oil

10 REFERENCE EXAMPLE 37

A mixture of 4-(3-thienylmethoxy)-2-[2-(trimethylsilyl)ethoxy]methoxybenzoic acid (3.09 g), methyl aminoacetate hydrochloride (0.99 g), 1-hydroxybenzotriazole (1.28 g), triethylamine (0.8 g) and dimethylformamide (50 mL) is stirred at ambient temperature for 16 hours. It is then concentrated in vacuo and the resulting residue is dissolved in ethyl acetate (100 mL), washed quickly with hydrochloric acid (2 x 50 mL; 1 N), and then washed with saturated aqueous sodium bicarbonate solution (50 mL) and with water (50 mL), then dried over magnesium sulphate, filtered and concentrated in vacuo, to give 1-[N-(methoxycarbonylmethyl)carbamoyl]-4-(3-thienylmethoxy)-2-[2-(trimethylsilyl)ethoxy]methoxybenzene, in the form of a yellow oil.

By proceeding in a similar manner, but replacing the methyl aminoacetate hydrochloride by the appropriate quantities of the corresponding starting materials, there are prepared:-

1-[N-(3-imidazol-1-ylpropyl)carbamoyl]-4-(3-thienylmethoxy)-2-[2-(trimethylsilyl)ethoxy]methoxybenzene, in the form of a yellow oil; 1-[N-(2-methoxycarbonylethyl)carbamoyl]-4-(3-thienylmethoxy)-2-[2-(trimethylsilyl)ethoxy]methoxybenzene, in the form of a yellow oil; 1-[N-(cyanomethyl)carbamoyl]-4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxy)methoxybenzene, in the form of a yellow oil; 1-[N-(cyanoethyl)carbamoyl]-4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxy)methoxybenzene, in the form of a yellow oil; methyl 2-[2-(trimethylsilyl)ethoxy-4-(3-thienylmethoxy)benzoylamino]-3-phenylpropionate, thought to be the (R)-enantiomer, in the form of a yellow oil; and methyl 2-[2-(trimethylsilyl)ethoxy-4-(3-thienylmethoxy)benzoylamino]-3-phenylpropionate, thought to be the (S)-enantiomer, in the form of a yellow oil.

REFERENCE EXAMPLE 38

A mixture of methyl 4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxy)-methoxybenzoate (12.1 g) and aqueous potassium carbonate solution (50 mL; 10% w/v) in methanol (200 mL) is heated at reflux for 7 hours. It is then concentrated in vacuo and the residue is treated with water (200 mL) and acidified to pH 1 by treatment with hydrochloric acid (1 N), and the resulting solid is quickly filtered off and is then washed with water. The white solid is then azeotroped with cyclohexane in a Dean-Stark apparatus until no more water is removed. The solid is filtered off, to give 4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxy)methoxybenzoic acid (6.03 g), in the form of a white solid.

REFERENCE EXAMPLE 39

A stirred suspension of sodium hydride (1.53 g; 60% w/v dispersion in mineral oil; 38.3 mmol) in dimethylformamide (100 mL) is treated, portionwise, with methyl 2-hydroxy-4-(3-thienylmethoxy)benzoate (8.44 g) and stirred for 40 minutes. The mixture is treated with 2-(trimethyl-silyl)ethoxymethyl chloride (6.39 g), in one portion, and stirred at ambient temperature for three hours. It is then evaporated to dryness and the resulting residue is partitioned between water (100 mL) and ethyl acetate (100 mL). The organic layer is washed with brine (150 mL), dried over magnesium sulphate, filtered and concentrated in vacuo, to give methyl 4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxy)methoxybenzoate, in the form of a yellow oil.

REFERENCE EXAMPLE 40

A solution of triphenylphosphine (20 g) in tetrahydrofuran (300 mL) cooled at 0°C is treated dropwise during 5 minutes with diisopropyl azodicarboxylate (15 mL) and stirred for 15 minutes at 0°C. It is then treated, dropwise, with a solution of 2-fluoro-4-hydroxybenzonitrile (7 g) and 3-thienyl-methanol (7 g) in tetrahydrofuran (200 mL) during 2 hours, and stirring is continued at ambient temperature for 18 hours. The mixture is concentrated in vacuo and the residue is dissolved in ethyl acetate (500 mL). This solution is washed with water (200 mL) and aqueous sodium hydroxide solution (1 N), dried over magnesium sulphate and concentrated in vacuo, to give an oil, which is passed through a pad of silica gel (thickness 5 cm) with the aid of

dichloromethane. The resulting filtrate is concentrated in vacuo, to give an oil which crystallizes on standing. Recrystallization from isopropyl ether gives 2-fluoro-4-(3-thienyl-methoxy)benzonitrile (6 g), in the form of a white solid, m.p. 63-65°C.

5

REFERENCE EXAMPLE 41

A solution of triphenylphosphine (20 g) in tetrahydrofuran (300 mL) cooled at 0°C is treated dropwise during 5 minutes with diisopropyl
10 azodicarboxylate (15 mL) and stirred for 15 minutes at 0°C. It is then treated, dropwise, with a solution of 2-fluoro-4-hydroxybenzonitrile (7 g) and 3-pyridyl-methanol (7 g) in tetrahydrofuran (200 mL) during 2 hours, and stirring is continued at ambient temperature for 18 hours. The mixture is concentrated in vacuo and the residue is dissolved in ethyl acetate (500 mL) and the solution is
15 washed with hydrochloric acid (3 x 100 mL; 1 N) and the remaining organic layer is discarded. The combined acid washings are basified to pH 10 by treatment with solid sodium hydroxide and extracted with ethyl acetate (3 x 100 mL). This organic extract is dried over magnesium sulphate and concentrated in vacuo, and the residue is recrystallized from isopropyl ether, to
20 give 2-fluoro-4-(3-pyridylmethoxy)benzonitrile (10 g), in the form of a white solid, m.p. 105-110°C.

REFERENCE EXAMPLE 42

25 A mixture of 4-(2-methylphenyl)-4-oxobutanoic acid (40 g), concentrated sulphuric acid (4 mL) and t-butanol (4 mL) in dichloromethane (200 mL) is cooled to 0°C, and isobutylene (about 400 mL) is condensed into the mixture. It is then stirred at 0°C until t.l.c. shows reaction is complete. The excess isobutylene is allowed to evaporate, and then the mixture is treated carefully
30 with saturated aqueous sodium bicarbonate solution (200 mL), with vigorous stirring. The organic layer is separated, filtered through a pad of silica gel with the aid of more dichloromethane, and concentrated in vacuo, to give *t*-butyl 4-(2-methylphenyl)-4-oxobutanoate (5 g), in the form of a pale yellow oil.

35 REFERENCE EXAMPLE 43

A solution of *t*-butyl 4-(2-methylphenyl)-4-oxobutanoate (12 g) in dry diethyl ether (50 mL) is dried over molecular sieves (4 Å; 10 g) for 3 hours, and then it is transferred to a dry flask and placed under an argon atmosphere. The solution is cooled to -20°C, treated rapidly with bis[(1*S*,2*R*,3*S*,5*S*)-pinan-3-yl]chloroborane (25 g) and sealed under an atmosphere of argon. The reaction mixture is stored at -20°C for 48 hours, and then it is warmed to ambient temperature. It is then diluted with diethyl ether (50 mL) and treated with diethanolamine (10 g). The mixture is stirred vigorously at ambient temperature for 5 hours, then it is poured into pentane (500 mL) and filtered through a pad of diatomaceous earth. The filtrate is concentrated in vacuo at 0.5 mmHg/50°C for 3 hours, and the resulting residue is subjected to flash chromatography on silica gel, eluting with a mixture of methanol and dichloromethane (1:19 v/v), to give *t*-butyl 4-hydroxy-4-(2-methylphenyl)-butanoate, thought to be the (*R*)-enantiomer, (10 g) in the form of a colourless oil. $^{25}[\alpha]_D = +49^\circ$ ($c=0.01$; CHCl₃); [NMR(CDCl₃) (300MHz):- 1.45(s,9H), 2.00(m,2H), 2.32(s,3H), 2.40(m,2H), 4.99(m,1H), 7.10-7.50(m,4H).

The enantiomeric excess is determined to be greater than 99%, by reacting the product with (*R*)-(-)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride and considering the NMR (CDCl₃) spectrum of the resulting ester.

By proceeding in a similar manner, but replacing the bis[(1*S*,2*R*,3*S*,5*S*)-pinan-3-yl]-chloroborane by an appropriate quantity of bis[(1*R*,2*S*,3*R*,5*R*)-pinan-3-yl]chloroborane, there is prepared *tert*-butyl 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (*S*)-enantiomer, in the form of a colourless oil, $^{25}[\alpha]_D = -45^\circ$ ($c=0.02$; CHCl₃).

REFERENCE EXAMPLE 44

A solution of *t*-butyl 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (*R*)-enantiomer, (0.5 g) in tetrahydrofuran (3 mL) and methanol (3 mL) is treated with aqueous sodium hydroxide solution (5 N; 0.5 mL) and stirred at ambient temperature for 8 hours. The reaction mixture is concentrated to dryness. The resulting residue is treated with isopropanol (10 mL) and stirred for 30 minutes, and the resulting precipitate is collected and washed with isopropyl ether, to give sodium 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (*R*)-enantiomer, (300 mg) in the form of a white solid, m.p.

above 250°C; $^{25}[\alpha]_D = +34^\circ$ ($c=0.01$; CH₃OH); [NMR(DMSO-d₆) (300MHz):-
1.58-1.75 (m,2H),2.11(m,2H),2.22(s,3H),4.75(m,1H),7.05-7.48 (m.4H).

REFERENCE EXAMPLE 45

5

A mixture of 3-fluoro-4-nitrophenol (5.0 g) 3-chloromethylthiophene (4.2 g) and potassium carbonate (4.4 g) in dimethylformamide (25 mL) is stirred at 90°C for 16 hours. The solvent is removed in vacuo and the residue is partitioned between ethyl acetate (50 mL) and water (50 mL). The organic
10 phase is washed with water (2 x 50 mL), dried over magnesium sulphate, and evaporated to give 3-fluoro-4-nitrophenyl-5-(3-thienylmethoxy)benzene (7.66 g), in the form of a yellow solid.

REFERENCE EXAMPLE 46

15

A stirred solution of 2,4-dihydroxybenzaldehyde (31.8 g) in dry dimethylformamide (150 mL) at ambient temperature is treated with sodium hydride (11.96 g; 60% w/v dispersion in mineral oil; 300 mmol), portionwise, during 30 minutes. After a further period of 15 minutes, the mixture is treated,
20 dropwise, with a solution of 3-chloromethylthiophene (35 g) in dimethylformamide (50 mL). The mixture is stirred at 70°C for 2 hours, then it is cooled and the solvent is evaporated. The residue is partitioned between ethyl acetate (200 mL) and hydrochloric acid (200 mL; 0.5 M), and the aqueous layer is extracted further with ethyl acetate (2 x 100 mL). The combined organic
25 extracts are washed with brine (100 mL), dried over magnesium sulphate, and concentrated in vacuo. The residue is subjected to flash chromatography on silica gel, eluting with a mixture of ethyl acetate and pentane (1:4 v/v), to give a colourless oil, which crystallises on standing. This solid is recrystallised from diisopropyl ether, to give 2-hydroxy-4-(3-thienylmethoxy)benzaldehyde (8 g), in
30 the form of a white solid.

A mixture of 2-hydroxy-4-(3-thienylmethoxy)benzaldehyde (7.7 g), nitroethane (4.74 mL), sodium acetate (5.4 g) and glacial acetic acid (6.6 mL) is heated at reflux for 14 hours. The mixture is then cooled, diluted with water
35 (50 mL) and extracted with ethyl acetate (3 x 50 mL). The combined organic extracts are washed with saturated aqueous sodium bicarbonate solution

(3 x 50 mL), dried over magnesium sulphate, and concentrated in vacuo. The resulting residue is subjected to flash chromatography on silica gel, eluting with a mixture of ethyl acetate and pentane (3:7 v/v), to give 2-hydroxy-4-(3-thienylmethoxy)benzonitrile (7.6 g), in the form of a pale yellow solid.

5

REFERENCE EXAMPLE 47

A mixture of methyl (RS)-5-(3-iodophenyl)-4-(2-methylphenyl)-5-oxopentanoate (1.7 g), bis(triphenylphosphine)palladium (II) chloride (0.1 g),
10 tetrakis(triphenylphosphine)palladium (0) (0.13 g) and benzylthiotrimethylstannane (1.17 g) in toluene (37 mL) is heated at reflux for 24 hours. The reaction mixture is washed twice with 10% aqueous potassium fluoride (20 mL), with water (20 mL), dried over magnesium sulphate and evaporated. The residue (2.87 g) is purified by flash chromatography on silica eluting with a
15 mixture of pentane and ethyl acetate (98:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-5-(3-benzylthiophenyl)-4-(2-methylphenyl)-5-oxopentanoate as an orange oil (0.57 g).

A solution of 1-(3-iodophenyl)-2-(2-methylphenyl)ethanone (3.21 g) in
20 dry tetrahydrofuran (30 mL) is treated with methyl acrylate (0.9 mL) and potassium tertiary butoxide (0.107 g). The reaction mixture is heated at reflux for 18 hours, evaporated and the residue partitioned between ethyl acetate (60 mL) and water (30 mL). The organic phase is dried over magnesium
25 sulphate, evaporated and the residual solid washed with pentane to give methyl (RS)-5-(3-iodophenyl)-4-(2-methylphenyl)-5-oxopentanoate as a fawn coloured solid (3.39 g), m.p. 106-108°C.

A solution of 2-methylbenzyl chloride (4.57 mL) in ether (61 mL) is
30 added dropwise to magnesium (1.64 g) at a rate to maintain reflux. The mixture is stirred for 15 minutes, decanted from the excess magnesium and this solution is added dropwise to a stirred solution of N-methoxy-N-methyl-3-iodobenzamide (4.9 g) in ether (30 mL) at 5-10°C giving a cream coloured precipitate. The suspension is treated with 2 N hydrochloric acid (50 mL) and
35 the organic phase separated, washed with water (50 mL) and dried over magnesium sulphate. 1-(3-iodophenyl)-2-(2-methylphenyl)ethanone (0.69 g) is precipitated in the filtrate during filtration. Evaporation of the solution gives

an oily solid which is triturated with a mixture of pentane and ethyl acetate (9:1 v/v) to give a further quantity of 1-(3-iodophenyl)-2-(2-methylphenyl)ethanone (2.69 g), m.p.115-117°C. [Elemental analysis:- C,53.9; H,3.90%. Calculated :- C,53.6; H,3.90%].

5

A stirred solution of N-methoxymethylamine (3.47 g) and triethylamine (10.4 mL) in dichloromethane (100 mL) at 5°C is treated dropwise with 3-iodobenzoyl chloride (9.8 g). The reaction mixture is stirred at room temperature for 2 hours, washed twice with water (70 mL), dried over
10 magnesium sulphate, and evaporated to give N-methoxy-N-methyl-3-iodobenzamide (9.8 g) as a brown oil.

A stirred suspension of 3-iodobenzoic acid (8.84 g) in thionyl chloride (44 mL) is heated at reflux for 30 minutes. The dark brown solution is
15 evaporated to give 3-iodobenzoyl chloride (9.8 g) as a brown oil.

REFERENCE EXAMPLE 48

20 A stirred solution of ethyl (RS)-4-(2-acetyl-5-(3-thienyl-methoxy)phenoxy)-4-(2-methylphenyl)butanoate (0.5 g) in pyridine (2.5 mL) is treated with selenium dioxide (0.204 g) and the mixture is heated at 100°C for 4 hours. A further quantity of selenium dioxide (0.1 g) is added and the mixture is heated at 100°C for 18 hours. The suspension is filtered and the filtrate
25 evaporated to give an orange oil (0.76 g) which is dissolved in ethyl acetate (20 mL) and washed twice with 1 N acetic acid (10 mL), with water (10 mL) and dried over magnesium sulphate. Evaporation gives an orange oil (0.58 g) which is purified by flash chromatography on silica eluting with a mixture of dichloromethane and methanol (9:1 v/v). Fractions homogeneous in the
30 required product are combined and evaporated to give ethyl (RS)-4-(2-carboxycarbonyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoate (0.38 g) as a yellow gum.

REFERENCE EXAMPLE 49

35

Diisopropyl azodicarboxylate (6.45 mL) is added dropwise to a stirred, cooled (0°C) solution of triphenylphosphine (8.6 g) in dry tetrahydrofuran

(60 mL). After a further 30 minutes, a solution of 2'-hydroxy-4'-(pyridin-3-ylmethoxy)benzophenone (5 g) and tert-butyl 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (S)-enantiomer, (5.13 g) in dry tetrahydrofuran (60 mL) is added dropwise over 25 minutes and stirring continued for a further
5 3 hours at °C. The reaction is allowed to warm to ambient temperature overnight before being concentrated under vacuo. The resultant oil is purified by flash chromatography on silica eluting with a mixture of dichloromethane and ethyl acetate (9:1 v/v). Fractions homogeneous in the required product are combined and concentrated under reduced pressure to give *t*-butyl
10 4-[2-benzoyl-5-(pyridin-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate, thought to be the (R)-enantiomer, as a colourless oil (8.66 g).

A mixture of 2,4-dihydroxybenzophenone (21.4g), 3-chloromethylpyridine hydrochloride (25 g), potassium carbonate (80 g), potassium iodide
15 (0.5 g) and tetrabutylammonium bromide (0.2 g) in methyl ethyl ketone is heated at reflux with mechanical stirring for 18 hours. The reaction mixture is filtered through hyflo and the solids washed with several portions of methyl ethyl ketone. The filtrate and the washings are combined and concentrated under reduced pressure to leave a brown oil which is dissolved in ethyl acetate
20 (250 mL) and washed four times with water (200 mL). The organic layer is dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a residue which is crystallised from cyclohexane containing a little decolourizing charcoal to give 2'-hydroxy-4'-(pyridin-3-ylmethoxy)-benzophenone (21.2 g) as a pale yellow solid, m.p. 88-90 °C.

25

REFERENCE EXAMPLE 50

Dry sodium 4-hydroxy-4-(2-methylphenyl)butanoate, thought to be the (R)-enantiomer, (1 g) is added portionwise to a stirred suspension of sodium
30 hydride (0.6 g, 60% dispersion in mineral oil) in dry tetrahydrofuran (50 mL) under nitrogen. The reaction mixture is stirred at 50°C for 15 minutes when methyl 3-(3-fluoro-4-nitrophenoxy)methyl]benzoate (1.4 g) is added in one portion. After stirring at 50-60°C for 18 hours the reaction mixture is evaporated to dryness and the residue partitioned between ethyl acetate (50
35 mL) and 1 N hydrochloric acid (50 mL). The organic phase is washed three times with water (50 mL), dried over magnesium sulphate and evaporated to give a yellow-brown oil (2.47 g) which is purified by flash chromatography on

silica eluting initially with a mixture of cyclohexane and ethyl acetate (4:1 v/v) then eluting with a mixture of cyclohexane and ethyl acetate (3:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give methyl 3-[3-(3-carboxy-1-(2-methylphenyl)propoxy)-4-nitrophenoxyethyl]-benzoate (0.86 g) as a clear yellow oil.

2-Fluoro-4-hydroxynitrobenzene (3 g) is added portionwise to a stirred suspension of sodium hydride (0.9 g, 60% dispersion in mineral oil) in dry dimethylformamide (50 mL) under nitrogen. After stirring for 20 minutes methyl 3-bromomethylbenzoate is added in one portion and stirring continued for 3 hours. The reaction mixture is evaporated to dryness and the residue partitioned between ethyl acetate (50 mL) and water (50 mL). The organic phase is washed three times with water (50 mL), dried over magnesium sulphate and evaporated to give a yellow solid which is crystallised from a mixture of ethyl acetate and cyclohexane (1:3 v/v) to give methyl 3-(3-fluoro-4-nitrophenoxyethyl)benzoate (4.95 g) as a white solid.

REFERENCE EXAMPLE 51

Diisopropyl azodicarboxylate (2.6 g) is added portionwise to a stirred solution of triphenylphosphine (3.4 g) in dry tetrahydrofuran (40 mL) at 0°C under nitrogen. The reaction mixture is stirred at 0°C for 15 minutes then treated with a solution of 2-fluoro-4-hydroxybenzonitrile (0.87 g) and *t*-butyl 3-(3-hydroxymethylphenyl)propionate (1.5 g) in dry tetrahydrofuran (10 mL) whilst maintaining the temperature at 0-5°C. Stirring is continued at this temperature for 1 hour when the reaction mixture is allowed to warm up to room temperature. The reaction mixture is evaporated to low bulk and partitioned between ethyl acetate (100 mL) and water (100 mL). The organic phase is washed with water (50 mL), dried over magnesium sulphate and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of cyclohexane and ethyl acetate (4:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give *tert*-butyl 3-[3-(4-cyano-3-fluoro-phenoxyethyl)phenyl]propionate (0.3 g).

A stirred solution of *tert*-butyl 3-(3-formylphenyl)acrylate (5 g) in methanol (150 mL) at 0°C is treated with sodium borohydride (0.38 g) portionwise. After stirring at 0°C for 20 minutes the reaction mixture is

partitioned between ethyl acetate (200 mL) and water (100 mL). The organic phase is separated, dried over magnesium sulphate and evaporated to give tert-butyl 3-(3-hydroxymethylphenyl)acrylate which is dissolved in dry toluene (50 mL) and treated with tris(triphenylphosphine)rhodium(I) chloride (80 mg) and triethylsilane (8 mL). The mixture is stirred at room temperature for 3 hours, left standing at room temperature for 18 hours and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of cyclohexane and ethyl acetate (4:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give tert-butyl 3-(3-hydroxymethylphenyl)propionate (1.5 g).

To a stirred mixture of 3-bromobenzaldehyde (20 g), tert butyl acrylate (40 mL), triethylamine (44 mL) and tris(2-methylphenyl)phosphine (3.4 g) in dry dimethylformamide (300 mL) is added palladium (II) acetate in one portion. The resulting dark brown mixture is heated at 100°C for 18 hours then evaporated in vacuo. The residual semi-solid is partitioned between ethyl acetate (250 mL) and 2 N hydrochloric acid (250 mL). the organic phase is washed three times with water (250 mL), dried over magnesium sulphate and evaporated. The resulting dark brown oil (28.1 g) is purified by flash chromatography on silica eluting initially with a mixture of cyclohexane and dichloromethane (4:1 v/v) then with a mixture of cyclohexane and dichloromethane (1:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give tert-butyl 3-(3-formylphenyl)acrylate as a yellow oil (14.08 g).

REFERENCE EXAMPLE 52

Burgess reagent (1.3 g) is added to a solution of ethyl (RS)-4-[2-(2-hydroxyethylcarbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (2.5 g) in THF (50 mL) under nitrogen and the mixture refluxed for 30 minutes. The reaction is cooled to ambient temperature then partitioned between ethyl acetate (100 mL) and water (100 mL). The organic phase is dried over magnesium sulphate, filtered and evaporated to dryness. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and cyclohexane (1:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-[2-(4,5-

dihydrooxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate (1.5 g) as a white solid.

- 2-Hydroxy-N-(2-hydroxyethyl)-4-(3-thienylmethoxy)benzamide (2.93 g)
5 is added portionwise to a stirred suspension of sodium hydride (0.44 g of a 60% dispersion in mineral oil) in DMF (100 mL) at ambient temperature. The reaction is stirred at ambient temperature for 30 minutes then a solution of ethyl (R,S)-4-chloro-4-(2-methylphenyl) butanoate (3.6 g) in DMF (5 mL) is added in one portion and the reaction stirred for 4 hours at 100°C. The reaction is left at
10 ambient temperature for 16 hours then concentrated in vacuo. The residue is partitioned between ethyl acetate (150 mL) and water (100 mL). The organic phase is washed with water (100 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and cyclohexane (1:1 v/v).
15 Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-(2-(N-2-hydroxyethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy)-4-(2-methylphenyl) butanoate (3.9 g) as a yellow oil.

- A mixture of methyl 2-hydroxy-5-(3-thienylmethoxy)benzoate (9.96 g)
20 and ethanolamine (250 mL) is refluxed for 1 hour. The ethanolamine is neutralised with 2N HCl and the mixture extracted with ethyl acetate (2 x 250 mL). The combined organic extracts are washed with water (2 x 100 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure to give a yellow solid. Trituration with diethyl ether gives 2-Hydroxy-N-(2-
25 hydroxyethyl)-4-(3-thienylmethoxy)benzamide (6.8 g) in the form of a white solid.

REFERENCE EXAMPLE 53

- 30 A solution of ethyl (RS)-4-[2-(4,5-dihydrooxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (900 mg) and nickel peroxide (5.04 g) in toluene (50 mL) is stirred at 100°C for 2 hours. The reaction is left at ambient temperature for 16 hours then filtered through a pad of hyflo and the filtrate concentrated in vacuo. The residue is purified by flash chromatography
35 on silica eluting with a mixture of ethyl acetate and cyclohexane (15:85 v/v). Fractions homogeneous in the required product are combined and evaporated

to give ethyl (RS)-4-[2-(oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (140 mg) and starting material (400 mg).

REFERENCE EXAMPLE 54

5

Methyl (RS)-2-[2-hydroxy-4-(3-thienylmethoxy)phenyl]-4,5-dihydro-oxazole-4-carboxylate (200 mg) is added portionwise to a stirred suspension of sodium hydride (0.04 g, 60% dispersion in mineral oil) in DMF (25 mL) at ambient temperature. The reaction is stirred at ambient temperature for
10 30 minutes then a solution of ethyl (R,S)-4-chloro-4-(2-methylphenyl)butanoate (240 mg) in DMF (5 mL) is added in one portion and the reaction stirred for 4 hours at 100°C. The reaction is left at ambient temperature for 16 hours then concentrated in vacuo. The residue is partitioned between ethyl acetate (50 mL) and water (50 mL). The organic phase is washed with water
15 (50 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and cyclohexane (15:85 v/v). Fractions homogeneous in the required product are combined and evaporated to give methyl (RS)-2-[2-(3-ethoxycarbonyl-1-(2-methylphenyl)propoxy)-4-(3-thienylmethoxy)benzoyl-
20 amino]acrylate (250 mg) as a colourless oil.

Thionyl chloride (7.14 g) is added to a stirred solution of methyl (RS)-3-hydroxy-2-[4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxymethoxy)benzoyl]-propionate (6.5 g) in dichloromethane (100 mL) at ambient temperature. The
25 reaction is left to stand at room temperature for 18 hours. The reaction mixture is partitioned between ethyl acetate (250 mL) and 55% w/v potassium carbonate. The organic phase is washed with water (200 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is recrystallised from ethyl acetate/cyclohexane to give methyl (RS)-2-[2-hydroxy-
30 4-(3-thienylmethoxy)phenyl]-4,5-dihydrooxazole-4-carboxylate (1.9 g) in the form of an off white solid.

A mixture of 4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxymethoxy)-benzoic acid (6 g), (RS) serine methyl ester hydrochloride (2.7 g), 1-hydroxy-benzotriazole (2.6 g), triethylamine (2.4 mL), 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (3.6 g) and DMF (100 mL) is stirred at ambient
35 temperature for 16 hours. The reaction is concentrated in vacuo and the

residue partitioned between ethyl acetate (150 mL) and 1N HCl (100 mL). The organic phase is washed with water (2 x 100 mL) then dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting initially with a mixture of ethyl acetate and cyclohexane (3:7 v/v) then with a mixture of ethyl acetate and cyclohexane (1:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give methyl (RS)-3-hydroxy-2-[4-(3-thienylmethoxy)-2-(2-trimethylsilylethoxymethoxy)benzoyl]aminopropionate (6.5 g) as a colourless oil.

REFERENCE EXAMPLE 55

Methyl 2-[2-hydroxy-4-(3-thienylmethoxy)phenyl]oxazole-4-carboxylate (400 mg) is added portionwise to a stirred suspension of sodium hydride (100 mg, 60% dispersion in mineral oil) in dry DMF (30 mL). After stirring at 60°C for 15 minutes, ethyl (R,S)-4-chloro-4-(2-methylphenyl) butanoate (0.6 g) is added in one portion and the reaction stirred for 2 hours at 90°C. The reaction is concentrated in vacuo and the residue partitioned between ethyl acetate (100 mL) and water (100 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:4 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-[2-(4-methoxycarbonyloxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (300 mg) as an yellow oil.

A mixture of methyl 2-[2-hydroxy-4-(3-thienylmethoxy)phenyl]-4,5-dihydrooxazole-4-carboxylate (1g) and (2,3-dichloro-5,6-dicyano-benzoquinone (0.8 g) in toluene (100 mL) is stirred at reflux for 2 hours. The reaction is concentrated in vacuo and the residue passed through a pad of silica eluting with 20% ethyl acetate in cyclohexane. The eluant is concentrated in vacuo and the residue partitioned between ethyl acetate (100 mL) and 1N HCl (50 mL). The organic phase is washed with water (20 mL), dried over magnesium sulphate and concentrated in vacuo to give methyl 2-[2-hydroxy-4-(3-thienylmethoxy)phenyl]oxazole-4-carboxylate (400 mg).

REFERENCE EXAMPLE 56

2-Hydroxy-4-(3-thienylmethoxy)benzonitrile (0.5 g) is added portionwise to a stirred suspension of sodium hydride (90 mg, 60% dispersion in mineral oil) in dry DMF (100 mL). After stirring at ambient temperature for 30 minutes, ethyl (RS)-4-chloro-4-(2-chloro-6-fluorophenyl)butanoate (0.62 g) is added in one portion and the reaction stirred for 48 hours at 90°C. The reaction mixture is concentrated in vacuo and the residue partitioned between ethyl acetate (100 mL) and water (100 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-(2-chloro-6-fluorophenyl)-4-(2-cyano-5-(3-thienylmethoxy)phenoxy)butanoate (360 mg) as an orange oil.

A solution of 2-chloro-6-fluorobenzaldehyde (15 g) in dichloromethane (20 mL) is added dropwise to a stirred 1M solution of titanium (IV) chloride in dichloromethane (95 mL) at 0°C under nitrogen. The mixture is stirred for 5 minutes and [(1-ethoxycyclopropyloxy)trimethyl]silane (21.1 mL) is added dropwise maintaining the temperature at 0-5°C. The mixture is stirred for 16 hours at ambient temperature then quenched with water (200 mL). The mixture is extracted three times with dichloromethane (100 mL) and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-chloro-4-(2-chloro-6-fluorophenyl)butanoate (24.3 g) as a yellow oil.

REFERENCE EXAMPLE 57

3-(3-Fluoro-4-nitrophenoxy)methylpyridine (1.53 g) is added to a stirred suspension of sodium hydride (0.48 g, 60% dispersion in mineral oil) in THF (50 mL) at ambient temperature. A solution of tert-butyl (R)-4-hydroxy-4-(2-methylphenyl)butanoate (3.08 g) in THF (30 mL) is added dropwise to the mixture over 2 hours. The reaction mixture is concentrated under reduced pressure and the residue partitioned between water (100 mL) and ethyl acetate (100 mL). The aqueous layer is extracted three times with ethyl acetate (100 mL) and the combined organic phases were dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash

chromatography on silica eluting with ethyl acetate. Fractions homogeneous in the required product are combined and evaporated to give tert-butyl (R)-4-(2-methylphenyl)-4-(2-nitro-5-(3-pyridylmethoxy)phenoxy)butanoate (0.85 g) as a yellow gum.

5

3-Fluoro-4-nitrophenol (7.5 g) is added to a stirred suspension of sodium hydride (2.2 g, 60% dispersion in mineral oil) in DMF (50 mL). The mixture is stirred at ambient temperature for 30 minutes. To this mixture is added a solution prepared by the addition of sodium hydride (2.2 g, 60% dispersion in mineral oil) to a stirred suspension of 3-chloromethylpyridine hydrochloride (8.2 g) in DMF (50 mL). The resulting mixture is stirred at 90°C for 16 hours then cooled to ambient temperature. The reaction mixture is concentrated in vacuo and the residue partitioned between dichloromethane (100 mL) and water (100 mL). The aqueous phase is extracted four times with dichloromethane (100 mL). The combined organic phases are washed with brine (100 mL), dried over magnesium sulphate, filtered and concentrated in vacuo to give an orange solid. Recrystallisation from ethyl acetate gives 3-(3-fluoro-4-nitrophenoxy)methylpyridine (4.6 g) as pale orange/brown crystals.

15
20

REFERENCE EXAMPLE 58

2-Hydroxy-4-(3-thienylmethoxy)benzonitrile (1.5 g) is added portionwise to a stirred suspension of sodium hydride (440 mg, 60% dispersion in mineral oil) in dry DMF (50 mL). After stirring at ambient temperature for 30 minutes, ethyl (RS)-4-chloro-4-(2,5-dimethylphenyl)butanoate (2.5 g) is added in one portion and the reaction stirred for 48 hours at 90°C. The reaction is concentrated in vacuo and the residue partitioned between ethyl acetate (50 mL) and water (50 mL). The aqueous layer is extracted twice with ethyl acetate (50 mL). The combined organic phases are dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-(2-cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoate (500 mg) as a pale brown oil.

30
35

A solution of 2,5-dimethylbenzaldehyde (10 g) in dichloromethane

(20 mL) is added dropwise to a stirred 1M solution of titanium (IV) chloride in dichloromethane (75 mL) at 0°C under nitrogen. The mixture is stirred for 5 minutes and [(1-ethoxycyclopropyloxy)trimethyl]silane (16.6 mL) is added dropwise maintaining the temperature at 0-5°C. The mixture is stirred for 5 16 hours at ambient temperature then quenched with water (200 mL). The mixture is extracted three times with dichloromethane (100 mL) and the combined organic extracts are washed with brine (50 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-chloro-4-(2,5-dimethylphenyl) butanoate (17.3 g) as a yellow oil.

10

REFERENCE EXAMPLE 59

A solution of ethyl (RS)-4-(benzo[1,3]dioxol-4-yl)-4-hydroxybutanoate (2.52 g) in ethanol (56 mL) is treated with 1 N sodium hydroxide solution 15 (10 mL). The reaction mixture is stirred at room temperature for 4 hours, evaporated and the residue azeotroped with toluene to give sodium (RS)-(benzo[1,3]dioxol-4-yl)-4-hydroxybutanoate (2.22 g) as a brown gum.

20 A mixture of 2,3-methylenedioxybenzaldehyde (1.5 g) and [1-ethoxy-cyclopropyl]oxy]trimethylsilane (2.7 g) in dry dichloromethane (15 mL) is added dropwise to a stirred suspension of zinc iodide (10 mg) in dry dichloromethane (20 mL) under nitrogen. The temperature of the reaction mixture reaches 25-35°C. The reaction flask is wrapped in aluminium foil and stirring is continued 25 for 3 hours. The reaction mixture is treated with dry pyridine (20 drops) and stirring continued for 15 minutes. The reaction mixture is evaporated, washed twice with water (50 mL), twice with brine (50 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-(benzo[1,3]dioxol-4-yl)-4-hydroxybutanoate (6.01 g) as a yellow oil.

30

REFERENCE EXAMPLE 60

A solution of ethyl (RS)-4-(2,3-dimethylphenyl)-4-hydroxybutanoate (3.66 g) in ethanol (100 mL) is treated with 1 N sodium hydroxide solution 35 (15.5 mL). The reaction mixture is stirred at room temperature for 18 hours, evaporated and the residue azeotroped with toluene to give sodium (RS)-4-(2,3-dimethylphenyl)-4-hydroxy-butanoate (3.2 g) as a brown solid.

A mixture of 2,3-dimethylbenzaldehyde (2 g) and [1-ethoxycyclopropyl]-oxy]trimethylsilane (3.6 mL) in dry dichloromethane (20 mL) is added dropwise to a stirred suspension of zinc iodide (<100 mg) in dry dichloromethane (25 mL) under nitrogen. The reaction flask is wrapped in aluminium foil and stirring is continued for 3 hours. The reaction mixture is treated with dry pyridine (10 mL) and stirring continued for 15 minutes. The reaction mixture is washed three times with water (50 mL), twice with brine (50 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-(2,3-dimethyl-phenyl)-4-hydroxybutanoate (3.6 g) as a yellow oil.

REFERENCE EXAMPLE 61

Sodium (RS)-4-hydroxy-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy-phenyl)butanoate (1.81 g) is added to a stirred suspension of sodium hydride (0.6 g, 60% dispersion in mineral oil) in tetrahydrofuran (50 mL). The mixture is stirred at room temperature for 5 minutes then at 50°C for 1.5 hours when 2-fluoro-4-(3-pyridylmethoxy)benzonitrile (1.07 g) is added in one portion. The reaction mixture is heated at reflux for 18 hours and evaporated. The residue is dissolved in water, the solution acidified to pH 1 with 1 N hydrochloric acid and the mixture extracted with ethyl acetate (200 mL). The organic phase is washed with water (200 mL), brine (200 mL), dried over magnesium sulphate and evaporated to give (RS)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy)phenyl)butanoic acid (1.86 g) as a cream solid.

A solution of ethyl (RS)-4-hydroxy-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxyphenyl)butanoate (3.68 g) in ethanol (50 mL) is treated with 1 N sodium hydroxide solution (10 mL). The reaction mixture is stirred at room temperature for 4.5 hours, evaporated and the residue azeotroped with toluene to give sodium (RS)-4-hydroxy-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy-phenyl)butanoate (3.24 g) as a yellow gum.

A mixture of 2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy)benzaldehyde (3.44 g) and [1-ethoxycyclopropyl]oxy]trimethylsilane (2.7 g) in dry dichloromethane (15 mL) is added dropwise to a stirred suspension of zinc iodide

- (10 mg) in dry dichloromethane (25 mL) under nitrogen. The temperature of the reaction mixture reaches 25-35°C. The reaction flask is wrapped in aluminum foil and stirring is continued for 3 hours. The reaction mixture is treated with dry pyridine (20 drops) and stirring continued for 15 minutes. The reaction mixture is evaporated, washed twice with water (100 mL), brine (100 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-hydroxy-4-(2-methyl-5-(2-(trimethylsilyl)ethoxymethoxyphenyl)butanoate (8.25 g) as a light-brown oil.
- 5
- 10 5-hydroxy-2-methylbenzaldehyde (3.4 g) is added portionwise to a stirred suspension of sodium hydride (1.2 g, 60% dispersion in mineral oil). The reaction mixture is stirred at room temperature for 45 minutes, 2-(trimethylsilyl)ethoxymethyl chloride (5 g) is added slowly and stirring continued for a further 3 hours. The mixture is evaporated and the residue partitioned between ethyl acetate (50 mL) and water (150 mL). The organic phase is washed with brine (150 mL) dried over magnesium sulphate and evaporated to give 2-methyl-5-(2-(trimethylsilyl)ethoxymethoxy)benzaldehyde (6.87 g) as a red-brown oil.
- 15
- 20 A stirred mixture of 5-methoxy-2-methylbenzaldehyde (24.9 g) and pyridine hydrochloride (100 g) is heated at 170°C for 7 hours. Water (250 mL) is added cautiously to the reaction mixture at 140°C and the very dark mixture is extracted twice with chloroform (250 mL), filtration through hyflo removes a small amount of insoluble black solid and aids separation. The organic phase is washed with 1 N hydrochloric acid (250 mL), with water (250 mL), with brine (250 mL), dried over magnesium sulphate and evaporated. The residue is washed with pentane to give 5-hydroxy-2-methylbenzaldehyde (10.66 g) as a cream solid, m.p. 116-118°C.
- 25
- 30 A solution of 2-(5-methoxy-2-methylphenyl)-[1,3]dioxolane (18.25 g) in tetrahydrofuran (180 mL) is treated with 1 N hydrochloric acid (27 mL). The reaction mixture is heated at reflux for 2 hours, evaporated and the residue is partitioned between ethyl acetate (150 mL) and water (150 mL). The organic phase is washed with water (150 mL), with brine (150 mL), dried over magnesium sulphate and evaporated to give 5-methoxy-2-methylbenzaldehyde (12 g) as an orange oil.
- 35

A stirred solution of 2-(2-bromo-5-methoxyphenyl)-[1,3]dioxolane (25.92 g) in tetrahydrofuran (400 mL) at -70°C is treated dropwise with a solution of *n*-butyl lithium in hexanes (48.0 mL, 2.5 M). After stirring for 10 minutes methyl iodide (21.29 g) is added dropwise whilst maintaining the temperature at -65°C. The reaction mixture is stirred at -65°C for 1 hour, allowed to warm to room temperature and stirring is continued for a further hour. The reaction mixture is evaporated and the residue partitioned between ethyl acetate (50 mL) and water (50 mL). The organic phase is washed with brine (500 mL), dried over magnesium sulphate and evaporated to give 2-(5-methoxy-2-methyl-phenyl)-[1,3]dioxolane (18.25 g) as a yellow oil.

A mixture of 2-bromo-5-methoxy-benzaldehyde (45.25 g), ethylene glycol (52.22 g) and 4-toluenesulphonic acid (0.4 g) in toluene (400 mL) is heated at reflux for 5 hours using a Dean and Stark apparatus to remove water formed in the reaction. The reaction mixture is washed five times with water (500 mL) and evaporated giving 2-(2-bromo-5-methoxyphenyl)-[1,3]dioxolane (51.9 g) as a yellow liquid.

A stirred solution of 3-methoxybenzaldehyde (27.23 g) in dichloromethane (150 mL) at 0°C is treated with a solution of bromine (32 g) in dry dichloromethane (50 mL) over 30 minutes. The reaction mixture is stirred at room temperature for 4 hours, combined with a previous preparation carried out on the same scale, and washed with saturated sodium metabisulphite solution (400 mL), with water (400 mL), brine (400 mL) and dried over magnesium sulphate. Evaporation and recrystallisation of the residue from petroleum ether (b.p. 40-60°C) gives 2-bromo-5-methoxybenzaldehyde (45.25 g) as a white solid, m.p. 73-74°C.

REFERENCE EXAMPLE 62

30

A suspension of 6-(2-methylphenyl)tetrahydropyran-2-one (2.5 g) in a mixture of tetrahydrofuran (20 mL) and methanol (20 mL) is treated with 5 M sodium hydroxide solution under nitrogen. The reaction mixture is stirred at room temperature for 24 hours, diluted with water (100 mL) and extracted with ethyl acetate (100 mL). The aqueous phase is washed three times with ethyl acetate (50 mL) and evaporated in vacuo. The residue is azeotroped with

toluene to give sodium (RS)-5-hydroxy-4-(2-methylphenyl)pentanoate (1.57 g) as a white solid.

To a stirred suspension of 5-oxo-5-(2-methylphenyl)pentanoic acid (10 g) in water (100 mL) is added potassium hydroxide pellets (6.35 g). The temperature of the mixture is maintained at 2-20°C whilst sodium borohydride (1.47 g) is added over 15 minutes. The reaction mixture is left at room temperature for 18 hours when 1 N hydrochloric acid (100 mL) is cautiously added followed by water (50 mL) and ethyl acetate (100 mL). The organic phase is separated and the aqueous phase extracted twice with ethyl acetate (100 mL). The combined organic phases are washed with brine (100 mL), dried over magnesium sulphate and evaporated to give 6-(2-methylphenyl)-tetrahydropyran-2-one (5.27 g) as a yellow oil.

A stirred solution of glutaric anhydride (22.82 g) in dry tetrahydrofuran (300 mL) at -78°C is treated with a solution of 2-methylphenyl magnesium chloride in tetrahydrofuran (200 mL, 0.1 M) over 45 minutes. The mixture is stirred at -78°C for 2.5 hours and the grey suspension is poured into 1 N hydrochloric acid (300 mL). The mixture is extracted with ethyl acetate (300 mL). The organic phase is dried over magnesium sulphate and evaporated to give 5-oxo-5-(2-methylphenyl)pentanoic acid (36.74 g) as a red-brown oil.

REFERENCE EXAMPLE 63

2-Fluoro-6-hydroxy-4-(3-thienylmethoxy)benzonitrile (200 mg) is added to a stirred suspension of sodium hydride (36 mg, 60% dispersion in mineral oil) in dry DMF (10 mL). After stirring at ambient temperature for 30 minutes, a solution of ethyl (R,S)-4-chloro-4-(2-methylphenyl)butanoate (220 mg) in DMF (5 mL) is added in one portion and the reaction stirred for 3 hours at ambient temperature. The reaction mixture is concentrated in vacuo and the residue partitioned between diethyl ether (20 mL) and water (20 mL). The aqueous layer is extracted four times with diethyl ether (20 mL). The combined organic phases are washed with brine (20 mL), dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane

(1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-(2-cyano-3-fluoro-5-(3-thienylmethoxy)-phenoxy)-4-(2-methylphenyl) butanoate (200 mg) as a pale yellow oil.

- 5 A mixture of 2-fluoro-6-hydroxy-4-(3-thienylmethoxy)benzaldehyde (1 g), nitroethane (0.6 g) sodium acetate (0.66 g) and glacial acetic acid (1 mL) is heated at reflux for 2 hours. The reaction mixture is cooled to ambient temperature, water (50 mL) is added and the mixture is extracted three times with ethyl acetate (50 mL). The combined organic phases are washed with
10 saturated brine (50 mL), dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a brown oil. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give 2-fluoro-6-hydroxy-4-(3-thienylmethoxy)-
15 benzonitrile (250 mg), as a yellow solid.

- Sodium hydride (0.52 g, 60% dispersion in mineral oil) is added portionwise over 30 minutes to a solution of 2,4-dihydroxy-6-fluoro-benzaldehyde (2 g) in DMF (50 mL) at 0°C. The mixture is stirred at 0°C for
20 30 minutes then a solution of 3-chloromethylthiophene (1.72 g) in DMF (50 mL) is added and the mixture stirred at 60°C for 16 hours. The reaction is concentrated in vacuo and the residue partitioned between diethyl ether (50 mL) and water (50 mL). The aqueous phase is extracted four times with diethyl ether (50 mL) and the combined organic phases washed with brine
25 (50 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:4 v/v). Fractions homogeneous in the required product are combined and evaporated to give a pale yellow solid which is triturated with diisopropyl ether to give 6-fluoro-2-
30 hydroxy-4-(3-thienylmethoxy)benzaldehyde (1.1 g) as a yellow solid.

- Dimethylformamide (DMF) (12.7 g) is added to vigorously stirred phosphorous oxychloride (14.4 g) at 0°C. The reaction mixture is stirred at this temperature for 30 minutes, then 3,5-dihydroxyfluorobenzene (6 g) is added.
35 The sticky red syrup is allowed to warm to room temperature and stirred for 2 hours, then left to stand for 14 hours. Water (100 mL) is added and the mixture extracted three times with ethyl acetate (100 mL). The combined organic

phases are washed with brine (100 mL), dried over magnesium sulphate and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give an orange solid
5 which is triturated with ethyl acetate to give 2,4-dihydroxy-6-fluorobenzaldehyde (2 g) as a yellow solid.

REFERENCE EXAMPLE 64

10 To a stirred suspension of sodium hydride (0.24 g of a 60% w/w dispersion in mineral oil) in toluene (30 mL) is added a solution of (RS)-2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)acetophenone (1.42 g) and diethyl oxalate (2.45 g) in toluene (20 mL) dropwise. The reaction is heated at reflux for two hours and concentrated under reduced pressure to
15 leave ethyl (RS)-2,4-dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)phenyl]butanoate as a brown oil.

To a stirred, cooled (0°C) solution of triphenylphosphine (8.4 g) in dry tetrahydrofuran (60 mL) is added diisopropyl azodicarboxylate (6.5 mL)
20 dropwise. When a cream suspension is formed, a solution of 2-hydroxy-4-(3-thienylmethoxy)acetophenone (3.5 g) and (RS)-1-(2-methylphenyl)ethanol (2.2 g) in tetrahydrofuran (75 mL) is added dropwise and the reaction stirred at ambient temperature for 48 hours. The reaction is partitioned between ethyl acetate and 1 N aqueous sodium hydroxide. The organic layer is washed with
25 water, dried over magnesium sulphate and concentrated under reduced pressure. The residue is chromatographed on silica gel eluting with dichloromethane to give (RS)-2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)acetophenone (2.61 g) as a colourless oil.

30 REFERENCE EXAMPLE 65

A solution of tert-butyl (R)-4-(2-acetyl-5-(3-pyridylmethoxy)phenoxy)-4-(2-methylphenyl)butanoate (1.46 g) and diethyl oxalate (1.46 g) in toluene (20 mL) is added dropwise to a stirred suspension of sodium hydride (0.15 g, 60%
35 dispersion in mineral oil) in toluene (50 mL) at ambient temperature. The mixture is stirred at reflux for three hours, cooled to ambient temperature and concentrated in vacuo. The residue is taken up in ethanol (100 mL), treated

with hydrazine hydrate (0.16 mL) and acidified to pH 5 by the addition of glacial acetic acid. The mixture is refluxed for 3 hours, cooled to ambient temperature then concentrated in vacuo. The residue is partitioned between ethyl acetate (100 mL) and water (100 mL) and the organic phase washed with water (100 mL), dried over magnesium sulphate and passed through a pad of silica
5 eluting with ethyl acetate. The fractions containing product are concentrated in vacuo to give t-butyl (R)-4-[2-(5-ethoxycarbonylpyrazol-3-yl)-5-(3-pyridyl-methoxy)phenoxy]-4-(2-methylphenyl)butanoate (1.7 g).

10 Diisopropyl azodicarboxylate (8 g) is added portionwise to a stirred solution of triphenylphosphine (10.5 g) in THF (100 mL) at 0°C under nitrogen. Stirring is continued at 0°C for 15 minutes. A solution of 2-acetyl-5-(3-pyridyl-methoxy)phenol (4.5 g) and tert-butyl (S)-4-hydroxy-4-(2-methylphenyl)-
15 butanoate (4.9 g) in THF (100 mL) is added at such a rate that the temperature does not exceed 5°C. The reaction is allowed to warm to ambient temperature stirred for 16 hours, then concentrated in vacuo. The residue is partitioned between ethyl acetate (200 mL) and water (200 mL) and the organic phase dried over magnesium sulphate, filtered and concentrated in vacuo. The
20 residue is purified by flash chromatography on silica eluting initially with a mixture of ethyl acetate and cyclohexane (1:1 v/v) then with ethyl acetate. Fractions homogeneous in the required product are combined and evaporated to give t-butyl (R)-4-(2-acetyl-5-(3-pyridylmethoxy)phenoxy)-4-(2-methyl-
phenyl)butanoate (2 g).

25 REFERENCE EXAMPLE 66

A solution of ethyl bromodifluoroacetate (5.6 g) and 2-methyl-benzaldehyde (3 g) in THF (25 mL) is added dropwise to a refluxing suspension of activated zinc dust (2.1 g) in THF (25 mL) and refluxing
30 continued for a further 4 hours. The reaction is allowed to cool to ambient temperature, filtered through hyflo and concentrated in vacuo to give a yellow solid. The solid is taken up in ethyl acetate (50 mL) and washed with brine which leads to the formation of a white precipitate which is filtered off then partitioned between ethyl acetate (50 mL) and 1 N HCl (50 mL). The organic
35 phase is washed with water, dried over magnesium sulphate, filtered and concentrated in vacuo to give (RS)-2,2-difluoro-3-hydroxy-3-(2-methyl-phenyl)propanoic acid as a yellow solid (2.4 g)

REFERENCE EXAMPLE 67

A solution of ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2.91 g) and diethyl oxalate (2.82 g) in dry toluene are stirred at room temperature whilst adding sodium hydride (0.28 g, 60% dispersion in mineral oil) portionwise. The reaction mixture is heated at reflux for 5 hours then left at room temperature for 18 hours. Evaporation gives an orange oil which is dissolved in ethanol (100 mL) and treated with glacial acetic acid (1.5 mL). Hydroxylamine hydrochloride (0.49 g) is added and the mixture heated at reflux for 5 hours. The reaction mixture is partitioned between ethyl acetate (100 mL) and water (100 mL). The aqueous phase is extracted with three times ethyl acetate (50 mL). The combined organic phases are washed with brine (50 mL), with water (50 mL), dried over magnesium sulphate and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-[2-(5-ethoxycarbonylisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2.56 g) as an orange oil.

REFERENCE EXAMPLE 68

A solution of diisopropyl azodicarboxylate (40 g) in anhydrous THF (20 mL) is added dropwise with stirring under ice cooling to a solution of triphenylphosphine (52 g) in anhydrous THF (200 mL). The resulting thick white suspension is treated dropwise over about 1 h under ice cooling with a mixture of 2-acetyl-5-(3-pyridylmethoxy)phenol (23 g) and *tert*-butyl (R)-4-hydroxy-4-(2-methyl)butanoate (30 g) in anhydrous THF (100 mL). After stirring under ice cooling for 2 hours and then standing at room temperature overnight the mixture is partitioned between ethyl acetate (500 mL) and water (500 mL). The layers are separated and the organic layer washed with water, dried, and evaporated. The residue is partially purified by flash chromatography on silica eluting with a mixture of ethyl acetate and cyclohexane (60 to 100%). Fractions containing the required product are combined and evaporated to give impure *tert*-butyl (R)-4-[2-(acetyl-5-(3-pyridylmethoxy)phenoxy)-4-methylphenyl]butanoate (72 g). This material is

mixed with dimethyl oxalate (17.7 g) in anhydrous toluene (200 mL) and heated at reflux under a Dean-Stark water separator. When no more water separates the mixture is allowed to cool and sodium hydride (4.8 g, 60% dispersion) is added. The mixture is gradually returned to reflux; when a vigorous reaction sets in the heat is removed. When this reaction subsides the mixture is stirred at reflux for a further 15 minutes and then evaporated to dryness. The residue is dissolved in methanol (200 mL), treated with hydrazine hydrate (5 g) and taken to pH 6 with glacial acetic acid (about 7 g). The mixture is refluxed for 1 hour, further hydrazine hydrate (2 g) is added, and reflux continued for another 1 hour. The residue obtained after evaporation is partitioned between ethyl acetate (500 mL) and water (500 mL). The layers are separated and the organic layer washed with water, dried, and evaporated. The residue is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and cyclohexane (3:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give tert-Butyl (R)-4-[2-(5-methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate (20.4 g).

REFERENCE EXAMPLE 69

20

(R)-t-butyl 4-[2-(5-methoxycarbonyl-3-pyrazolyl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate (2 g) is dissolved in trifluoroacetic acid (10 mL) and allowed to stand at room temperature for 2 minutes. The solution is quenched with saturated aqueous sodium hydrogen carbonate solution (100 mL) and the product extracted into ethyl acetate (100 mL). This solution is washed with water, dried, and evaporated. The residue is purified by flash chromatography eluting with 60% ethyl acetate in cyclohexane to remove trace high running impurities, and then 20% methanol in dichloromethane to obtain the required product. Crystallisation from ethyl acetate/cyclohexane mixture gives (R)-4-[2-(5-methoxycarbonyl-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid (0.4 g) of a white solid, mp 155-6°C.

30

REFERENCE EXAMPLE 70

35

2-Hydroxy-4-(3-thienylmethoxy)benzonitrile (2.5 g) is added portionwise to a stirred suspension of sodium hydride (0.44 g, 60% dispersion in mineral oil) in dry dimethyl formamide (30 mL). After stirring at ambient temperature for

30 minutes, ethyl (RS)-4-chloro-4-(2-cyanophenyl)butanoate (2.87 g) is added in one portion and the reaction stirred for 16 hours at 100°C. The reaction is concentrated in vacuo and the residue partitioned between ethyl acetate (100 mL) and water (100 mL). The organic phase is dried over magnesium sulphate, filtered and concentrated in vacuo. The residual oil is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and pentane (1:2 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl (RS)-4-(2-cyanophenyl)-4-[2-cyano-5-(3-thienylmethoxy)phenoxy]butanoate (250 mg) as an orange oil.

10 A solution of 2-cyanobenzaldehyde (5 g) in dichloromethane (10 mL) is added dropwise to a stirred 1M solution of titanium (IV) chloride in dichloromethane (38 mL) at 0°C under nitrogen. The mixture is stirred for 5 minutes and [(1-ethoxycyclopropyloxy)trimethyl]silane (8.42 mL) is added dropwise maintaining the temperature at 0-5°C. The mixture is stirred for 16 hours at ambient temperature then quenched with water (200 mL). The mixture is extracted three times with dichloromethane (100 mL) and the combined organic extracts are washed with brine (100 mL), dried over magnesium sulphate and evaporated to give ethyl (RS)-4-chloro-4-(2-cyanophenyl)-butanoate (5.5 g) as a yellow oil.

REFERENCE EXAMPLE 71

Diisopropyl azodicarboxylate (1.19 g) is added to a stirred solution of triphenylphosphine (1.55 g) in tetrahydrofuran (30 mL) at 0°C under nitrogen and stirring is continued at 0°C for 30 minutes. A solution of isothiazol-4-ylmethanol (0.34 g) and 2-fluoro-4-hydroxybenzonitrile (0.4 g) in tetrahydrofuran (30 mL) is added at such a rate that the temperature did not exceed 0°C. After the addition is complete stirring is continued at 0°C for 30 minutes then at ambient temperature for 24 hours. The reaction mixture is concentrated in vacuo and the residue is partitioned between ethyl acetate (50 mL) and water (50 mL). The organic phase is washed with water (50 mL), with brine (50 mL), dried over magnesium sulphate and evaporated. The residue is purified by flash chromatography on silica eluting with dichloromethane. Fractions homogeneous in the required product are combined and evaporated to give 2-fluoro-4-(isothiazol-4-ylmethoxy)benzonitrile (0.24 g), m.p. 102-105°C.

Isothiazole-4-carboxylic acid (0.65 g) is added portionwise to a stirred mixture of lithium aluminum hydride (0.19 g) in tetrahydrofuran (50 mL) at ambient temperature under nitrogen the reaction mixture is heated at reflux for
5 4 hours then cooled to 0°C. Sodium hydroxide solution (3% w/v) is added dropwise and the mixture stirred at 0°C for 1 hour then filtered through hyflo. The filtrate is evaporated and azeotroped with toluene to give isothiazol-4-ylmethanol (0.35 g).

10 REFERENCE EXAMPLE 72

A stirred solution of 1-(3-benzyloxyphenyl)-2-phenylethanone in tetrahydrofuran (30 mL) is treated with a solution of lithium diisopropylamine in tetrahydrofuran (5.5 mL, 2 M) at -35°C under nitrogen. The resulting clear
15 yellow solution is cooled to -70°C and treated with a solution of bromine (1.6 g) in dichloromethane (5 mL). Stirring is continued at -70°C and treated with a solution of bromine (1.6 g) in dichloromethane (5 mL). Stirring is continued at -70°C for 5 minutes. Water (30 mL) is added and the reaction mixture allowed to warm to room temperature. The reaction mixture is extracted twice with ether
20 (50 mL), the combined extracts are washed with brine (40 mL), dried over magnesium sulphate and evaporated to give (RS)-1-(3-benzyloxyphenyl)-2-bromo-2-phenylethanone (3.92 g) in the form of a yellow oil.

Wilkinson's catalyst (18 mg) is added to a stirred solution of ethyl
25 3-(2,4-dibenzyloxy)phenyl propen-2-oate (1.52 g) in toluene (10 mL) at ambient temperature under nitrogen. Neat triethylsilane (1.6 mL) is added and the reaction stirred at ambient temperature for 16 hours. The reaction is concentrated in vacuo to give a dark brown oil which is purified by flash chromatography on silica eluting with a mixture of ethyl acetate and petrol
30 ether (5:1 v/v). Fractions homogeneous in the required product are combined and evaporated to give ethyl 3-(2,4-dibenzyloxy)phenyl propanoate (1.48 g) as a colourless oil.

Sodium hydride (1.32 g, 60% dispersion in mineral oil) is suspended in
35 THF (75 mL) and cooled to -10°C. Neat triethylphosphonoacetate (7.4 g) is added dropwise and stirred at -10°C for 15 minutes. A solution of 2,4-dibenzyl-oxybenzaldehyde (10 g) is added dropwise to the reaction. A thick viscous oil

is formed and the reaction is left at 0°C for one hour then quenched with saturated ammonium chloride solution (100 mL) and extracted with ethyl acetate (3 x 100 mL). The ethyl acetate extracts are combined, washed with water (2 x 100 mL), dried over magnesium sulphate, filtered and concentrated in vacuo to give ethyl 3-(2,4-dibenzyloxy)phenylpropen-2-oate (12 g) as a yellow oil.

REFERENCE EXAMPLE 74

10 A mixture of 2,4-dibenzyloxyphenol (3.1 g) and sodium hydride (0.44 g, 60% dispersion in mineral oil) in THF (75 mL) is refluxed for 30 minutes then refluxing is allowed to abate. Neat ethyl bromoacetate (1.8 g) is added and refluxing continued for 1 hour. The reaction is concentrated in vacuo and the residue partitioned between ethyl acetate (100 mL) and water (100 mL). The
15 organic phase is washed with water (100 mL), dried over magnesium sulphate, filtered and concentrated in vacuo to give a light brown oil. Flash chromatography of the residue eluting with dichloromethane affords ethyl 2,4-dibenzyloxyphenoxyacetate (3.2 g) after trituration with pentane.

20 REFERENCE EXAMPLE 75

A stirred solution of methyl 4-[2-hydroxy-4-(thiophen-3-ylmethoxy)-phenyl]-4-oxobutanoate (1.6 g) in DMF (20 mL) is treated with 60% sodium hydride (0.2 g) and stirred at 25° for 0.3 hours. The mixture is treated with ethyl
25 (R,S)-4-chloro-4-(2-methylphenyl)butanoate (1.2 g) and heated at 80°C for 6 hours. The solution is evaporated (hivac). The residue is dissolved in ethyl acetate, washed with water, dried and evaporated. The residue is purified by flash chromatography on silica, eluting with ethyl acetate/pentane : 1/1 to give (R,S)-4-[2-(3-methoxycarbonylpropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-
30 (2-methylphenyl)butyric acid, ethyl ester (1.9 g), an orange oil.

A stirred solution of 4-(2,4-dihydroxyphenyl)-4-oxobutyric acid, methyl ester 6.76 g in methyl ethyl ketone (150 mL) is treated with potassium carbonate (4.17 g), potassium iodide (5 g) tetrabutyl ammonium chloride (1 g)
35 and 3-chloromethylthiophene (4 g), and refluxed for 24 hours. The reaction mixture is filtered, the filtrate is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried, and evaporated. The residue is

recrystallised from ethyl acetate/cyclohexane to give 4-[2-hydroxy-4-(thiophen-3-ylmethoxy)phenoxy]-4-oxo-butyric acid, methyl ester (7.16 g), a pale cream solid, mp. 100-102°C.

5 REFERENCE EXAMPLE 76

A stirred solution of (R,S)-4-[2-(amino-hydroxyiminomethyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid, ethyl ester (400 mg) in trimethyl orthoformate (8 mL) is refluxed for 2 hours. The solution is
10 evaporated. The residue is purified by flash chromatography on silica, eluting with pentane/ether: 2/1 to give (R,S)-4-[2-(1,2,4-oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid, ethyl ester (74 mg).

A solution of ethyl (R,S)-4-[2-cyano-5-(thiophen-3-ylmethoxy)phenoxy]-
15 4-(2-methylphenyl)butanoate (2.3 g) and hydroxylamine hydrochloride (0.4 g) in ethanol (80 mL) is treated with a solution of NaOH (0.23 g) in water (8 mL) and refluxed for 60 hours. The solution is evaporated. The residue is dissolved in ethyl acetate, washed with water, dried and evaporated. The residual yellow oil is purified by flash chromatography on silica, eluting with
20 pentane/ethyl acetate: 3/1 to give ethyl (R,S)-4-[2-(amino-hydroxyiminomethyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (0.63 g).

REFERENCE EXAMPLE 77

25 A stirred solution of ethyl (R,S)-4-[2-(2-thiocarbamoyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (1.17 g) in ethanol (12 mL) and DMF (6 mL) is treated with 50% aq. chloroacetaldehyde (1.17 g) and refluxed for 24 hours. The solution is evaporated. The residue is partitioned between ethyl acetate and water. The organic layer is washed with water, dried
30 and evaporated. The residual solid is purified by flash chromatography on silica eluting with ethyl acetate/pentane: 1/3, followed by trituration with ether to give (R,S)-4-[2-(thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid, ethyl ester, a colourless solid (1.02 g), m.p. 127-129°C.

35 A stirred solution of ethyl (R,S)-4-[2-(2-carbamoyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate (2.44 g) in THF (49 mL) is treated with Lawesson's reagent (1.08 g) and maintained at 25°C for 24 hours.

The solution is evaporated. The residual gum is purified by flash chromatography on silica, eluting with pentane/ethyl acetate: 2/1, followed by trituration with ether to give (R,S)-4-[2-(2-thiocarbamoyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid, ethyl ester, a pale yellow solid (1.79 g), mp. 134-135°C.

Compounds of formula I and their pharmaceutically acceptable salts exhibit pharmacological activity and accordingly are of use for the preparation of pharmaceutical compositions for the treatment of humans and other animals.

10

Compounds of formula I exhibit useful pharmacological activity and accordingly are incorporated into pharmaceutical compositions and used in the treatment of patients suffering from certain medical disorders. More especially, they are endothelin inhibitors, in particular endothelin A inhibitors.

15

The present invention provides compounds of formula I, and compositions containing compounds of formula I, which are of use in a method for the treatment of a patient suffering from, or subject to, conditions which can be ameliorated by the administration of an inhibitor of endothelin. For example, compounds within the present invention are useful for the treatment of diseases and conditions characterized by, or having an etiology involving pathogenic endothelin levels. Examples of disease states and conditions which can be ameliorated by the administration of inhibitors of endothelin such as compounds of formula I include vascular ischaemia, for example cerebrovascular disease including cerebral ischaemia such as stroke and subarachnoid hemorrhage, coronary disorders such as myocardial infarction including acute myocardial infarction, coronary heart disease, angina including unstable and vasospastic angina, preeclampsia, essential and pulmonary hypertension and congestive heart failure, renal disorders such as acute renal insufficiency and chronic renal insufficiency, cyclosporin induced nephrotoxicity, erythropoietin induced renal complications and hypertension, gastrointestinal disorders such as ulceration and irritable bowel syndrome, poor peripheral skeletal muscle disorders such as peripheral vascular disease, intermittent claudication and critical limb ischaemia, glaucoma, atherosclerosis and related diseases, hypertension, asthma, migraine, endotoxin shock, Raynauds disease, benign prostatic hyperplasia, bone loss such as

35

osteoporosis and restenosis after angioplasty. Compounds of the present invention are also useful as a therapy for promoting wound healing.

According to a further feature of the invention there is provided a method
5 for the treatment of a human or animal patient suffering from, or subject to, conditions which can be ameliorated by the administration of an inhibitor of endothelin, especially ETA, for example conditions as hereinbefore described, which comprises the administration to the patient of an effective amount of compound of formula I or a composition containing a compound of formula I.
10 "Effective amount" is meant to describe an amount of compound of the present invention effective in inhibiting endothelin and thus producing the desired therapeutic effect.

The present invention also includes within its scope pharmaceutical
15 formulations which comprise at least one of the compounds of formula I in association with a pharmaceutically acceptable carrier or coating.

In practice compounds of the present invention may generally be administered parenterally, rectally or orally. The compounds of the invention
20 may also be administered topically to treat peripheral vascular diseases.

The products according to the invention may be presented in forms permitting administration by the most suitable route and the invention also relates to pharmaceutical compositions containing at least one product
25 according to the invention which are suitable for use in human or veterinary medicine. These compositions may be prepared according to the customary methods, using one or more pharmaceutically acceptable adjuvants or excipients. The adjuvants comprise, inter alia, diluents, sterile aqueous media and the various non-toxic organic solvents. The compositions may be
30 presented in the form of tablets, pills, granules, powders, aqueous solutions or suspensions, injectable solutions, elixirs, creams, ointments or syrups, and can contain one or more agents chosen from the group comprising sweeteners, flavorings, colorings, or stabilizers in order to obtain pharmaceutically acceptable preparations.

35

The choice of vehicle and the content of active substance in the vehicle are generally determined in accordance with the solubility and chemical

properties of the product, the particular mode of administration and the provisions to be observed in pharmaceutical practice. For example, excipients such as lactose, sodium citrate, calcium carbonate, dicalcium phosphate and disintegrating agents such as starch, alginic acids and certain complex
5 silicates combined with lubricants such as magnesium stearate, sodium lauryl sulfate and talc may be used for preparing tablets. To prepare a capsule, it is advantageous to use lactose and high molecular weight polyethylene glycols. When aqueous suspensions are used they can contain emulsifying agents or agents which facilitate suspension. Diluents such as sucrose, ethanol,
10 polyethylene glycol, propylene glycol, glycerol and chloroform or mixtures thereof may also be used.

For parenteral administration, emulsions, suspensions or solutions of the products according to the invention in vegetable oil, for example sesame
15 oil, groundnut oil or olive oil, or aqueous-organic solutions such as water and propylene glycol, injectable organic esters such as ethyl oleate, as well as sterile aqueous solutions of the pharmaceutically acceptable salts, are used. The solutions of the salts of the products according to the invention are especially useful for administration by intramuscular or subcutaneous injection.
20 The aqueous solutions, also comprising solutions of the salts in pure distilled water, may be used for intravenous administration with the proviso that their pH is suitably adjusted, that they are judiciously buffered and rendered isotonic with a sufficient quantity of glucose or sodium chloride and that they are sterilized by heating, irradiation or microfiltration.

25 Suitable compositions containing the compounds of the invention may be prepared by conventional means. For example, compounds of the invention may be dissolved or suspended in a suitable carrier for use in a nebulizer or a suspension or solution aerosol, or may be absorbed or
30 adsorbed onto a suitable solid carrier for use in a dry powder inhaler.

Solid compositions for rectal administration include suppositories formulated in accordance with known methods and containing at least one compound of formula I.

35 Compositions for topical administration include creams and ointments formulated in accordance with known methods, such as a topical carrier such

as Plastibase® (mineral oil gelled with polyethylene) and containing at least one compound of formula I.

5 The percentage of active ingredient in the compositions of the invention may be varied, it being necessary that it should constitute a proportion such that a suitable dosage shall be obtained. Obviously, several unit dosage forms may be administered at about the same time. The dose employed will be determined by the physician, and depends upon the desired therapeutic effect, the route of administration and the duration of the treatment, and the condition
10 of the patient. In the adult, the doses are generally from about 0.01 to about 100 mg/kg, preferably about 0.02 to about 50 mg/kg, and more preferably about 0.1 to about 25 mg/kg body weight (or from about 1 to about 5000 mg, preferably from about 5 to about 2000 mg) in single or 2 to 4 divided daily doses. In each particular case, the doses will be determined in accordance
15 with the factors distinctive to the subject to be treated, such as age, weight, general state of health and other characteristics which can influence the efficacy of the medicinal product.

The products according to the invention may be administered as
20 frequently as necessary in order to obtain the desired therapeutic effect. Some patients may respond rapidly to a higher or lower dose and may find much weaker maintenance doses adequate. For other patients, it may be necessary to have long-term treatments at the rate of 1 to 4 doses per day, in accordance with the physiological requirements of each particular patient. Generally, the
25 active product may be administered orally 1 to 4 times per day. It goes without saying that, for other patients, it will be necessary to prescribe not more than one or two doses per day.

The compounds of the present invention can also be administered in
30 combination with endothelin converting enzyme inhibitors, angiotensin II receptor antagonists, renin inhibitors, angiotensin converting enzyme inhibitors, α - and β -adrenoceptor agonists and antagonists, diuretics, potassium channel activators, calcium channel antagonists, nitrates, antiarrhythmic agents, positive inotropic agents, serotonin receptor agonists
35 and antagonists, platelet activating factor antagonists, histamine receptor antagonists, proton pump inhibitors, antithrombotic and thrombolytic agents, lipid lowering agents, antibiotic agents and phosphodiesterase inhibitors.

If formulated as a fixed dose, such combination products employ the compounds of the present invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. The compounds of the present invention may also be formulated with or useful
5 in conjunction with antifungal and immunosuppressive agents such as amphotericin B, cyclosporins and the like to counteract the hypertension and nephrotoxicity secondary to such compounds. The compounds of the present invention may also be used in conjunction with haemodialysis.

10 Compounds within the scope of the present invention exhibit marked pharmacological activities according to tests described in the literature which tests results are believed to correlate to pharmacological activity in humans and other mammals. The following pharmacological test results are typical characteristics of compounds of the present invention.

15

IN VITRO TESTS

A) Preparation of ET_A receptors:

20 A10 cells are grown to confluence in Dulbecco's modified essential medium containing 10% foetal calf serum. Two days after the final medium change cells are harvested by scraping from the base of the flask and centrifuged at 1500 rpm for 10 minutes at 4°C in a bench centrifuge. The resulting pellets are washed in 50 mM Hepes buffer pH 7.3 containing calcium
25 chloride (1 mM) and magnesium chloride (5 mM) and resuspended at a density of 140,000 cells/mL in the same. Cell suspensions are then frozen using a mixture of methanol and solid carbon dioxide and stored at -20°C until required. For use in the assay cells are diluted to the required density with Hepes buffer pH 7.3.

30

B) Preparation of ET_B receptors:

Rats are killed by cervical dislocation and the cerebellum tissue is
35 removed into ice cold Tris buffer pH 7.4 containing sucrose (0.25 M), ethylenediaminetetracetic acid (3 mM), and a cocktail of protease inhibitors. After homogenizing using a glass/teflon manual homogenizer, the samples are

centrifuged at 4°C for 17 minutes at 1000 g, and the resulting supernatants are retained. This material is centrifuged at 4000 g for 35 minutes at 4°C and the pellets are resuspended in 50 mM Tris buffer pH 7.4, and the protein concentration is measured. Aliquots of 100 mL are frozen in a mixture of
5 methanol and solid carbon dioxide and stored at -20°C until required. For use in the assay samples are diluted to the required concentration with Tris buffer pH 7.4 containing 0.1% bovine serum albumin.

C) Assay methodology:

10

Assays are performed using Millipore 96 well filtration plates with 0.22 µm filters in a final volume of 250 µL. Mixtures consisting of test compound and [¹²⁵I]-ET-1 (20 pM) in Tris buffer pH 7.4 containing 0.1% bovine serum albumin are treated with either A10 cells or cerebellum protein.
15 Total and non-specific binding are measured in the absence and presence of unlabelled ET-1 (100 nM). Approximately 60,000 A10 cells are used per well or 5 µg of cerebellum protein. Plates are incubated for 2 hours at 37°C before the reaction is terminated by vacuum filtration. Plates are washed twice with assay buffer at 4°C and the filters are punched out for gamma counting.

20

RESULTS:

The IC₅₀ values (50% inhibition of ET-1 binding) for compounds of the invention in the antagonism of ET-1 at ET_A and ET_B receptors are from about
25 10⁻⁹ M to about 10⁻⁴ M.


IN VIVO TESTS

Compounds are evaluated *in vivo* after administration either
30 intravenously or orally to pithed rats (prepared according to the method described by Gillespie, J.S. & Muir, T.C., (1967), Br. J. Pharmacol. Chemother., 30, 78-87). The degree of inhibition produced by the test compounds of the biphasic depressor and pressor responses to ET-1 given intravenously is measured.

35

WHAT IS CLAIMED IS:

1. A compound of formula I



5

wherein

10 R^1 is hydrogen, $-(\text{lower alkyl})_q(\text{CO}_2R^6 \text{ or OH})$, $-\text{CN}$, $-\text{C}(R^7)=\text{NOR}^8$, $-\text{NO}_2$, $-\text{O}(\text{lower alkyl})R^9$, $-\text{C}\equiv\text{C}-R^{10}$, $-\text{CR}^{11}=\text{C}(R^{12})(R^{13})$, $-\text{C}(=\text{O})\text{CH}_2\text{C}(=\text{O})\text{CO}_2\text{H}$, $-\text{CO}(R^{14})$, alkylthio, alkylsulphinyl, alkylsulphonyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, substituted thiocarbamoyl, sulphamoyl or an optionally substituted nitrogen-containing ring;

15 R^2 is aryl lower alkoxy, heteroaryl lower alkoxy, aryl lower alkylthio or heteroaryl lower alkylthio;

20 R^3 is hydroxy, alkoxy, aryloxy, cycloalkyl(lower alkyl)oxy, cycloalkenyl(lower alkyl)oxy, aryl lower alkoxy, heteroaryl lower alkoxy, aryl lower alkylthio, heteroaryl lower alkylthio or aralkynyl;

R^4 is $-\text{Y}-\text{CH}(R^{15})(\text{alkyl or alkenyl})R^{16}$;

25 R^5 is alkyl, alkenyl or halo;

R^6 , R^7 , R^{17} , R^{18} and 19 are independently hydrogen or alkyl;

R^8 is hydrogen, aralkyl or $-(\text{lower alkyl})\text{CO}_2R^{17}$;

30 R^9 is $-\text{CN}$, $-\text{CO}_2R^{19}$, $-\text{CH}_2\text{OH}$, or carbamoyl;

R^{10} is $-\text{CO}_2\text{H}$ or carboxyphenyl;

R^{11} is hydrogen, alkyl or aralkyl;

R¹² and R¹³ are independently hydrogen, -CO₂R¹⁸, -CN, aryl lower alkyl, heteroaryl lower alkyl or -NHC(=O)aryl, provided that one of R¹² and R¹³ is -CO₂H;

5

R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, aralkenyl, heteroaralkenyl or carboxy;

R¹⁵ is aryl or heteroaryl;

10

R¹⁶ is carboxy or acid isostere;

Y is oxygen or carbonyl; and

15

m, n, o, p and q are independently zero or 1,

with the provisos that (i) when o is zero then m and n are both 1 (ii) when o is 1 then at least one of m and n is 1 (iii) when o is zero then R¹ cannot represent hydrogen,

20

or a pharmaceutically acceptable salt, N-oxide or prodrug thereof.

2. A compound of claim 1 wherein

25

R¹ is hydrogen, -(lower alkyl)_q(CO₂H or OH), -CN, -C(R⁷)=NOR⁸, -NO₂, -O(lower alkyl)R⁹, -C≡C-R¹⁰, -CR¹¹=C(R¹²)(R¹³), -C(=O)CH₂C(=O)CO₂H, -CO(R¹⁴), alkylthio, alkylsulphinyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted tetrazolyl, 1,2,4-oxadiazolyl, substituted isoxazolyl, pyrazolyl, substituted pyrazolyl, pyridyl, isothiazolyl, oxazolyl, substituted oxazolyl or dihydrooxazolyl;

30

R² is aryl lower alkoxy or heteroaryl lower alkoxy;

R⁴ is -O-CH(aryl)(alkyl)R¹⁶;

35

R⁵ is halo;

R^8 is hydrogen or $-(\text{lower alkyl})\text{CO}_2\text{H}$;

R^9 is $-\text{CN}$, $-\text{CO}_2\text{H}$ or carbamoyl;

5 R^{10} is $-\text{CO}_2\text{H}$;

R^{12} and R^{13} are independently hydrogen, $-\text{CO}_2\text{H}$, $-\text{CN}$ or $-\text{NHC}(=\text{O})\text{aryl}$, provided that one of R^{12} and R^{13} is $-\text{CO}_2\text{H}$;

10 R^{14} is hydrogen, alkyl, $-(\text{lower alkyl})\text{carboxy}$, heteroarylalkenyl or carboxy; and

m , o and p are 1 and n is zero.

15 3. A compound of claim 1 wherein

R^1 is hydrogen, $-(\text{lower alkyl})_q(\text{CO}_2\text{H or OH})$, $-\text{CN}$, $-\text{C}(\text{R}^7)=\text{NOR}^8$, $-\text{NO}_2$, $-\text{O}(\text{lower alkyl})\text{R}^9$, $-\text{C}\equiv\text{C}-\text{R}^{10}$, $-\text{CR}^{11}=\text{C}(\text{R}^{12})(\text{R}^{13})$, $-\text{C}(=\text{O})\text{CH}_2\text{C}(=\text{O})\text{CO}_2\text{H}$, $-\text{CO}(\text{R}^{14})$, alkylthio, alkylsulphinyl, carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted tetrazolyl, 1,2,4-oxadiazolyl, substituted isoxazolyl, pyrazolyl, substituted pyrazolyl, pyridyl, oxazolyl, substituted oxazolyl or dihydrooxazolyl;

25 R^2 is aryl lower alkoxy or heteroaryl lower alkoxy;

R^4 is $-\text{O}-\text{CH}(\text{aryl})(\text{alkyl})\text{R}^{16}$;

R^8 is hydrogen, aralkyl or $-(\text{lower alkyl})\text{CO}_2\text{H}$

30 R^9 is $-\text{CN}$, $-\text{CO}_2\text{H}$, $-\text{CH}_2\text{OH}$, or carbamoyl;

R^{10} is $-\text{CO}_2\text{H}$;

R^{12} and R^{13} are independently hydrogen, $-\text{CO}_2\text{H}$, $-\text{CN}$, or $-\text{NHC}(=\text{O})\text{aryl}$, provided that one of R^{12} and R^{13} is $-\text{CO}_2\text{H}$;

R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, heteroarylalkenyl or carboxy; and

m and o are 1 and n and p are zero.

5

4. A compound of claim 1 wherein

R¹ is -(lower alkyl)_q(CO₂H or OH), -CN, -C(R⁷)=NOR⁸, -NO₂, -O(lower alkyl)R⁹, -C≡C-R¹⁰, -CR¹¹=C(R¹²)(R¹³), -C(=O)CH₂C(=O)CO₂H, -CO(R¹⁴),
10 carbamoyl, thiocarbamoyl, substituted carbamoyl, tetrazolyl, substituted tetrazolyl, substituted isoxazolyl, substituted pyrazolyl or substituted oxazolyl;

R² is aryl lower alkoxy or heteroaryl lower alkoxy;

15

R³ is hydroxy, alkoxy, aryl lower alkoxy, heteroaryl lower alkoxy or aralkynyl;

R⁸ is hydrogen, aralkyl or -(lower alkyl)CO₂H;

20

R⁹ is -CN, -CO₂H, -CH₂OH, or carbamoyl;

R¹⁰ is -CO₂H;

R¹² and R¹³ are independently hydrogen, -CO₂H, -CN, aryl lower alkyl,
25 heteroaryl lower alkyl or -NHC(=O)aryl, provided that one of R¹² and R¹³ is -CO₂H;

R¹⁴ is hydrogen, alkyl, -(lower alkyl)carboxy, heteroarylalkenyl or carboxy; and

30

m and n are 1 and o and p are zero.

5. A compound according to any one of claims 1, 2 and 3 wherein

35

o is 1; and

R^1 is $-\text{CO}_2\text{H}$, $-\text{C}\equiv\text{CCO}_2\text{H}$, $-\text{C}(\text{R}^{11})=\text{CHCO}_2\text{H}$, $-(\text{CH}_2)_2\text{CO}_2\text{H}$, $-\text{O}(\text{lower alkyl})\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}(\text{lower alkyl})\text{CO}_2\text{H}$, (carboxy)pyrazolyl, (carboxy)(lower alkyl)pyrazolyl, (alkoxycarbonyl)pyrazolyl, $-\text{CN}$, $-\text{NO}_2$, (carboxy)oxazolyl, (carboxy)isoxazolyl, pyrazolyl, formyl, oxazolyl, 1,2,4-oxadiazolyl, acetyl, 5 carboxymethoxyimino, dihydrooxazolyl, pyridyl, carboxamide, thiocarboxamide, pyridylpyrazolyl, cyano(lower alkyl)carboxamide, carboxamidopyrazolyl, isothiazolyl or oxime, wherein R^{11} is hydrogen or lower alkyl.

10 6. A compound of claim 5 wherein

R^1 is $-\text{CN}$, NO_2 , substituted pyrazolyl, substituted oxazolyl or substituted isoxazolyl.

15 7. A compound according to claim 1 or claim 4 wherein

o is zero; and

20 R^1 is $-\text{CO}_2\text{H}$, $-\text{C}\equiv\text{CCO}_2\text{H}$, $-\text{C}(\text{R}^{11})=\text{CHCO}_2\text{H}$, $-\text{C}(=\text{O})\text{NH}(\text{lower alkyl})\text{CO}_2\text{H}$, (carboxy)pyrazolyl, (carboxy)(lower alkyl)pyrazolyl or (carboxy)oxazolyl.

8. A compound of claim 7 wherein

25 R^1 is $-\text{C}(=\text{O})\text{NH}(\text{lower alkyl})\text{CO}_2\text{H}$, (carboxy)pyrazolyl or (carboxy)(lower alkyl)pyrazolyl.

9. A compound according to any relevant preceding claim wherein

30 o is zero; and

R^2 and R^3 are substituted on the phenyl moiety at the (2- and 4-) positions relative to R^1 .

35 10. A compound according to any relevant preceding claim wherein

o is 1; and

R² is substituted on the phenyl moiety at the 4- position relative to R¹
and R⁴ is substituted on the phenyl moiety at the 2-position relative to R¹.

5 11. A compound according to any relevant preceding claim wherein
p is 1; and

R⁵ is substituted on the phenyl moiety at the ortho position relative to R¹.
10

12. A compound according to any relevant preceding claim wherein
o is 1; and

15 R¹⁵ is aryl.

13. A compound of claim 12 wherein R¹⁵ is phenyl or phenyl substituted at
the ortho position relative to the attachment of the phenyl group to the rest of
the R⁴ moiety, and is optionally further substituted.
20

14. A compound of claim 12 wherein R¹⁵ is phenyl substituted at the ortho
position relative to the attachment of the phenyl group to the rest of the R⁴
moiety by methyl, and is optionally further substituted.

25 15. A compound of claims 12 to 14 wherein R¹⁵ is phenyl substituted by
one or more substituents selected from lower alkyl, halo, CF₃, CN or lower
alkoxy.

30 16. A compound according to any relevant preceding claim wherein

R² and R³ are independently aryl lower alkoxy or heteroaryl lower
alkoxy.

35 17. A compound according to any relevant preceding claim wherein

o is zero;

R² is 1-(aryl)ethoxy; and

R³ is benzyloxy, (3-thienyl)methoxy, (3-pyridyl)methoxy or (4-isothiazolyl)methoxy.

5

18. A compound of claim 17 wherein R² is 1-(o-tolyl)ethoxy.

19. A compound according to any relevant preceding claim wherein

10

o is 1; and

R² is heteroaryl lower alkoxy.

20. A compound of claim 19 wherein R² is (3-thienyl)methoxy,
15 (3-pyridyl)methoxy or (4-isothiazolyl)methoxy.

21. A compound according to any relevant preceding claim wherein

20

o is 1; and

Y is oxygen.

22. A compound according to any relevant preceding claim wherein

25

o is 1; and

n is zero.

23. A compound according to any relevant preceding claim wherein

30


o is 1; and

R⁴ is -Y-CH(R¹⁵)(C₁₋₃alkyl)R¹⁶

35 24. A compound according to claim 23 wherein

R⁴ is carboxy.

25. A compound of formula Ia



5

wherein R¹, R² and R⁴ are as defined in claim 1, or a physiologically acceptable salt, N-oxide or prodrug thereof.

26. A compound according to claim 25 wherein R¹ is CN, NO₂, substituted
10 pyrazolyl, substituted oxazolyl or substituted isoxazolyl.

27. A compound according to claim 25 or claim 26 wherein R¹ is CN.

28. A compound according to any one of claims 25 to 27 wherein R² is
15 heteroarylmethoxy.


29. A compound according to claim 28 wherein R² is pyridylmethoxy, thienylmethoxy or isothiazolylmethoxy.

20 30. A compound according to any one of claims 25 to 29 wherein R⁴ is -OCH(R¹⁵)(alkyl)R¹⁶ wherein R¹⁵ is aryl and R¹⁶ is carboxy.

31. A compound according to claim 30 wherein R¹⁵ is phenyl substituted in
25 the ortho position relative to the attachment of the phenyl group to the rest of the R⁴ moiety by a lower alkyl or chloro substituent and is optionally further substituted by one or more of halo, lower alkyl, CF₃, CN or lower alkoxy.

32. A compound according to claim 30 or claim 31 wherein within
-OCH(R¹⁵)(alkyl)R¹⁶ the alkyl portion is -CH₂CH₂-.
30

33. A compound of formula Ib



wherein R² and R¹⁵ are as defined in claim 1, or a physiologically acceptable salt, N-oxide or prodrug thereof.

5

34. A compound according to claim 33 wherein R² is heteroarylmethoxy and R¹⁵ is aryl.

10

35. A compound according to claim 34 wherein R¹⁵ is phenyl substituted in the ortho position by lower alkyl or chloro and is optionally further substituted by one or more of halo, lower alkyl, CN, CF₃ and lower alkoxy.

36. A compound according to any of claims 33 to 35 wherein R² is pyridylmethoxy, thienylmethoxy or isothiazolylmethoxy.

15

37. A compound according to claim 1 which is:

20

2-Benzyloxy-4-(4-chlorobenzyloxy)benzoic acid;

2-Benzyloxy-4-(3-phenylpropyloxy)benzoic acid;

2,4-Di-(4-chlorobenzyloxy)benzoic acid;

2-Benzyloxy-4-(2-(3-indolyl)ethoxy)benzoic acid;

2-Hydroxy-4-benzyloxybenzoic acid;

2-(3-Phenylpropyloxy)-4-benzyloxybenzoic acid;

2-Benzyloxy-4-(2-naphthylmethoxy)benzoic acid;

25

2-Benzyloxy-4-(1-naphthylmethoxy)benzoic acid;

2-Benzyloxy-4-(3,4-methylenedioxybenzyloxy)benzoic acid;

2-(2-Pyridylmethoxy)-4-benzyloxybenzoic acid;

2-(2-Phenylethoxy)-4-benzyloxybenzoic acid;

2-Cyclohexylmethoxy-4-benzyloxybenzoic acid;

30

2-(4-Chlorobenzyloxy)-4-benzyloxybenzoic acid;

2-Benzyloxy-4-(2-phenylethoxy)benzoic acid;

2-Benzyloxy-4-(2-methylpropoxy)benzoic acid;

2-Benzyloxy-4-(4-nitrobenzyloxy)benzoic acid;

- 2-Benzyloxy-4-(4-fluorobenzyloxy)benzoic acid;
Benzyloxy-2-(2-methylpropoxy)benzoic acid;
2-(4-Pyridylmethoxy)-4-benzyloxybenzoic acid;
2-Benzyloxy-4-(3,4-dichlorobenzyloxy)benzoic acid;
5 2-Benzyloxy-4-(4-methoxybenzyloxy)benzoic acid;
2-Benzyloxy-4-(3-methoxybenzyloxy)benzoic acid;
2,4-Dibenzyloxybenzoic acid;
(E)-3-(2,4-Dibenzyloxyphenyl)prop-2-enoic acid;
3-(2,4-Dibenzoyloxyphenyl)prop-2-ynoic acid;
10 N-(2,4-Dibenzyloxybenzoyl)glycine;
N-(4-Benzyloxy-2-(1-o-tolyloethoxy)benzoyl)glycine;
3-(2,4-Dibenzyloxyphenyl)-pyrazole-5-carboxylic acid;
2-Benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid;
1-Benzyl-5-(4-benzyloxyphenyl)-2H-pyrazole-3-carboxylic acid;
15 5-(2,4-Dibenzyloxyphenyl)-2H-tetrazole;
(Z)-2-Benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid;
(E)-2-Benzoylamino-3-(2,4-dibenzyloxyphenyl)acrylic acid;
E-2-Benzyl-3-(2,4-dibenzyloxyphenyl)acrylic acid;
(E)-3-(2,4-Dibenzyloxyphenyl)-2-cyanoacrylic acid;
20 3-(2,4-Dibenzyloxyphenyl)isoxazole-5-carboxylic acid;
4-(2,4-Dibenzyloxyphenyl)-2,4-dioxobutanoic acid;
2,4-Dibenzyloxybenzaldehyde oxime;
2,4-Dibenzyloxybenzaldehyde;
2,4-Dibenzyloxybenzyl alcohol;
25 2,4-Dibenzyloxybenzonitrile;
2,4-Dibenzyloxynitrobenzene;
2,4-Dibenzyloxyacetophenone;
2,4-Dibenzyloxyphenol;
2,4-Dibenzyloxyacetophenone oxime;
30 2-Benzyloxy-4-phenylethynylbenzoic acid;
2-(2,4-Dibenzyloxyphenoxy)ethanol hemihydrate;
2-(2,4-Dibenzyloxyphenoxy)acetamide;
(RS)-5-(3-Benzyloxyphenyl)-5-oxo-4-phenylpentanoic acid;
(RS)-5-(3-Benzyloxyphenyl)-4-(2-chlorophenyl)-5-oxopentanoic acid;
35 (RS)-5-(3-Benzyloxyphenyl)-4-(2-methoxyphenyl)-5-oxopentanoic acid;
(RS)-5-(3-Benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
(RS)-5-(3-Benzyloxyphenyl)-4-(2-ethylphenyl)-5-oxopentanoic acid;

- (RS)-5-(3-Benzyloxyphenyl)-4-(2-chloro-6-fluorophenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxyphenyl)-4-(2,6-dichlorophenyl)-5-oxopentanoic acid;
- 5 (RS)-5-(3-Benzyloxyphenyl)-4-(4-methoxyphenyl)-5-oxopentanoic acid;
- (RS)-5-(3-benzyloxyphenyl)-4-(3,4-dichlorophenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxyphenyl)-4-(4-chlorophenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxyphenyl)-4-(4-methylphenyl)-5-oxopentanoic acid;
- 10 (RS)-5-(3-Benzyloxyphenyl)-4-(3-chlorophenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxyphenyl)-4-(3-methoxyphenyl)-5-oxopentanoic acid;
- (RS)-5-(4-Benzyloxyphenyl)-5-oxo-4-phenylpentanoic acid;
- (RS)-4-(3-Benzyloxyphenyl)-4-oxo-3-phenylbutanoic acid;
- (RS)-6-(3-Benzyloxyphenyl)-6-oxo-5-phenylhexanoic acid;
- 15 (RS)-7-(3-Benzyloxyphenyl)-7-oxo-6-phenylheptanoic acid;
- (RS)-6-(3-Benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoic acid;
- (RS)-5-(3-Benzyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoic acid;
- (3RS,4RS)-5-(3-Benzyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoic acid;
- 20 (2RS,4RS)-5-(3-Benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoic acid;
- (RS)-5-(3-Methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(2-Methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid;
- (RS)-5-[3-(3,4-Methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoic acid;
- 25 (RS)-4-(2-Methylphenyl)-5-[3-(2-methylpropoxy)phenyl]-5-oxopentanoic acid;
- (RS)-5-[3-(4-Chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 30 (RS)-5-(3-Cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(3-Thienylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(2-Fluorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 35 (RS)-5-[3-(2-Furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;

- (RS)-5-[3-(3-Furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(3-Chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 5 (RS)-5-[3-(4-Methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(2-pyridylmethoxy)phenyl]pentanoic acid;
- (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]pentanoic acid;
- 10 acid;
- (RS)-5-(2-Methyl-3-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-(3,5-Dibenzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 15 (RS)-5-[3-(3-Methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(3-Aminobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- (RS)-5-[3-(3-Isothiazolylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;
- 20 oxopentanoic acid;
- (RS)-5-(3-benzyloxy-5-hydroxyphenyl)-4-(2,3-dimethylphenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 25 (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(4-pyridylmethoxy)phenyl]pentanoic acid;
- (RS)-4-(2-Methylphenyl)-5-[3-(3-methyl-2-thienylmethoxy)phenyl]-5-oxopentanoic acid;
- (RS)-4-(2-Methylphenyl)-5-oxo-5-[3-(3-pyridylmethoxy)phenyl]pentanoic acid;
- 30 acid;
- (RS)-5-(3-Benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoic acid;
- (RS)-5-(3-Benzyloxy-6-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 35 (RS)-5-(3-Benzyloxyphenyl)-4-(3-methylpyrid-4-yl)-5-oxopentanoic acid;
- (RS)-5-[3-(1,2,4-Oxadiazol-3-ylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoic acid;

- Methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylpentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chlorophenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methoxyphenyl)-5-oxopentanoate;
5 Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-ethylphenyl)-5-oxopentanoate;
10 Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2-chloro-6-fluorophenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(2,6-dichlorophenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3,4-dichlorophenyl)-5-oxopentanoate;
15 Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(4-chlorophenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(4-methylphenyl)-5-oxopentanoate;
20 Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3-chlorophenyl)-5-oxopentanoate,
Methyl (RS)-5-[3-(benzyloxy)phenyl]-4-(3-methoxyphenyl)-5-oxopentanoate;
Methyl (RS)-5-(4-benzyloxyphenyl)-5-oxo-4-phenylpentanoate;
25 Methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-(2-pyridyl)pentanoate,
Methyl (3RS,4RS)-5-(3-benzyloxyphenyl)-3-methyl-5-oxo-4-phenylpentanoate;
Methyl (2RS,4RS)-5-(3-benzyloxyphenyl)-2-methyl-5-oxo-4-phenylpentanoate;
30 Methyl (RS)-5-[3-(2-methoxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate;
Methyl (RS)-5-[3-(3,4-methylenedioxybenzyloxy)phenyl]-5-oxo-4-phenylpentanoate;
Methyl (RS)-5-(3-benzyloxy-2-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
35 Methyl (RS)-5-(3,5-benzyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;

- Methyl (RS)-5-(3-benzyloxy-4-methylphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- Methyl (RS)-5-(3-benzyloxyphenyl)-4-(2,5-dimethylphenyl)-5-oxopentanoate;
- 5 Methyl (RS)-5-(3-benzyloxyphenyl)-5-oxo-4-phenylbutanoate;
Ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhexanoate;
Ethyl (RS)-7-(3-benzyloxyphenyl)-7-oxo-6-phenylheptanoate;
Ethyl (RS)-6-(3-benzyloxyphenyl)-6-oxo-5-phenylhex-2-enoate;
Methyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 10 Ethyl (RS)-5-[3-(4-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-(3-cyclopentylmethoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- 15 Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(3-thienylmethoxy)phenyl]pentanoate;
Methyl (RS)-5-[3-(2-fluorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(2-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 20 Methyl (RS)-5-[3-(3-furylmethoxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(3-chlorobenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 25 Methyl (RS)-5-[3-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-pyridylmethoxy)phenyl]pentanoate;
Methyl (RS)-4-(2-methylphenyl)-5-oxo-5-[3-(2-thienylmethoxy)phenyl]pentanoate;
- 30 Methyl (RS)-5-[3-(3-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(3-isothiazolylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- 35 Methyl (RS)-5-[3-(4-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;

- Methyl (RS)-5-[3-(3-nitrobenzyloxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
Methyl (RS)-5-[3-(3-pyridylmethoxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- 5 Methyl (RS)-5-(3-methoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-(3-isopropoxyphenyl)-4-(2-methylphenyl)-5-oxopentanoate;
- Methyl (RS)-5-[3-(1,2,4-oxadiazol-3-yl-methoxy)phenyl]-4-(2-methylphenyl)-5-oxo-pentanoate;
- 10 Methyl (RS)-5-[3-(3-methyl-2-thienylmethoxy)-phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
Methyl (RS)-5-[3-(3-aminobenzyloxy)phenyl]-5-oxo-4-(2-methylphenyl)pentanoate;
- Ethyl (RS)-5-[3-benzyloxy-5-(4-methoxybenzyloxy)phenyl]-4-(2-methylphenyl)-5-oxopentanoate;
- 15 (RS)-5-(3-Benzyloxy-5-hydroxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
(RS)-5-[4-(3-Benzyloxyphenyl)-4-oxo-3-phenylbutyl]-1H-tetrazole;
(RS)-5-[3-(3-Benzyloxyphenyl)-3-oxo-2-(2-methylphenyl)propyl]-1H-tetrazole;
- 20 (RS)-4-(3-Benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylsulphonic acid;
(RS)-4-[3-(3-Thienylmethoxy)phenyl]-3-(2-methylphenyl)-4-oxobutylsulphonic acid;
- 25 (RS)-6-(3-Benzyloxyphenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide;
(RS)-6-(3-(3-Thienylmethoxy)phenyl)-5-(2-methylphenyl)-3,4-dihydro-1,2-oxathiine-2,2-dioxide;
- Ethyl (RS)-4-(3-benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate;
- 30 Diethyl (RS)-4-(3-benzyloxyphenyl)-3-(2-methylphenyl)-4-oxobutylphosphonate;
(RS)-4-(3-Benzyloxyphenoxy)-4-phenylbutanoic acid;
(RS)-4-(2-Carboxy-5-benzyloxyphenoxy)-4-phenylbutanoic acid;
- 35 (RS)-4-[2-(N-Ethylcarbamoyl)-5-benzyloxy-phenoxy]-4-phenylbutanoic acid;
(RS)-4-(2-Acetyl-5-benzyloxyphenoxy)-4-phenylbutanoic acid;

- (RS)-4-[3-(3-Thienylmethoxy)phenoxy]-4-phenylbutanoic acid;
(RS)-4-(3-Benzoyloxy)phenoxy-4-(2-methylphenyl)butanoic acid;
(RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-
butanoic acid;
- 5 (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-phenylbutanoic acid;
(RS)-4-[2-Acetyl-5-(3-pyridylmethoxy)-phenoxy]-4-phenylbutanoic acid;
(RS)-4-[2-Carbamoyl-5-benzoyloxyphenoxy]-4-phenylbutanoic acid;
(RS)-4-[2-Cyano-5-benzoyloxyphenoxy]-4-phenylbutanoic acid;
(RS)-4-[2-(N,N-Dimethylcarbamoyl)-5-benzoyloxyphenoxy]-4-
- 10 phenylbutanoic acid;
(RS)-4-[2-(3-Pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-
methylphenyl)butanoic acid;
(RS)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-
methylphenyl)butanoic acid;
- 15 Ethyl (RS)-4-(3-benzoyloxyphenoxy)-4-phenylbutanoate;
Ethyl (RS)-4-(2-methoxycarbonyl-5-benzoyloxyphenoxy)-4-
phenylbutanoate;
Ethyl (RS)-4-[(3-benzoyloxy)phenoxy]-4-(2-methylphenyl)butanoate;
Ethyl (RS)-4-[3-(3-thienylmethoxy)phenoxy]-4-phenylbutanoate;
- 20 Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-
phenylbutanoate;
Ethyl (RS)-4-[2-(N-ethylcarbamoyl)-5-benzoyloxyphenoxy]-4-
phenylbutanoate;
Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-
- 25 methylphenyl)butanoate;
Ethyl (RS)-4-[2-acetyl-5-(3-pyridylmethoxy)-phenoxy]-4-
phenylbutanoate;
Ethyl (RS)-4-[2-carbamoyl-5-benzoyloxyphenoxy]-4-phenylbutanoate;
Ethyl (RS)-4-[2-N,N-dimethylcarbamoyl-5-benzoyloxyphenoxy]-4-
- 30 phenylbutanoate;
Ethyl (RS)-4-[2-carbamoyl-5-(3-thienyl-methoxy)phenoxy]-4-(2-
methylphenyl)butanoate
Ethyl (RS)-4-(2-acetyl-5-benzoyloxyphenoxy)-4-phenylbutanoate;
Ethyl (RS)-4-(2-cyano-5-benzoyloxyphenoxy)-4-phenylbutanoate;
- 35 Ethyl (RS)-4-[2-cyano-5-(3-thienylmethoxy)-phenoxy]-4-(2-
methylphenyl)butanoate;

- Ethyl (RS)-4-[2-(3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- (R)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- (S)-5-(3-Benzoyloxyphenyl)-4-(2-methylphenyl)-5-oxopentanoic acid;
- 5 (R)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid,
- (S)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid.
- (RS)-4-[2-Acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 10 (RS)-4-(2-Methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- 15 (RS)-4-(2-Methylphenyl)-4-[2-thiocarbamoyl-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- (E)-(RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)prop-2-enoyl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- Ethyl (RS)-4-[2-acetyl-5-(5-pyrimidinylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate,
- 20 Ethyl (RS)-4-(2-methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoate;
- Ethyl (RS)-4-(2-methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]butanoate;
- 25 Ethyl (RS)-4-(2-methylphenyl)-4-[5-(3-thienylmethoxy)-2-thiocarbamoylphenoxy]butanoate;
- (RS)-4-[5-Benzoyloxy-2-(methylthio)phenoxy]-4-phenylbutanoic acid;
- (E)-(RS)-4-[5-Benzoyloxy-2-(2-carboxyethenyl)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 30 (RS)-4-[2-(1-Methyl-2-carboxyethenyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-[5-Benzoyloxy-2-hydroxyiminomethylphenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(Carboxymethoxy)iminomethyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 35 (RS, RS)-4-[2-(1-Hydroxyethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;

- (RS)-4-(2-Methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)-phenoxy]butanoic acid;
- (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 5 (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)-butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]-
- 10 butanoic acid;
- (RR,RS,SR,SS)-4-(2-Methylphenyl)-4-[2-methylsulphinyl-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoic acid;
- 15 (RS)-4-(5-Benzoyloxy-2-formylphenoxy)-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-Formyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[5-(3-thienylmethoxy)-2-(trifluoroacetyl)-
- 20 phenoxy]butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)-phenoxy]butanoic acid;
- (RS)-4-[2-Cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 25 (RS)-4-[2-Carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(3-Imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(2-Carboxyethyl)carbamoyl}-5-(3-thienylmethoxy)-
- 30 phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(Carboxymethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-[2-(N-(2-Cyanoethyl)carbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid ;
- 35 (RS)-4-[2-(5-Carboxy-1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;

- (RS)-4-[2-{N-(Carbamoylmethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(Methoxycarbonylmethyl)-carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 5 Ethyl (RS)-4-[5-benzyloxy-2-(methylthio)phenoxy]-4-phenylbutanoate;
- Ethyl (RS)-4-(2-methylphenyl)-4-[2-methylthio-5-(3-thienylmethoxy)-phenoxy]butanoate;
- Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoate;
- 10 Ethyl (RS)-4-(5-benzyloxy-2-formylphenoxy)-4-(2-methylphenyl)-butanoate;
- Ethyl (RS)-4-[2-formyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- Ethyl (RS)-4-[5-(3-thienylmethoxy)-2-trifluoroacetylphenoxy]-4-(2-methylphenyl)butanoate;
- 15 Ethyl (RS)-4-[2-carbamoyl-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoate;
- Ethyl (RS)-4-[2-acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-bromophenyl)-butanoate;
- 20 Ethyl (RS)-4-(2-methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]-butanoate;
- Ethyl (RS)-4-[2-{N-methoxycarbonylmethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- Ethyl (RS)-4-[2-{N-(3-imidazol-1-ylpropyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- 25 Ethyl (RS)-4-[2-{N-(2-cyanoethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- Ethyl (RS)-4-[2-{N-cyanomethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- 30 Ethyl (RS)-4-[2-{N-(2-methoxycarbonylethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- Ethyl (RS, RS)-4-(2-methylphenyl)-4-[2-methylsulphinyl-5-(3-thienylmethoxy)phenoxy]butanoate;
- Ethyl (E)-(RS)-4-(2-methylphenyl)-4-[2-(2-methoxycarbonylethenyl)-5-benzyloxyphenoxy]-butanoate;
- 35 Ethyl (RS)-4-[2-(2-methoxycarbonyl-1-methylethenyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;

- Ethyl (RS)-4-(5-benzyloxy-2-hydroxyiminomethylphenoxy)-4-(2-methylphenyl)butanoate;
- Ethyl (RS)-4-[2-hydroxyiminomethyl-5-(3-thienylmethoxy))phenoxy]-4-(2-methylphenyl)-butanoate;
- 5 (RS)-4-[2-{N-(ethoxycarbonylmethoxy)iminomethyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- Ethyl (RS,RS)-4-[2-(1-hydroxyethyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoate;
- Ethyl (RS)-4-(2-methylphenyl)-4-[2-(propen-2-yl)-5-(3-thienylmethoxy)-phenoxy]butanoate;
- 10 Ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-pyridylmethoxy)-phenoxy]-4-(2-ethylphenyl)butanoate;
- Ethyl (RS)-4-[2-(5-ethoxycarbonyl-3-pyrazolyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoate;
- 15 Ethyl (RS)-4-(2-methylphenyl)-4-[2-pentafluoroethyl-5-(3-thienylmethoxy)phenoxy]-butanoate;
- Ethyl (RS)-4-[2-cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate.
- Methyl (RS)-4-[2-(5-methoxycarbonyl-1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- 20 (RS)-N-[4-{2-Cyano-5-(3-thienylmethoxy)-phenoxy}-4-(2-methylphenyl)butanoyl]benzenesulphonamide;
- (R)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 25 (S)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (R)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- Dicyclohexylammonium (R)-4-[2-cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoate;
- 30 (S)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-5-(3-Benzylthiophenyl)-4-(2-methylphenyl)-5-oxopentanoic acid hemihydrate;
- 35 (RS)-4-(2-Methylphenyl)-5-oxo-(1-oxopyrid-3-ylmethoxy)pentanoic acid hemihydrate;

- (RS)-4-[2-(Benzyloxyiminomethyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid hydrate;
- (RS)-4-(2-Carboxycarbonyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoic acid;
- 5 (R)-4-[2-Benzoyl-5-(pyridin-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid;
- (RS)-3-[3-(3-Carboxy-1-(2-methylphenyl)propoxy)-4-nitrophenoxy-methyl]benzoic acid;
- (RS)-4-{5-[3-(2-Carboxyethyl)benzyloxy]-2-cyano-phenoxy}-4-(2-methylphenyl)butanoic acid;
- 10 (RS)-4-[2-(4,5-Dihydrooxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-((1R)-1-Carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid ;
- 15 (RS)-4-[2-((1S)-1-carboxy-2-phenylethylcarbamoyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-(Oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-2-[2-(3-Carboxy-1-(2-methylphenyl)propoxy)-4-(3-thienyl-methoxy)benzoylamino]acrylic acid;
- 20 (RS)-4-[2-(4-Carboxyoxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-(2-Chloro-6-fluorophenyl)-4-(2-cyano-5-(3-thienylmethoxy)-phenoxy)butanoic acid;
- 25 (R)-4-(2-Methylphenyl)-4-(2-nitro-5-(3-pyridylmethoxy)phenoxy)butanoic acid;
- (RS)-4-(2-Cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoic acid;
- (RS)-4-(Benzo[1,3]dioxol-4-yl)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]butanoic acid;
- 30 (RS)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2,3-dimethylphenyl)-butanoic acid;
- (RS)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(5-hydroxy-2-methylphenyl)-butanoic acid;
- 35 (RS)-5-[2-Cyano-5-(3-thienylmethoxy)-phenoxy]-5-(2-methylphenyl)-pentanoic acid;

- (RS)-4-(2-Cyano-3-fluoro-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl) butanoic acid;
- (RS)-2,4-Dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)-phenyl]butanoic acid;
- 5 (RS)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-(5-Carboxypyrazol-3-yl)-5-(pyrid-3-ylmethoxy)phenoxy]-4-(2-methylphenyl) butanoic acid;
- 10 (RS)-3-(2-Cyano-5-(pyridylmethoxy)phenoxy)-2,2-difluoro-3-(2-methylphenyl)-propanoic acid;
- (RS)-4-[2-(5-Carboxisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 15 (R)-4-[2-(5-Methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-(5-Carbamoylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[(2-Acetyl-5-(3-thienylmethoxy)phenoxy)]-4-(2-cyanophenyl)-butanoic acid;
- 20 (R)-4-[(2-Cyano-5-(isothiazol-4-ylmethoxy)phenoxy)]-4-(2-methylphenyl)-butanoic acid;
- (R)-4-[2-Nitro-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid;
- 25 (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}methane sulphonamide;
- (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}trifluoromethane sulphonamide;
- (R,S)-4-[2-(3-Carboxypropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid;
- 30 (R,S)-4-[2-(1,2,4-Oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid; or
- (R,S)-4-[2-(Thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid,
- 35 or a physiologically acceptable salt, N-oxide or prodrug thereof.

38. A compound according to claim 1 which is:

- (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-[2-(3-Pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 5 (RS)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-(2-pyridyl)-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- 10 (RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)pyrazol-5-yl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-thiocarbamoyl-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- (E)-(RS)-4-(2-Methylphenyl)-4-[2-{3-(2-pyridyl)prop-2-enoyl}-5-(3-thienylmethoxy)phenoxy]butanoic acid;
- 15 (RS)-4-[2-{N-(Carboxymethoxy)iminomethyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 20 (RS)-4-[2-(5-Carboxy-3-pyrazolyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-(2-Methylphenyl)-4-[2-nitro-5-(3-thienylmethoxy)phenoxy]-butanoic acid;
- (RS)-4-[2-Acetyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-chloro-6-fluorophenyl)butanoic acid;
- 25 (RS)-4-[2-Formyl-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (RS)-4-[2-Cyano-3-methyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 30 (RS)-4-[2-Carbamoyl-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(2-Carboxyethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(Carboxymethyl)carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 35 (RS)-4-[2-(N-(2-Cyanoethyl)carbamoyl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid ;

- (RS)-4-[2-(5-Carboxy-1-methyl-3-pyrazolyl)-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-{N-(Carbamoylmethyl)carbamoyl}-5-(3-thienylmethoxy)-phenoxy]-4-(2-methylphenyl)butanoic acid;
- 5 (RS)-4-[2-{N-(Methoxycarbonylmethyl)-carbamoyl}-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-Cyano-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- (R)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2-
- 10 methylphenyl)butanoic acid;
- (RS)-4-(2-Carboxycarbonyl-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl)butanoic acid;
- (RS)-4-[2-(4,5-Dihydrooxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-
- 15 methylphenyl)butanoic acid;
- (RS)-4-[2-(Oxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (RS)-2-[2-(3-Carboxy-1-(2-methylphenyl)propoxy)-4-(3-thienylmethoxy)benzoylamino]acrylic acid;
- (RS)-4-[2-(4-Carboxyoxazol-2-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-
- 20 methylphenyl)butanoic acid;
- (RS)-4-(2-Chloro-6-fluorophenyl)-4-(2-cyano-5-(3-thienylmethoxy)-phenoxy)butanoic acid;
- (R)-4-(2-Methylphenyl)-4-(2-nitro-5-(3-pyridylmethoxy)phenoxy)butanoic acid;
- 25 (RS)-4-(2-Cyano-5-(3-thienylmethoxy)phenoxy)-4-(2,5-dimethylphenyl)butanoic acid;
- (RS)-4-(Benzo[1,3]dioxol-4-yl)-4-[2-cyano-5-(3-pyridylmethoxy)-phenoxy]butanoic acid;
- (RS)-4-[2-Cyano-5-(3-pyridylmethoxy)phenoxy]-4-(2,3-dimethylphenyl)-
- 30 butanoic acid;
- (RS)-5-[2-Cyano-5-(3-thienylmethoxy)-phenoxy]-5-(2-methylphenyl)-pentanoic acid;
- (RS)-4-(2-Cyano-3-fluoro-5-(3-thienylmethoxy)phenoxy)-4-(2-methylphenyl) butanoic acid;
- 35 (RS)-2,4-Dioxo-4-[2-(1-(2-methylphenyl)ethoxy)-4-(3-thienylmethoxy)-phenyl]butanoic acid;

- (RS)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-(5-Ethoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 5 (R)-4-[2-(5-Carboxypyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl) butanoic acid;
- (RS)-4-[2-(5-Carboxyisoxazol-3-yl)-5-(3-thienylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[2-(5-Methoxycarbonylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- 10 (R)-4-[2-(5-Carbamoylpyrazol-3-yl)-5-(3-pyridylmethoxy)phenoxy]-4-(2-methylphenyl)butanoic acid;
- (R)-4-[(2-Cyano-5-(isothiazol-4-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butanoic acid;
- 15 (R)-4-[2-Nitro-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)-butyric acid;
- (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}methane sulphonamide;
- (R,S)-N-{4-[2-Cyano-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyryl}trifluoromethane sulphonamide;
- 20 (R,S)-4-[2-(3-Carboxypropionyl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid;
- (R,S)-4-[2-(1,2,4-Oxadiazol-3-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid; or
- 25 (R,S)-4-[2-(Thiazol-2-yl)-5-(thiophen-3-ylmethoxy)phenoxy]-4-(2-methylphenyl)butyric acid,
- or a physiologically acceptable salt, N-oxide or prodrug thereof.
39. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier or excipient.
40. A pharmaceutical composition for use in the treatment of a disease state associated with physiological detrimental amounts of endothelin comprising an endothelin inhibitory effective amount of the compound according to claim 1.
41. A compound according to claim 1 for use in therapy.

42. The use of a compound according to claim 1 for the manufacture of a medicament for the treatment of a disease state associated with physiological detrimental amounts of endothelin.
- 5 43. A method for treating a disease state capable of being modulated by inhibiting endothelin comprising administering to a patient suffering from said disease state an effective amount of a compound of claim 1.
- 10 44. The method of claim 43 wherein the disease state is selected from the group consisting of vascular ischaemia, coronary disorders, coronary heart disease, angina, preeclampsia, essential hypertension, pulmonary hypertension, congestive heart failure, renal disorders, cyclosporin induced nephrotoxicity, erythropoietin induced renal complications, erythropoietin induced hypertension, gastrointestinal disorders, poor peripheral skeletal
15 muscle disorders, intermittent claudication, critical limb ischaemia, glaucoma, atherosclerosis, atherosclerotic related diseases, hypertension, asthma, migraine, endotoxin shock, Raynauds disease, benign prostatic hyperplasia and restenosis after angioplasty.
- 20 45. A method for promoting wound healing capable of being modulated by inhibiting endothelin comprising administering to a patient suffering from said wound an effective amount of a compound of claim 1.
- 25 46. A process for the preparation of a compound according to claim 1 substantially as hereinbefore described.
47. A compound substantially as hereinbefore described with reference to the Examples.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 94/02499

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07C65/24 C07C65/26 C07C65/28 C07C59/68 C07C59/90

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 051 819 (F.HOFFMANN-LA ROCHE & CO.) 19 May 1982 see page 24, line 20 - page 25, line 4 ---	1,4,7
X	EP,A,0 072 926 (MERCK PATENT GESELLSCHAFT) 2 March 1983 see page 41, line 18 ---	1,4,7
X	EP,A,0 419 905 (EISAI CO.) 3 April 1991 see claim 1 ---	1,4
A	EP,A,0 248 594 (ORTHO PHARMACEUTICAL CO.) 9 December 1987 see page 19, line 1 - line 35 ---	1,39,41
A	EP,A,0 068 250 (BAYER AG) 5 January 1983 see claims 1,5 ---	1,39,41
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

8 March 1995

Date of mailing of the international search report

15. 03. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Klag, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 94/02499

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,2 792 418 (DRUEY ET AL.) 14 May 1957 see column 1, line 37 - line 42 see claim 1 ---	1,39,41
A	US,A,3 914 166 (RUDOLPH ET AL.) 21 October 1975 see claim 1 -----	1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 94/02499

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0051819	19-05-82	AT-T- 11766	15-02-85
		AU-B- 543311	18-04-85
		AU-A- 7730581	20-05-82
		CA-A- 1206964	01-07-86
		JP-A- 57122039	29-07-82
EP-A-0072926	02-03-83	DE-A- 3132754	03-03-83
		DE-A- 3139064	14-04-83
		AU-B- 557714	08-01-87
		AU-A- 8669282	24-02-83
		CA-A- 1207791	15-07-86
		CA-A- 1294277	14-01-92
		JP-C- 1723132	24-12-92
		JP-B- 4006713	06-02-92
		JP-A- 58039681	08-03-83
		US-A- 4616090	07-10-86
		US-A- 5104881	14-04-92
EP-A-0419905	03-04-91	AU-B- 637138	20-05-93
		AU-A- 6225890	14-03-91
		CA-A- 2024479	12-03-91
		CN-A- 1050182	27-03-91
		JP-A- 3188042	16-08-91
		NO-B- 174292	03-01-94
		US-A- 5385942	31-01-95
		US-A- 5210239	11-05-93
EP-A-0248594	09-12-87	US-A- 4826868	02-05-89
		AU-B- 596844	17-05-90
		AU-A- 7360887	10-12-87
		DE-D- 3788244	05-01-94
		DE-T- 3788244	14-04-94
		ES-T- 2059377	16-11-94
		US-A- 5164381	17-11-92
EP-A-0068250	05-01-83	JP-A- 63022080	29-01-88
		DE-A- 3125059	05-01-83
		JP-A- 58000928	06-01-83
		US-A- 4590291	20-05-86

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 94/02499

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-2792418	14-05-57	NONE	
US-A-3914166	21-10-75	AT-A, B 305631	15-01-73
		AT-A- 294053	15-10-71
		BE-A- 741343	06-05-70
		DE-A- 1807301	04-06-70
		FR-A- 2022723	07-08-70
		GB-A- 1229082	21-04-71
		NL-A- 6916372	11-05-70
		SE-B- 360350	24-09-73
		US-A- 3915823	28-10-75
		US-A- 3891524	24-06-75
		US-A- 3824284	16-07-74

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB94/02499

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.: Claims searched incompletely: 1-23; 25-42, 46, 47
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
As the drafting of the claims is not clear and concise and encompasses such enormous amount of products; a complete search is not possible on economic grounds. So the search has been based on the examples.
Claims not searched 24 (contradiction to other claims.)
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.